Non-Hodgkin's lymphoma and occupation in Sweden: a registry based analysis

Martha S Linet, Hans S R Malker, Joseph K McLaughlin,
Jan A Weiner, William J Blot, Jan L E Ericsson, Joseph F Fraumeni Jr

Abstract

Incidence of non-Hodgkin's lymphoma in different employment categories was evaluated from the Swedish Cancer-Environment Registry, which links cancer incidence during 1961 to 1979 with occupational information from the 1960 census. New associations were found for men employed in shoemaking and shoe repair, porcelain and earthenware industries, education, and other white collar occupations. Several findings supported associations found in other countries, including excesses among woodworkers, furniture makers, electric power plant workers, farmers, dairy workers, lorry drivers, and other land transport workers. Risks were not increased among chemists, chemical or rubber manufacturing workers, or petrochemical refinery workers. Caution must be used in drawing causal inferences from these linked registry data because information on exposure and duration of employment is not available. Nevertheless, this study has suggested new clues to possible occupational determinants of non-Hodgkin's lymphoma.

(British Journal of Industrial Medicine 1993;50:79-84)

During the past few decades, incidence of non-Hodgkin's lymphoma has been increasing in many countries including Sweden, particularly among the elderly.¹⁻⁴ Although some of this increase may reflect diagnostic improvements, environmental and occupational factors have also been implicated.⁵ The relation of occupational exposures to non-Hodgkin's lymphoma has not been extensively evaluated,

although excess risks have been reported among farmers in several geographical areas.⁶⁻¹¹ Among the agricultural exposures implicated in non-Hodgkin's lymphoma are pesticides, fertilisers, and livestock.^{7 12-15} Not all studies have found an increased occurrence among persons in agriculture related occupations or those exposed to pesticides, however.¹⁶⁻¹⁹ Workers exposed to organic solvents such as chemical manufacturing workers,^{12 13 17 22 23} chemists,²⁴ and petrochemical workers,^{17 25} may also have an increased incidence.^{20 21} To evaluate a possible role for these occupations in the aetiology of non-Hodgkin's lymphoma, we utilised the Swedish Cancer-Environment Registry (CER), a national linked registry of cancer and employment data. Because earlier reports²⁶⁻²⁸ have examined agricultural and forestry workers with this linked registry, our focus is on employment outside these areas.

Materials and methods

The CER merges incidence reports from the National Swedish Cancer Registry (NSCR) for the period 1961 to 1979 with occupational information from the 1960 census.²⁹ The unique 10 digit number assigned to each Swedish citizen allows linkage of these two data sources for individual subjects. According to Swedish law, all malignant tumours and certain precancerous conditions and benign lesions must be reported to the NSCR.³⁰

The CER employs the International Classification of Diseases (ICD) 7th revision for coding of site and a modified World Health Organisation coding system for histopathological classification. During the past 25 years, numerous morphology based classification schemes have been proposed for non-Hodgkin's lymphoma and at present no single system is in widespread use.³¹ The problems related to reproducibility for a given pathologist or in attaining agreement among pathologists, using either a single classification or different schemes, have been documented.³² These difficulties in conjunction with the failure of the ICD coding system to reflect the rapid changes in non-Hodgkin's lymphoma classification over time make it difficult to assess occupational association by subtype over a long

Epidemiology and Biostatistics Program, Division of Cancer Etiology, National Cancer Institute, Rockville, Maryland

M S Linet, J K McLaughlin, W J Blot, J F Fraumeni
National Board of Occupational Safety and Health, Solna, Sweden

H S R Malker, J A Weiner
National Board of Health and Welfare, Stockholm, Sweden

J L E Ericsson

period of follow up. Thus in this analysis non-Hodgkin's lymphoma will be considered as a single entity.

The standardised cumulative incidence ratio (SIR) was used to estimate the risk of non-Hodgkin's lymphoma for each occupational and industrial group. The SIR is the ratio of observed to expected number of cases in a specific employment category. The expected number was calculated by applying five year birth cohort, sex, and region specific non-Hodgkin's lymphoma incidence rates in the Swedish male population for 1961-79 to the birth cohort and sex specific distributions of the industrial and occupational categories. To evaluate statistical significance it was assumed that the observed number of cases of non-Hodgkin's lymphoma followed a Poisson distribution.³³

The employment classifications, derived from international standards, and used in the 1960 census were taken to define industrial and occupational groups.^{34,35} The SIRs were calculated for all major (one digit code), general (two digit code), and specific (three digit code) industrial and occupational categories.

Results

There were 4496 employed Swedish men diagnosed with non-Hodgkin's lymphoma (all microscopically verified) during the 19 year follow up (1961 to 1979).

The SIRs for non-Hodgkin's lymphoma were initially evaluated by the 10 major (one digit code) industry and occupational categories. Incidence was

not significantly increased or decreased in any of the major industrial or occupational employment categories (data not shown).

There were few significantly increased SIRs among the 46 general (two digit) industries, or among the 60 general (two digit) occupations. For general industries, moderately increased risks were seen in the furniture and furnishings industry (SIR 1.3; 55 cases; $p < 0.05$) and in the electrical power industry (SIR 1.3; 81 cases; $p < 0.05$). For the general (two digit) occupation categories, non-Hodgkin's lymphoma was moderately increased among men working in education (SIR 1.3; 86 cases; $p < 0.05$).

Table 1 presents results for specific (three digit) industries and occupations with statistically significant ($p < 0.05$) SIRs. Workers in the dairy, shoe repair, and porcelain and earthenware industries had significantly increased SIRs for non-Hodgkin's lymphoma, as did those in the wholesale paper industry, banking, research and scientific institutes, and in other public jobs. The significant excess of non-Hodgkin's lymphoma at the two digit level for the furniture and furnishings industry was not seen at the three digit level, although furniture making was still increased (Industrial code (Ind) 250; SIR = 1.3; 44 cases) as was other related furniture work (Ind 252; SIR = 1.9; four cases). The significant excess for the electrical power industry at the two digit level was not seen at the three digit level, although the SIR for electrical power plants was still increased (Ind 500; SIR = 1.3; 52 cases), as were SIRs for gas plants (Ind 501; SIR = 2.0; eight cases) and waterworks

Table 1 Statistically significant ($p < 0.05$) SIRs for non-Hodgkin's lymphomas by specific industries and occupations among Swedish men, 1961-79

Code		Cases	Number expected	SIR†
<i>Industry</i>				
201	Dairy	25	15.2	1.6*
232	Shoe repair	23	12.6	1.8*
332	Porcelain and earthenware	14	7.2	1.9*
616	Wholesale paper	16	7.9	2.0*
670	Banking	25	14.1	1.8**
830	Research and scientific institute	13	3.7	3.5**
837	Other public jobs	10	3.7	2.7**
<i>Specific Occupation</i>				
005	Engineers, technicians in mining	20	10.3	1.9*
050	School teachers	14	6.8	2.1*
052	Special teachers	24	14.8	1.6*
203	Bank cashier	5	1.5	3.3*
290	Secretaries, typists	11	5.4	2.1*
297	Health insurance clerks	6	1.9	3.2**
721	Shoemakers	23	13.4	1.7*
875	Lorry drivers	42	29.2	1.4*

* $p < 0.05$; ** $p < 0.01$.

†Adjusted for age and region.

(Ind 502; SIR = 1.8; nine cases). For specific occupations (table 1, lower panel), significantly raised SIRs were seen for mining engineers and technicians, school and special teachers, bank cashiers, secretaries, health insurance clerks, shoemakers, and lorry drivers.

To further examine non-Hodgkin's lymphoma and employment, we evaluated risks for non-Hodgkin's lymphoma for occupations within industries. Table 2 lists the occupations within industry having significantly high SIRs. Of interest and not already identified in table 1 were the increased risks for engineers and technicians employed in power plants, mechanical engineers and technicians in car factories, and other woodworkers in sawmill and planing mills.

Discussion

This CER survey for non-Hodgkin's lymphoma has uncovered new associations among shoemakers and workers in the porcelain and earthenware industry. There have been some links between shoe workers and leukaemia, aplastic anaemia,^{36,37} and non-Hodgkin's lymphoma (two cases)³⁸ in clinical reports, but no excess of non-Hodgkin's lymphoma has been found in analytical studies in the United States,³⁹⁻⁴¹ the United Kingdom,⁴² or Italy.⁴³ Shoemakers and shoe repair workers have been exposed to a large number of agents including benzene, other solvents, glues, adhesives, polishes, and stains.⁴⁴ Workers in the porcelain and earthenware industry have been found to be at increased risk for silicosis, and to have increased rates of lung cancer (although not in Sweden) but non-Hodgkin's lymphoma has not previously been linked with this industry.⁴⁵⁻⁴⁸ Risk of non-Hodgkin's lymphoma was significantly increased among men in Iowa and Minnesota residing near earthenware and pottery factories, however.⁴⁹

Workers in the porcelain and earthenware industry are exposed to fine dust particles and also to dyes, paints, arsenic, lead, and other compounds.⁵⁰

The excesses we found for office workers and bank employees have not been previously described⁵¹⁻⁵³ although these associations may reflect a relation of non-Hodgkin's lymphoma to higher socioeconomic state,^{53,54} rather than a role for specific occupational exposures.

The increased risks of lymphoma among workers in the furniture and furnishings industry and among woodworkers in sawmills and planing mills are consistent with the excesses of non-Hodgkin's lymphoma previously reported for carpenters, sawmill workers, and pulp and paper mill workers in Washington state⁵⁵ and British Columbia.⁵⁶ Potential exposures include not only wood dusts, but also paints, lacquers, varnishes, thinning solvents, and in some situations, antifungal agents, insecticides, and dioxin.⁵⁷

Significantly increased risks were found among engineers and technicians in the electrical power industry. Occupational exposures to electric and magnetic fields have been associated with increased occurrence of non-Hodgkin's lymphoma among Swedish workers in a large telecommunications company,⁵⁸ United States telephone company employees,^{59,60} Canadian power and telephone linemen,⁶¹ and electrical engineers and technicians in surveys in Massachusetts⁶² and Washington State.⁶³ No significant excesses were reported among Swedish electrical engineering graduates,⁶⁴ Swedish electric power linemen and power station operators,⁶⁵ or Swiss electricians.⁶⁶ Residential exposures to magnetic fields have been associated with increased risk of lymphomas in Denver⁶⁷ and the United Kingdom.⁶⁸ As well as possible exposures to magnetic and electrical fields, the Swedish workers may have had

Table 2 Significant ($p < 0.05$) SIRs for non-Hodgkin's lymphomas by specific occupations within industries among Swedish men, 1961-79

Code Occ/Ind	Specific occupation within industry	Cases	Number expected	SIR [†]
203/670	Bank cashier in banking industry	5	0.9	5.8**
201/613	Bookkeeper in wholesale fules	4	0.8	4.8*
001/500	Engineer or technician in electric power plants	4	0.9	4.7*
635/620	Messenger in food business	4	0.9	4.3*
883/233	Warehouse worker in ready made garment industry	5	1.2	4.3*
883/642	Warehouse worker in hardware and machine business	4	1.0	3.9*
050/813	School teacher in other education	5	1.3	3.8*
005/340	Mining engineer or technician in iron and steel industry	10	3.0	3.4**
297/674	Health insurance clerk in health insurance industry	6	1.9	3.2*
003/363	Mechanical engineer or technician in car factories	8	2.5	3.2**
321/614	Travelling salesman in wholesale tools	6	2.1	2.9*
827/201	Dairy worker in dairies	15	8.1	1.8*
778/240	Other woodworkers in sawmills and planing mills	17	9.4	1.8*
721/232	Shoemaker in shoe repairs	22	12.3	1.8*

* $p < 0.05$; ** $p < 0.01$.

[†]Adjusted for age and region.

contact with various solvents, other chemicals, and non-chemical exposures.⁶⁵ In the absence of specific information on exposures of engineers and technicians in the electric power industry, however, it is difficult to determine the role of any agents in the increased risk for non-Hodgkin's lymphoma.⁶⁹

Although we found a significant excess risk among dairy farmers, earlier case-control⁷⁰ and CER²⁸ studies in Sweden found no increased risk of non-Hodgkin's lymphoma among farmers overall or various subgroups of agricultural workers including owners and tenant farmers employed in land or animal husbandry, horticulture, and related occupations (reindeer husbandry, fur farming, care of animals other than livestock, and miscellaneous).²⁶ Some studies from the United States,^{6-9, 62 71-73} the United Kingdom,⁷⁴ Italy,⁷⁵ New Zealand,¹⁰ and Tasmania¹¹ have reported an increased risk for non-Hodgkin's lymphoma among farmers and other agricultural workers, although no excesses were found in other United States^{16 52 76 77} or United Kingdom¹⁸ investigations. The exposures linked to non-Hodgkin's lymphoma among farmers include pesticides, herbicides, fertilisers, and livestock.^{7 13 15} Also, slaughterhouse workers in New Zealand have been reported to be at excess risk of non-Hodgkin's lymphoma,⁷⁸ suggesting that zoonotic agents may have an aetiological role.⁷⁹

HHfle Hcess risk we found among lorry drivers supports earlier reports of significantly increased non-Hodgkin's lymphoma among highway and road transport workers.^{71 80 81} Recognised exposures in this type of work include exhausts from petrol and diesel driven vehicles, benzene, and various solvents.^{82 83}

The raised SIRs for teachers and those employed in research and scientific institutes are consistent with findings from other occupational surveys.^{62 63 84} Whether carcinogenic exposures occur among these workers is not clear, although socioeconomic and dietary factors may also be important.^{53 54 85}

We found no excess of non-Hodgkin's lymphoma among chemists, chemical or rubber manufacturing workers, or in men working in petrochemical refineries as described in other investigations.^{17 22-25 86 87} Also, no excess risk of non-Hodgkin's lymphoma was seen among physicians and dentists as reported in other surveys,^{62 63 71} among postal workers or maintenance workers as reported elsewhere.^{62 63 84}

The shortcomings of the CER such as the lack of direct information on exposures and duration of employment have been previously described.^{26 88} Nevertheless, the CER has identified known high risk occupational activities—for example, mesothelioma among shipyard workers, nasal adenocarcinoma among furniture makers, and liver angiosarcoma among certain chemical workers.^{88 89} Our results, however, may be affected by the problem of multiple comparisons, with some findings reflecting chance

events rather than true occupational associations.

In summary, this nationwide survey of occupation and non-Hodgkin's lymphoma has shown new aetiological leads as well as associations reported in other studies. The overall results, however, suggest that occupational risk factors have a limited role in non-Hodgkin's lymphoma and the rising incidence trends reported for this tumour.

Requests for reprints to: M S Linet, National Cancer Institute, EPN 415B, Rockville, Maryland 20892, USA.

- 1 Doll R, Muir CS, Waterhouse J, eds. *Cancer incidence in five continents*. Vol II. Berlin: Springer-Verlag, 1970.
- 2 Waterhouse J, Muir CS, Correa P, Powell J, eds. *Cancer incidence in five continents*. Vol III. Lyon: IARC, 1986. (IARC publ No 15.)
- 3 Waterhouse J, Muir C, Shanmugaratnam K, Powell J, eds. *Cancer incidence in five continents*. Vol IV. Lyon: IARC, 1986. (IARC sci publ No 42.)
- 4 Muir C, Waterhouse J, Mack T, Powell J, Whelan S, eds. *Cancer incidence in five continents*. Vol V. Lyon: IARC, 1987. (IARC sci publ No 88.)
- 5 Greene MH: Non-Hodgkin's lymphoma and mycosis fungoides. In: Schottenfeld D, Fraumeni JF Jr, eds. *Cancer epidemiology and Prevention*. Philadelphia: WB Saunders Co, 1982;754-78.
- 6 Cantor KP. Farming and mortality from non-Hodgkin's lymphoma: a case-control study. *Int J Cancer* 1982;29:239-47.
- 7 Hoar SK, Blair A, Holmes FF, et al. Agricultural herbicide use and risk of lymphoma and soft-tissue sarcoma. *JAMA* 1986;256:1141-7.
- 8 Dubrow R, Paulson JO, Indian RW. Farming and malignant lymphoma in Hancock County, Ohio. *Br J Ind Med* 1988;45: 25-8.
- 9 Brownson RC, Reif JS. A cancer registry-based study of occupational risk for lymphoma, multiple myeloma and leukaemia. *Int J Epidemiol* 1988;17:27-32.
- 10 Pearce NE, Sheppard RA, Smith AH, Teague CA. Non-Hodgkin's lymphoma and farming: an expanded case-control study. *Int J Cancer* 1987;39:155-61.
- 11 Giles GG, Lickiss JN, Baikie MJ, Lowenthal RM, Panton J. Myeloproliferative and lymphoproliferative disorders in Tasmania. 1972-80: occupational and familial aspects. *J Natl Cancer Inst* 1984;72:1233-39.
- 12 Hardell L, Eriksson M, Lennner P, Lundgren E. Malignant lymphoma and exposure to chemicals, especially organic solvents, chlorophenols, and phenoxyacids. A case control study. *Br J Cancer* 1981;43:169-76.
- 13 Woods JS, Polissar L, Severson RK, Heuser LS, Kulander BG. Soft tissue sarcoma and non-Hodgkin's lymphoma in relation to phenoxyherbicide and chlorinated phenol exposure in western Washington. *J Natl Cancer Inst* 1987;78:899-910.
- 14 Weissenberger DD. Environmental epidemiology of non-Hodgkin's lymphoma in eastern Nebraska. *Am J Ind Med* 1990;18:303-5.
- 15 Burmeister LF, Everett GD, Van Lier SF, Isacson P. Selected cancer mortality and farm practices in Iowa. *Am J Epidemiol* 1983;118:72-7.
- 16 Schumacher MC, Delzell E. A death-certificate case-control study of non-Hodgkin's lymphoma and occupation in men in North Carolina. *Am J Ind Med* 1988;13:317-30.
- 17 Franceschi S, Serraino D, Bidoli E, et al. The epidemiology of non-Hodgkin's lymphoma in the north-east of Italy: a hospital-based case-control study. *Leuk Res* 1989;13:465-72.
- 18 Cartwright RA, McKinney PA, O'Brien C, et al. Non-Hodgkin's lymphoma: case control epidemiological study in Yorkshire. *Leuk Res* 1988;12:81-7.
- 19 Wiklund K, Dick J, Holm L-E. Risk of malignant lymphoma in Swedish pesticide applicators. *Br J Cancer* 1987;56:505-8.
- 20 Brandt L. Leukaemia and lymphoma risks derived from solvents. *Med Oncol Tumor Pharmacother* 1987;4:199-205.
- 21 Olsson H, Brandt L. Risk of non-Hodgkin's lymphoma among men occupationally exposed to organic solvents. *Scand J Work Environ Health* 1988;14:246-51.

- 22 Wong O. An industry wide mortality study of chemical workers occupationally exposed to benzene. II Dose response analyses. *Br J Ind Med* 1987;44:382-95.
- 23 Ott MG, Teta MJ, Greenberg HL. Lymphatic and hematopoietic tissue cancer in a chemical manufacturing environment. *Am J Ind Med* 1989;16:631-43.
- 24 Li FP, Fraumeni JF Jr, Mantel N. Cancer mortality among chemists. *J Natl Cancer Inst* 1969;43:1159-64.
- 25 Tabershaw IR, Cooper WC. *A Mortality study of petroleum workers*. Medical Research Report No EA7402. Prepared for the American Petroleum Institute. Tabershaw/Cooper Associates Inc, Berkeley, California, September, 1974.
- 26 Wiklund K, Lindefors B-M, Holm L-E. Risk of malignant lymphoma in Swedish agricultural and forestry workers. *Br J Ind Med* 1988;45:19-24.
- 27 Wiklund K, Holm L-E. Soft tissue sarcoma in Swedish agricultural and forestry workers. *J Natl Cancer Inst* 1986;86:229-34.
- 28 Blair A, Malker H, Cantor KP, Burmeister L, Wiklund K. Cancer among farmers: A review. *Scand J Work Environ Health* 1985;11:397-407.
- 29 Wiklund K, Einhorn J, Wenstrom G, Rapaport E. A Swedish cancer environment registry available for research. *Scand J Work Environ Health* 1981;7:63-7.
- 30 National Board of Health and Welfare. *Cancer incidence in Sweden, 1988*. Stockholm: Allmanna Forlaget, 1991.
- 31 NCI non-Hodgkin's classification writing committee. Classification of non-Hodgkin's lymphoma. *Cancer* 1982;55:91-5.
- 32 Dick F, Van Lier S, Banks P, et al. Use of the working formulation for non-Hodgkin's lymphoma in epidemiologic studies: agreement between reported diagnosis and a panel of experienced pathologists. *J Natl Cancer Inst* 1984;78:1137-44.
- 33 Bailar JC, Ederer F. Significance factors for the ratio of a Poisson variable to its expectation. *Biometrics* 1964;20:639-43.
- 34 International Labor Office. *International standard classification of occupations*. Geneva: International Labour Office, 1958.
- 35 United Nations Statistical Office. *International standard classification of occupations*. Geneva: International Labour Office, 1958.
- 36 Vigliani EC, Saita B. Benzene and leukemia. *N Engl J Med* 1964;271:872-6.
- 37 Aksoy M. Malignancy due to occupational exposure to benzene. *Am J Ind Med* 1985;7:395-402.
- 38 Aksoy M. Different types of malignancies due to occupational exposure to benzene: a review of recent observations in Turkey. *Environ Res* 1980;23:181-90.
- 39 Decouffe P, Walrath J. Proportionate mortality among US shoeworkers, 1966-1977. *Am J Ind Med* 1983;4:523-32.
- 40 Garabrant DH, Wegman DH. Cancer mortality among shoe and leather workers in Massachusetts. *Am J Ind Med* 1984;5:303-14.
- 41 Walrath J, Decouffe P, Thomas TL. Mortality among workers in a shoe manufacturing company. *Am J Ind Med* 1987;12:615-23.
- 42 Pippard EC, Acheson ED. The mortality of boot and shoe makers with special reference to cancer. *Scand J Work Environ Health* 1985;11:249-55.
- 43 Paci E, Buiatti E, Seniori Constantini A, et al. Aplastic anemia, leukemia and other cancer mortality in a cohort of shoe workers exposed to benzene. *Scand J Work Environ Health* 1989;15:313-8.
- 44 International Agency for Research on Cancer. Wood, leather, and some associated industries. *IARC Monogr* 1981;25:249-77.
- 45 Goldsmith DF, Winn DM, Shy CM. *Silica, silicosis and cancer*. New York: Praeger Publications, 1986.
- 46 International Agency for Research on Cancer. Silica and some silicates. *IARC Monogr* 1987;42:39-143.
- 47 Simonato L, Fletcher AC, Saracci R, Thomas TL. *Occupational Exposure to silica and cancer risk*. Lyon: IARC, 1990. (IARC sci publ No 97.)
- 48 Westerholm P. Silicosis. *Scand J Work Environ Health* 1980;6(suppl 2):34-7.
- 49 Linos A, Blair A, Gibson RW, et al. Leukemia and non-Hodgkin's lymphoma and residential proximity to industrial plants. *Arch Environ Health* 1991;46:70-4.
- 50 Key MM, Henschel AF, Butler J, Ligo RN, Tabershaw IR. *Occupational disease: a guide to their recognition*. USDHEW, PHS, CDC, NIOSH, publ No 77-181, 1977.
- 51 Registrar General for England and Wales. *The Registrar General's decennial supplement, England and Wales. Occupational Mortality, Part II, Vol 2*. London: HMSO, 1958.
- 52 Guralnick L. Mortality by occupation (or industry) and cause of death among men 20-64 years of age: United States, 1950. Washington, DC: Vital statistics special report 1963;53:95-433.
- 53 Cantor KP, Fraumeni JF Jr. Geographic and temporal patterns of non-Hodgkin's lymphoma mortality in US counties, 1950-1975. *Cancer Res* 1980;40:2645-52.
- 54 Vagero D, Persson G. Occurrence of cancer in socioeconomic groups in Sweden. An analysis based on the Swedish Cancer Environment Registry. *Scand J Soc Med* 1986;14:151-8.
- 55 Milham S Jr, Demers RY. Mortality among pulp and paper workers. *J Occup Med* 1984;26:844-6.
- 56 Svirchev LM, Gallagher RP, Band PR, Threlfall WJ, Spinelli JJ. Gastric cancer and lymphosarcoma among wood and pulp workers. *J Occup Med* 1986;28:264-5.
- 57 Wills JH. Nasal carcinoma in woodworkers: a review. *J Occup Med* 1982;24:526-30.
- 58 Vagero D, Ahlbom A, Olin R, Sahlsten S. Cancer morbidity among workers in the telecommunications industry. *Br J Ind Med* 1985;42:191-5.
- 59 Milham S Jr. Mortality from leukaemia in workers exposed to electrical and magnetic fields. *N Engl J Med* 1982;307:249.
- 60 Matanoski G, Elliott E, Breyse P. *Poster presented at the annual Department of Energy/Electric Power Research Institute contractors review on biologic effects from electric and magnetic fields*. Portland, Oregon; November, 1989.
- 61 Howe GR, Lindsay JP. A follow-up study of a ten-percent sample of the Canadian Labour Force. I. Cancer mortality in males, 1965-73. *J Natl Cancer Inst* 1983;70:37-44.
- 62 Dubrow R, Wegman DH. Cancer and occupation in Massachusetts: A death certificate study. *Am J Ind Med* 1984;6:207-30.
- 63 Milham S, Jr. Occupational mortality in Washington State 1950-1979. Cincinnati, Ohio: USDHEW (PHS) (CDC) (NIOSH). DHSS publ No 83-116, 1983.
- 64 Olin R, Vagero D, Ahlbom S. Mortality experience of electrical engineers. *Br J Ind Med* 1985;42:211-2.
- 65 Tornqvist S, Norell S, Ahlbom A, Knave B. Cancer in the electric power industry. *Br J Ind Med* 1986;43:212-3.
- 66 Guberan E, Usel M, Raymond L, Tissot R, Sweetman PM. Disability, mortality, and incidence of cancer among Geneva painters and electricians. *Br J Ind Med* 1989;46:16-23.
- 67 Wertheimer N, Leeper E. Adult cancer related to electrical wires near the home. *Int J Epidemiol* 1982;11:345-55.
- 68 McDowall ME. Mortality of persons resident in the vicinity of electrical transmission facilities. *Br J Cancer* 1986;53:271-9.
- 69 Savitz DA, Pearce NE, Poole C. Methodological issues in the epidemiology of electromagnetic fields and cancer. *Epidemiol Rev* 1989;11:59-78.
- 70 Persson P, Dahlander A-M, Frederiksson M, et al. Malignant lymphomas and occupational exposures. *Br J Ind Med* 1989;46:516-20.
- 71 Williams RR, Stegens NL, Goldsmith JR. Associations of cancer site and type with occupation and industry from the Third National Cancer Survey Interview. *J Natl Cancer Inst* 1977;59:1147-85.
- 72 Burmeister LF. Cancer mortality in Iowa farmers, 1971-78. *J Natl Cancer Inst* 1981;66:461-4.
- 73 Goldsmith JR, Guidotti TL. Environmental factors in the epidemiology of lymphosarcoma. *Pathol Annu* 1977;12:411-25.
- 74 Fox AJ, Goldblatt PO. *Longitudinal study of sociodemographic mortality differentials, 1971-75*. Office of Population Censuses and Surveys. London: HMSO, 1980.
- 75 LaVecchia C, Negri E, D'Avanzo B, Franceschi S. Occupation and lymphoid neoplasms. *Br J Cancer* 1989;60:385-8.
- 76 Decouffe P, Stanislawsky K, Houten L, Bross IDJ, Viadana E. A retrospective survey of cancer in relation to occupation. Cincinnati Ohio: NIOSH. (DHEW (NIOSH) publ No 77-178.)
- 77 Fasal E, Jackson EW, Klauber MR. Leukemia and lymphoma mortality and farm residence. *Am J Epidemiol* 1968;87:267-74.
- 78 Pearce N, Smith AH, Reif JS. Increased risks of soft tissue sarcoma, malignant lymphoma, and acute myeloid leukemia in abattoir workers. *Am J Ind Med* 1988;14:63-72.
- 79 Miller JM, Miller LD, Olson C, Gillette K. Virus-like particles in phytohemagglutinin-stimulated lymphocyte cultures with reference to bovine lymphosarcoma. *J Natl Cancer Inst* 1969;43:1297-305.
- 80 Balarajan R. Malignant lymphomas in road transport workers. *J Epidemiol Comm Health* 1983;37:279-80.
- 81 Maizlish N, Beaumont J, Singleton J. Mortality among Californian highway workers. *Am J Ind Med* 1988;13:363-79.

- 82 Boffetta P, Stellman S, Garfinkel L. Diesel exhaust exposure and mortality among males in the American Cancer Society prospective study. *Am J Ind Med* 1988;14:403-15.
- 83 Wallace LA. Major sources of benzene exposure. *Environ Health Perspect* 1989;82:165-9.
- 84 Dubrow R, Wegman DH. Setting priorities for occupational cancer research and control: Synthesis of the results of occupational disease surveillance studies. *J Natl Cancer Inst* 1983;71:1123-42.
- 85 Franceschi S, Serraino D, Carbone A, Talamini R, La Vecchia C. Dietary factors and non-Hodgkin's lymphoma: a case-control study in the northeastern part of Italy. *Nutr Cancer* 1989;12:333-41.
- 86 Olin GR, Ahlbom A. The cancer mortality among Swedish chemists graduated during three decades. *Environ Res* 1980;22:154-61.
- 87 Wong O, Raabe GK. Critical review of cancer epidemiology in petroleum industry employees with a quantitative meta-analysis by cancer site. *Am J Ind Med* 1989;15:283-310.
- 88 Malaker HSR. Register-epidemiology in the identification of cancer risks. *Arbete Och Halsa* 1988;21:7-50.
- 89 McLaughlin JK, Malaker HSR, Malaker BK, et al. Registry-based analysis of occupational risks for primary liver cancer in Sweden. *Cancer Res* 1987;47:287-91.

Accepted 30 March 1992

Destruction of manuscripts

From 1 July 1985 articles submitted for publication will not be returned. Authors whose papers are rejected will be advised of the decision and the manuscripts will be kept under security for three months to deal with any inquiries and then destroyed.