Results Estimates of exposure for 138 EMF sources were obtained from measurements (1424 aggregated records) extracted from 71 papers and hygiene reports (1974–2013). For each source, exposure was calculated by frequency band and dosimetry type, as the arithmetic and geometric means of all measurements identified. Standard deviations were included in order to characterise the variability of the estimates.

Conclusions A source-exposure matrix has been constructed for the most common sources of EMF in the workplace, based on the responses to the INTERPHONE-INTEROCC study questionnaire. This database currently represents the most comprehensive source of information on occupational EMF exposure and is available on request to researchers.

0286 OCCUPATIONAL USE OF INSECTICIDES, FUNGICIDES AND FUMIGANTS AND RISK OF NON-HODGKIN LYMPHOMA AND MULTIPLE MYELOMA IN THE AGRICULTURAL HEALTH STUDY

¹Michael Alavanja, ¹Jonathan Hofmann, ²Charles Lynch, ³Cynthia Hines, ¹Kathryn Barry, ⁴Joseph Barker, ⁴Dennis Buckman, ⁵Kent Thomas, ⁶Dale Sandler, ⁷Jane Hoppin, ¹Stella Koutros, ¹Gabriella Andreotti, ¹Jay Lubin, ¹Aaron Blair, ¹Laura Beane Freeman. ¹Occupational and Environmental Epidemiology Branch, National Cancer Institute, Bethesda, MD, USA; ²University of Iowa, Department of Epidemiology, Iowa City, IA, USA; ³National Institute for Occupational Safety and Health, Cincinnati, OH, USA; ⁴IMS, Calverton, MD, USA; ⁵National Exposure Research Laboratory, U. S. Environmental Protection Agency, Research Triangle, NC, USA; ⁶National Institute of Environmental Health Sciences, Research Triangle, NC, USA; ⁷North Carolina State University, Raleigh, NC, USA

10.1136/oemed-2014-102362.111

Objectives Farming and exposure to pesticides have been linked to non-Hodgkin lymphoma (NHL), and multiple myeloma (MM) in previous studies. We evaluated use of insecticides, fungicides and fumigants and risk of NHL, including MM and other NHL sub-types in the Agricultural Health Study, a US-based prospective cohort study.

Method A total of 527 cases occurred among 55 875 pesticide applicators from enrollment (1993–1997) through 2011 in Iowa and 2010 in North Carolina. Information on pesticide use, other agricultural exposures and other factors was obtained from questionnaires at enrollment and follow-up approximately five years later (1999–2005). Information from these questionnaires was used to create lifetime-days and intensity-weighted lifetime-days of pesticide use. Poisson regression and polytomous logit models were used to calculate relative risks (RR) and 95% confidence intervals (CI) to evaluate associations between 26 pesticides and NHL and five NHL-subtypes including multiple myeloma, while adjusting for potential confounding factors.

Results Statistically significant positive exposure-response trends occurred between overall NHL risk and lindane (p-trend = 0.004) and DDT (p-trend = 0.02). In addition, ever use of terbufos was associated with NHL overall (RR=1.2; CI=1.0–1.5), but with no exposure-response trend. In sub-type analyses, terbufos and DDT were associated with small cell lymphoma/chronic lymphocytic leukaemia/marginal cell lymphoma. In addition, lindane and diazinon were associated with follicular lymphoma and permethrin with MM although tests of homogeneity did not show significant differences in exposure-response among NHL-subtypes for any chemical.

Conclusions These findings are among the first to suggest links between DDT, lindane, permethrin, diazinon and terbufos and specific NHL subtypes.

0288 GENE-SPECIFIC DNA METHYLATION AS A VALUABLE TOOL FOR RISK ASSESSMENT: THE CASE OF OCCUPATIONAL EXPOSURE TO DIFFERENT VOC'S IN MEXICAN WORKERS

¹Octavio Jiménez-Garza, ²Andrea Baccarelli, ²Hyang-Min Byun, ³Giovanni Battista Bartolucci, ³Mariella Carrieri. ¹University of Guanajuato Campus León, Health Sciences Division, León, Guanajuato, Mexico; ²Harvard School of Public Health, Laboratory of Human Environmental Epigenomics, Boston, Massachusetts, USA; ³University of Padova, Department of Medicine, Padova, Italy

10.1136/oemed-2014-102362.112

Objectives To determine gene-specific methylation levels (promoter region) on genes from critical cellular pathways in persons occupationally exposed to a single volatile organic compound (VOC) or to a mixture of them

Method Workers from two tanneries, two shoe factories, and two gas stations were included (exposed groups to VOC's). We also included administrative workers (reference group), all of them from the city of León Guanajuato, México. In tannery workers we measured individual exposure levels to three different VOC's; for the rest of the groups we measured up to 7 VOC's. After exposure characterisation, we took blood samples and extracted DNA in order to determine, by PCR-pyrosequencing, methylation levels in genes involved in inflammation, DNA repair, oxidative stress and xenobiotic metabolism pathways.

Results Only toluene environmental levels were higher in tannery workers. Workers from the leather shoe factory showed the highest exposure levels for up to five different VOC's. There was no statistical significant difference in gene-specific methylation for tannery workers when compared to the control group. For the leather shoe factory workers, we found hypermethylation in the TNF α , SOD1 and TOP2A promoter regions compared to the control group or to other exposed groups. Gas station attendants showed hypermethylation for the IL6 gene compared to controls

Conclusions Occupational exposure to a mixture of VOC's has important effects on the methylation status of genes involved in inflammation, DNA repair and oxidative stress. These epigenetic changes, detectable at a pre-clinical stage, represent a valuable tool for performing an early risk assessment in these populations

0289 PREVENTING NEEDLESTICKS AND OTHER SHARPS INJURIES TO HOME CARE AIDES: RESULTS OF A SURVEY TO IDENTIFY HAZARDS DURING HOME VISITS

¹David Kriebel, ¹Natalie Brouillette, ¹Pia Markkanen, ¹Catherine Galligan, ¹Susan Sama, ¹Rebecca Gore, ²Angela Laramie, ¹Daniel Okyere, ¹Chuan Sun, ²Letitia Davis, ¹Margaret Quinn. ¹University of Massachusetts Lowell, Lowell, MA, USA; ²Massachusetts Department of Public Health, Boston MA, USA

10.1136/oemed-2014-102362.113

Objectives To follow up on our previous finding that home care (HC) aides have a substantial risk of injury with used sharp medical devices, this study's goal was to identify modifiable aspects of aide-client encounters during HC visits that increase risk of sharps injuries (SI).

Method A survey of 1249 HC aides was conducted in eastern Massachusetts. Approximately half the participants (634) were employed by HC agencies, and half (615) directly by HC clients and their families. A questionnaire gathered data on aides' most recent home visits, enabling quantification of hazardous working conditions. The specific investigation described here focused on understanding determinants of hazardous conditions likely to increase the risk of SI including: the aide finding used sharps