ation of these microtraumas might be the cause of a radiologically visible zone of decalcification.

The presence of these lesions is a sensitive indicator of exposure to light tool vibrations. The lesions appear before clinical symptoms become manifest. This is of great importance for the implementation of preventive measures. The question whether the lesions disappear after withdrawal from the risk cannot yet be answered. We need therefore a longitudinal study that is not yet finished.

There is some evidence that protection of the hand with gloves, covered with special absorbing materials of different thicknesses, provokes a clear diminution of the vibrations: from  $4\cdot10^{-1}$ m/sec<sup>2</sup> to  $10\cdot1$  m/sec<sup>2</sup> by 125 Hz (a damping factor  $0\cdot18$  to  $0\cdot72$ ). Further research on this means of protection is necessary.

#### References

Brailsford J. Pathological changes in bones and joints induced by injuries. Br J Med 1936;ii:657.

- <sup>2</sup> Heidenhoffer J. Zur ursächlichen Stellung der Lunatum malacie in Rahmen der örtlichen Malazien. Fortschr Rontgenstr 1949;71:472-5.
- <sup>3</sup> Hunter D. The diseases of occupations. London: English Universities Press, 1964.
- 4 Iowata H. Effects of rockdrillers on operator. Ind Health 1968;6:37-46.
- <sup>5</sup> Linder GS. Mechanical vibration effects on human beings. Aerospace Medicine 1962;33:939-50.
- 6 Hempstock TI, O'Conner DE. The measurement of hand-arm vibration. In: Taylor W, Pelmear PL, eds. Vibration white finger in industry. London; Academic Press, 1975.
- Agate JH. Druett HA. A study of vibrating tools in relation to the clinical effects which they produce. Br J Ind Med 1947;4:141-63.
- Stewart MA, Goda D. Vibration syndrome. Br J Ind Med 1970;27:19-27.
- Hattner RA, McMillan DE. Influence of weightlessness upon the skeleton: a review. Aerospace Medicine 1968;39:849-55.
- <sup>10</sup> Pace N. Weightlessness: A matter of gravity. N Engl J Med 1977;7:32-6.
- Teissinger J, Louda L. Vascular disorders resulting from vibrating tools. JOM 1972;14:129-33.

## **Notice**

### Course in occupational neurotoxicology 17-21 September 1984

A course in occupational neurotoxicology will be held at the Institute of Neurology, Queen Square. Topics to be covered will include clinical and neuropathological aspects of peripheral neuropathy, the epidemiology of occupational neuropathy and psychopathy, clinical and psychological aspects of toxic organic psychoses, neurological examination, psychological testing in the field, and neurophysiological methods. In addition, a whole day will be spent at the MRC Toxicology Unit at Carshalton. The fee for the course will be £150 including registration. Participants will be responsible for their own travel and accommodation. Further details of the course may be obtained from Dr H A Waldron, TUC Centenary Institute of Occupational Health, London School of Hygiene and Tropical Medicine, Keppel Street (Gower Street), London WC1E 7HT.

#### Correction

# Reversibility of skeletal fluorosis (November 1983)

We regret that figure 3 (p. 458) was printed upside down. This was due to incorrect positioning by the printers.