

Subject index

- 2,3,7,8-tetrachlorodibenzo-p-dioxin**, Cancer risk for chemical workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin, **672**
- 2-EEA**, Haematological effects among silk screening workers exposed to 2-ethoxy ethyl acetate, **e7**
- 2-methoxyethanol**, Follow up study of haematological effects in workers exposed to 2-methoxyethanol, **130**
- 8-hydroxydeoxyguanosine**, Chromium (VI) induced oxidative damage to DNA: increase of urinary 8-hydroxydeoxyguanosine concentrations (8-OHdG) among electroplating workers, **590**
- 8-OHdG, 8-hydroxy-2'-deoxyguanosine**, Temporal variation of hydroxyl radical generation and 8-hydroxy-2'-deoxyguanosine formation by coarse and fine particulate matter, **315**
- absenteeism**, Return to work of cancer survivors: a prospective cohort study into the quality of rehabilitation by occupational physicians, **352**
- acoustic rhinometry**, Upper airway inflammation in waste handlers exposed to bioaerosols, **444**
- activity**, Knee disorders in the general population and their relation to occupation, **794**
- acute leukaemia**, Occupational exposure to electromagnetic fields and acute leukaemia: analysis of a case-control study, **577**
- adaptation**, The impact of a week of simulated night work on sleep, circadian phase, and performance, **e13**
- adjustment disorder**, Reducing long term sickness absence by an activating intervention in adjustment disorders: a cluster randomised controlled design, **429**
- age-period-cohort model**, Update of predictions of mortality from pleural mesothelioma in the Netherlands, **50**
- agreement**, Agreement between hearing thresholds measured in non-soundproof work environments and a soundproof booth, **667**
- agricultural workers**, Ischaemic heart disease among livestock and agricultural workers, **e1**
- agriculture**, Endotoxin: is it an environmental factor in the cause of Parkinson's disease?, **378**
Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
- air cyst**, Should computed chest tomography be recommended in the medical certification of professional divers? A report of three cases with pulmonary air cysts, **606**
- air pollution**, Acute effects of winter air pollution on respiratory function in schoolchildren in southern England, **82**
Gas stove use and respiratory health among adults with asthma in NHANES III, **759**
How much does the environment contribute to cancer?, **150**
Personal exposure assessment in the epidemiology of air pollutants, **458**
Respiratory effects of sulphur dioxide: a hierarchical multicity analysis in the APHEA 2 study, **e2**
Risk factors among elderly for short term deaths related to high levels of air pollution, **684**
The biological effects of coarse and fine particulate matter, **313**
The effect of sulphurous air pollutant exposures on symptoms, lung function, exhaled nitric oxide, and nasal epithelial lining fluid antioxidant concentrations in normal and asthmatic adults, **e15**
Use of personal exposure modelling in risk assessment of air pollutants, **529**
William Harvey and air pollution, **147**
- air quality**, NO₂, as a marker of air pollution, and recurrent wheezing in children: a nested case-control study within the BAMSE birth cohort, **876**
- air sampling**, Municipal waste incinerators: air and biological monitoring of workers for exposure to particles, metals, and organic compounds, **563**
- aircraft maintenance**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
- airflow obstruction**, Chronic obstructive pulmonary disease due to occupational exposure to silica dust: a review of epidemiological and pathological evidence, **237**
- airline crew**, SARS and occupational health in the air, **539**
- airway irritation**, Effects of bioaerosol polluted outdoor air on airways of residents: a cross sectional study, **336**
- airway responsiveness**, A longitudinal observation of early pulmonary responses to cotton dust, **115**
- alcohol**, Driving impairment due to sleepiness is exacerbated by low alcohol intake, **689**
- allergen**, Estimating the incidence of occupational asthma and rhinitis from laboratory animal allergens in the UK, 1999–2000, **604**
- aluminium potrooms**, World at work: Hazards and controls in aluminium potrooms, **989**
- ambulance workers**, Acute and chronic job stressors among ambulance personnel: predictors of health symptoms, **i40**
- amino acid metabolism**, Increased plasma homocysteine levels in shift working bus drivers, **662**
- ammonia**, Potential effects on human health of an ammonia rich atmospheric environment in an archaeologically important cave in southeast Asia, **986**
- ammonium bisulphate**, The effect of sulphurous air pollutant exposures on symptoms, lung function, exhaled nitric oxide, and nasal epithelial lining fluid antioxidant concentrations in normal and asthmatic adults, **e15**
- amyl nitrite**, Alternative methods of administering amyl nitrite to victims of cyanide poisoning, **147**
- anaesthetics**, Multiple sclerosis in nurse anaesthetists, **66**
- ANCOVA**, Fatigue, burnout, and chronic fatigue syndrome among employees on sick leave: do attributions make the difference?, **i26**
- annoyance**, Health status as a potential effect modifier of the relation between noise annoyance and incidence of ischaemic heart disease, **739**
- antioxidants**, Oxidative stress: its role in air pollution and adverse health effects, **612**
The effect of sulphurous air pollutant exposures on symptoms, lung function, exhaled nitric oxide, and nasal epithelial lining fluid antioxidant concentrations in normal and asthmatic adults, **e15**
- archaeology**, Potential effects on human health of an ammonia rich atmospheric environment in an archaeologically important cave in southeast Asia, **986**
- arthritis**, Knee disorders in the general population and their relation to occupation, **794**
- asbestos**, Earlier study on asbestos workers, ILO scores, and oxygenation more comprehensive, **611**
Evaluation of the quality of coding of job episodes collected by self questionnaires among French retired men for use in a job-exposure matrix, **438**
Occupational exposures and lung cancer in New Caledonia, **584**
- asbestos exposures**, Asbestos related diseases from environmental exposure to crocidolite in Da-yao, China. I. Review of exposure and epidemiological data, **35**
- asbestosis**, Earlier study on asbestos workers, ILO scores, and oxygenation more comprehensive, **611**
Radiographic (ILO) readings predict arterial oxygen desaturation during exercise in subjects with asbestosis, **201**
- ascertainment strategies**, Under-ascertainment of multiple myeloma among participants in UK atmospheric atomic and nuclear weapons tests, **e18**
- ash**, Respiratory effects of volcanic emissions, **529**
- asthma**, Dermatological powder as hidden cause of occupational allergy due to casein: a case report, **609**
Estimating the incidence of occupational asthma and rhinitis from laboratory animal allergens in the UK, 1999–2000, **604**
Gas stove use and respiratory health among adults with asthma in NHANES III, **759**
NO₂, as a marker of air pollution, and recurrent wheezing in children: a nested case-control study within the BAMSE birth cohort, **876**
Occupational asthma, **711**
Occupational asthma due to IgE mediated allergy to the flower *Molucella laevis* (Bells of Ireland), **701**
Oxidative stress: its role in air pollution and adverse health effects, **612**
Respiratory effects of sulphur dioxide: a hierarchical multicity analysis in the APHEA 2 study, **e2**

- Volcanic ash and respiratory symptoms in children on the island of Montserrat, British West Indies, **207**
- World at work: Hazards and controls in aluminium potrooms, **989**
- atherosclerosis**, Increased plasma homocysteine levels in shift working bus drivers, **662**
- atopy**, A longitudinal observation of early pulmonary responses to cotton dust, **115**
- Australia**, The right treatment to the right patient at the right time, **235**
- autoimmune**, Occupational silica exposure and risk of various diseases: an analysis using death certificates from 27 states of the United States, **122**
- back disorders**, Effects of measurement strategy and statistical analysis on dose-response relations between physical workload and low back pain, **942**
- back pain**, Generalised estimating equations and low back pain, **378**
Sickness absence due to low back pain, **306**
The management of low back pain, **617**
The relative importance of whole body vibration and occupational lifting as risk factors for low-back pain, **715**
- background radiation**, No change in sex ratio in Ramsar (north of Iran) with high background of radiation, **146**
- bakeries**, Prevalence of sensitisation to cellulase and xylanase in bakery workers, **802**
- bakery workers**, Prevalence of sensitisation to cellulase and xylanase in bakery workers, **802**
- Bayesian statistics**, Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
- benzene**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
Lymphohaematopoietic cancer mortality among workers with benzene exposure, **676**
Lymphohaematopoietic system cancer incidence in an urban area near a coke oven plant: an ecological investigation, **187**
- bias**, Mortality of petroleum refinery workers, **304**
- biliary tract cancer**, Wastewater exposure and health—a comparative study of two occupational groups, **595**
- bioaerosol exposure**, Effects of bioaerosol polluted outdoor air on airways of residents: a cross sectional study, **336**
Upper airway inflammation in waste handlers exposed to bioaerosols, **444**
- biological monitoring**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
Biological monitoring of occupational exposure to N,N-dimethylacetamide with identification of a new metabolite, **746**
Can salivary lead be used for biological monitoring of lead exposed individuals?, **696**
Exposure to nitroaromatic explosives and health effects during disposal of military waste, **483**
Follow up study of haematological effects in workers exposed to 2-methoxyethanol, **130**
Municipal waste incinerators: air and biological monitoring of workers for exposure to particles, metals, and organic compounds, **563**
Occupational exposure of midwives to nitrous oxide on delivery suites, **958**
Surveillance in occupational health, **705**
- biomarker**, Exposure assessment of monoterpenes and styrene: a comparison of air sampling and biomonitoring, **599**
- biotransformation**, Biological monitoring of occupational exposure to N,N-dimethylacetamide with identification of a new metabolite, **746**
- black smoke**, Risk factors among elderly for short term deaths related to high levels of air pollution, **684**
- bladder cancer**, Mortality of employees of a perfluorooctanesulphonyl fluoride manufacturing facility, **722**
- blood lead**, Yearly measurements of blood lead in Swedish children since 1978: an update focusing on the petrol lead free period 1995–2001, **370**
- blood pressure**, Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
The effect on ambulatory blood pressure of working under favourably and unfavourably perceived supervisors, **468**
- Borneo**, Potential effects on human health of an ammonia rich atmospheric environment in an archaeologically important cave in southeast Asia, **986**
- brain neoplasm**, Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
- break duration**, The amount of sleep obtained by locomotive engineers: effects of break duration and time of break onset, **e17**
- breast cancer**, Breast cancer and serum organochlorine residues, **348**
- breath**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
- bronchial epithelium**, Repeated daily exposure to 2 ppm nitrogen dioxide upregulates the expression of IL-5, IL-10, IL-13, and ICAM-1 in the bronchial epithelium of healthy human airways, **892**
- building related symptoms**, Inflammatory potential of dust from schools and building related symptoms, **e5**
- burnout**, Acute and chronic job stressors among ambulance personnel: predictors of health symptoms, **i40**
Fatigue, burnout, and chronic fatigue syndrome among employees on sick leave: do attributions make the difference?, **i26**
How to conduct research on burnout: advantages and disadvantages of a unidimensional approach in burnout research, **i16**
Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
- byssinosis**, Respiratory symptoms and cotton dust exposure; results of a 15 year follow up observation, **935**
- cabin crew**, Cancer incidence in airline cabin crew, **805**
Cancer incidence in airline cabin crew: experience from Sweden, **810**
- cancer**, How much does the environment contribute to cancer?, **150**
Lung cancer in heavy equipment operators and truck drivers with diesel exhaust exposure in the construction industry, **516**
Mortality and cancer incidence in a cohort of meatworkers, **e4**
Mortality in workers in poultry slaughtering/processing plants: the Missouri poultry cohort study, **784**
World at work: Hazards and controls in aluminium potrooms, **989**
- cancer incidence**, Cancer incidence in airline cabin crew, **805**
Cancer incidence in airline cabin crew: experience from Sweden, **810**
Follow up of mortality and incidence of cancer 1952–98 in men from the UK who participated in the UK's atmospheric nuclear weapon tests and experimental programmes, **165**
Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
Mortality and cancer morbidity in a cohort of Canadian petroleum workers, **918**
- cancer mapping**, Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
- cancer survivors**, Return to work of cancer survivors: a prospective cohort study into the quality of rehabilitation by occupational physicians, **352**
- carbon monoxide**, Are personal and static samples related?, **224**
Memory one month after acute carbon monoxide intoxication: a prospective study, **212**
- cardiorespiratory fitness**, Cardiorespiratory fitness and risk of disability pension: a prospective population based study in Finnish men, **765**
- cardiovascular disease**, Need for recovery after work and the subsequent risk of cardiovascular disease in a working population, **i83**
Workplace bullying and the risk of cardiovascular disease and depression, **779**
- cardiovascular risk**, Downsizing of staff is associated with lowered medically certified sick leave in female employees, **e9**
Health status as a potential effect modifier of the relation between noise annoyance and incidence of ischaemic heart disease, **739**
- case-control study**, Cancer risk around the nuclear power plants of Trillo and Zorita (Spain), **521**
Musicians playing wind instruments and risk of lung cancer: is there an association?, **143**
Occupational exposure to electromagnetic fields and acute leukaemia: analysis of a case-control study, **577**
- case-control study**, Occupational risk factors for symptomatic lumbar disc herniation; a case-control study, **821**
Parental occupation at periconception: findings from the United Kingdom Childhood Cancer Study, **901**
- casein**, Dermatological powder as hidden cause of occupational allergy due to casein: a case report, **609**
- case-referent study**, Lung cancer risk in workers exposed to poly (vinyl chloride) dust: a nested case-referent study, **423**
- cellulase**, Prevalence of sensitisation to cellulase and xylanase in bakery workers, **802**
- childhood asthma**, Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools, **385**

- childhood cancer**, Parental occupation at periconception: findings from the United Kingdom Childhood Cancer Study, **901**
- childhood cancer, pesticide**, Cancer risk in offspring of male pesticide applicators in agriculture in Sweden, **798**
- children**, Acute effects of winter air pollution on respiratory function in schoolchildren in southern England, **82**
- chlorinated hydrocarbons**, Increased mortality odds ratio of male liver cancer in a community contaminated by chlorinated hydrocarbons in groundwater, **364**
- chlorine**, Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools, **385**
- chlorothalonil**, Occupational asthma from fungicides fluzinam and chlorothalonil, **76**
- chromium (VI)**, Chromium (VI) induced oxidative damage to DNA: increase of urinary 8-hydroxydeoxyguanosine concentrations (8-OHdG) among electroplating workers, **590**
- chronic bronchitis**, Chronic obstructive pulmonary disease due to occupational exposure to silica dust: a review of epidemiological and pathological evidence, **237**
- chronic fatigue syndrome**, Fatigue, burnout, and chronic fatigue syndrome among employees on sick leave: do attributions make the difference?, **i26**
- chronic obstructive pulmonary disease**, Chronic obstructive pulmonary disease due to occupational exposure to silica dust: a review of epidemiological and pathological evidence, **237**
Respiratory effects of sulphur dioxide: a hierarchical multicity analysis in the APHEA 2 study, **e2**
- chronic pain**, The right treatment to the right patient at the right time, **235**
- circadian rhythms**, Increased plasma homocysteine levels in shift working bus drivers, **662**
- clinical features**, Fatigue, burnout, and chronic fatigue syndrome among employees on sick leave: do attributions make the difference?, **i26**
- coal tar pitch volatiles**, World at work: Hazards and controls in aluminium potrooms, **989**
- cohort study**, Cardiorespiratory fitness and risk of disability pension: aprospective population based study in Finnish men, **765**
Early prognostic factors for duration on temporary total benefits in the first year among workers with compensated occupational soft tissue injuries, **244**
Fatigue as a risk factor for being injured in an occupational accident: results from the Maastricht Cohort Study, **i88**
Occupations with increased risk of pancreatic cancer in the Swedish population, **570**
Organisational justice and health of employees: prospective cohort study, **27**
Work schedules and fatigue: a prospective cohort study, **i47**
- cold challenge plethysmography**, Skin temperature recovery from cold provocation in workers exposed to vibration: a longitudinal study, **962**
- colour vision**, Relation between colour vision loss and occupational styrene exposure, **222**
- compensation**, The right treatment to the right patient at the right time, **235**
- competence**, Competence at the workplace, **225**
- computed tomography**, Should computed chest tomography be recommended in the medical certification of professional divers? A report of three cases with pulmonary air cysts, **606**
- confounders**, Confounding and confounders, **227**
- confounding**, Confounding and confounders, **227**
- congenital anomaly**, Hazardous waste landfill sites and congenital anomalies, **79**
- construction industry**, Radiographic abnormalities among construction workers exposed to quartz containing dust, **410**
- contaminated land**, Land contamination and urinary abnormalities: cause for concern?, **463**
- contrast sensitivity**, Visual and ocular changes associated with exposure to two tertiary amines, **69**
- COPD**, Occupational silica exposure and risk of various diseases: an analysis using death certificates from 27 states of the United States, **122**
Respiratory effects of sulphur dioxide: a hierarchical multicity analysis in the APHEA 2 study, **e2**
- corneal opacity**, Visual and ocular changes associated with exposure to two tertiary amines, **69**
- coronary disease**, Increased plasma homocysteine levels in shift working bus drivers, **662**
- cortisol**, Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery, **373**
Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
- cosmic radiation**, Breast cancer risk in airline cabin attendants: a nested case-control study in Iceland, **807**
Cancer incidence in airline cabin crew, **805**
Cancer incidence in airline cabin crew: experience from Sweden, **810**
Risk factors for cutaneous malignant melanoma among aircrews and a random sample of the population, **815**
- cost-benefit analysis**, The economic burden of pneumoconiosis in China, **383**
- cotton dust**, A longitudinal observation of early pulmonary responses to cotton dust, **115**
Respiratory symptoms and cotton dust exposure; results of a 15 year follow up observation, **935**
- crocidolite**, Asbestos related diseases from environmental exposure to crocidolite in Da-yao, China. I. Review of exposure and epidemiological data, **35**
- crystalline silica dust**, Chronic obstructive pulmonary disease due to occupational exposure to silica dust: a review of epidemiological and pathological evidence, **237**
- cyanide poisoning**, Alternative methods of administering amyl nitrite to victims of cyanide poisoning, **147**
- cytokines**, Repeated daily exposure to 2 ppm nitrogen dioxide upregulates the expression of IL-5, IL-10, IL-13, and ICAM-1 in the bronchial epithelium of healthy human airways, **892**
- dairy**, Animal production and wheeze in the Agricultural Health Study: interactions with atopy, asthma, and smoking, **e3**
- death**, Coverage of work related fatalities in Australia by compensation and occupational health and safety agencies, **195**
- decompression sickness**, Evaluation of decompression safety in an occupational diving group using self reported diving exposure and health status, **418**
- dehydration**, Fluid losses and hydration status of industrial workers under thermal stress working extended shifts, **90**
- demand-control model**, Does psychological distress influence reporting of demands and control at work?, **887**
- depression**, Mental health of British farmers, **181**
Workplace bullying and the risk of cardiovascular disease and depression, **779**
- dermal exposure**, Exposure related mutagens in urine of rubber workers associated with inhalable particulate and dermal exposure, **97**
- diabetes**, Fatigue in employees with diabetes: its relation with work characteristics and diabetes related burden, **i93**
- diesel exhaust**, Lung cancer in heavy equipment operators and truck drivers with diesel exhaust exposure in the construction industry, **516**
- differential misclassification**, Does psychological distress influence reporting of demands and control at work?, **887**
- diisocyanate**, Mortality results for polyurethane manufacture understated, **459**
- dimethylaminoethanol**, Visual and ocular changes associated with exposure to two tertiary amines, **69**
- dinitrotoluene**, Exposure to nitroaromatic explosives and health effects during disposal of military waste, **483**
- dioxin**, Cancer risk for chemical workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin, **672**
Risk of soft tissue sarcomas and residence in the neighbourhood of an incinerator of industrial wastes, **680**
- direct costs**, The economic burden of pneumoconiosis in China, **383**
- disability**, The right treatment to the right patient at the right time, **235**
- disease mapping**, Lymphohaematopoietic system cancer incidence in an urban area near a coke oven plant: an ecological investigation, **187**
- disinfection byproducts**, Effect of trihalomethane exposure on fetal development, **173**
- diving**, Should computed chest tomography be recommended in the medical certification of professional divers? A report of three cases with pulmonary air cysts, **606**
- DNA damage**, Urine mutagenicity and lymphocyte DNA damage in fruit growers occupationally exposed to the fungicide captan, **910**
- DNA protein crosslinks**, DNA-protein crosslinks and p53 protein expression in relation to occupational exposure to formaldehyde, **403**

- DNT**, Exposure to nitroaromatic explosives and health effects during disposal of military waste, **483**
- dose-response**, Investigating the dose-response relation between air pollution and total mortality in the APHEA-2 multicity project, **977**
Lung cancer mortality among chromate production workers, **451**
- downsizing**, Downsizing of staff is associated with lowered medically certified sick leave in female employees, **e9**
- drinking water**, Effect of trihalomethane exposure on fetal development, **173**
- driving**, Driving impairment due to sleepiness is exacerbated by low alcohol intake, **689**
- drug**, The right treatment to the right patient at the right time, **235**
- duration**, Dose-response patterns for vibration-induced white finger, **16**
- dust**, Inflammatory potential of dust from schools and building related symptoms, **e5**
Respiratory effects of volcanic emissions, **529**
- ecological study**, Lymphohaematopoietic system cancer incidence in an urban area near a coke oven plant: an ecological investigation, **187**
- economic burden**, The economic burden of pneumoconiosis in China, **383**
- elderly**, Risk factors among elderly for short term deaths related to high levels of air pollution, **684**
- electromagnetic fields**, Occupational exposure to electromagnetic fields and acute leukaemia: analysis of a case-control study, **577**
- electroplating workers**, Chromium (VI) induced oxidative damage to DNA: increase of urinary 8-hydroxydeoxyguanosine concentrations (8-OHdG) among electroplating workers, **590**
- emergency unit**, Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery, **373**
- emphysema**, Chronic obstructive pulmonary disease due to occupational exposure to silica dust: a review of epidemiological and pathological evidence, **237**
- employment status**, From insecure to secure employment: changes in work, health, health related behaviours, and sickness absence, **948**
- encephalopathy**, Competence at the workplace, **225**
- endocrine disrupter**, Epidemiology, biology, and endocrine disrupters, **541**
Risk of hypospadias in relation to maternal occupational exposure to potential endocrine disrupting chemicals, **543**
- endotoxin**, Respiratory symptoms and cotton dust exposure; results of a 15 year follow up observation, **935**
Will sewage workers with endotoxin related symptoms have the benefit of reduced lung cancer?, **144**
- environment**, How much does the environment contribute to cancer?, **150**
- environmental cadmium exposure**, Associations of lead biomarkers with renal function in Korean lead workers, **551**
- environmental exposition**, Breast cancer and serum organochlorine residues, **348**
- environmental exposure**, Cancer risk around the nuclear power plants of Trillo and Zorita (Spain), **521**
Endotoxin: is it an environmental factor in the cause of Parkinson's disease?, **378**
Gas stove use and respiratory health among adults with asthma in NHANES III, **759**
Occupational exposures and lung cancer in New Caledonia, **584**
- environmental health**, Evaluation of decompression safety in an occupational diving group using self reported diving exposure and health status, **418**
How important is personal exposure assessment in the epidemiology of air pollutants?, **143**
- environmental lead exposure**, Yearly measurements of blood lead in Swedish children since 1978: an update focusing on the petrol lead free period 1995–2001, **370**
- environmental monitoring**, Exposure assessment of monoterpenes and styrene: a comparison of air sampling and biomonitoring, **599**
Follow up study of haematological effects in workers exposed to 2-methoxyethanol, **130**
Haematological effects among silk screening workers exposed to 2-ethoxy ethyl acetate, **e7**
Occupational exposure of midwives to nitrous oxide on delivery suites, **958**
- enzymes**, Prevalence of sensitisation to cellulase and xylanase in bakery workers, **802**
- epidemiological research**, An epidemiological approach to study fatigue in the working population: the Maastricht Cohort Study, **132**
- epidemiology**, Knee disorders in the general population and their relation to occupation, **794**
Lung cancer risk in workers exposed to poly (vinyl chloride) dust: a nested case-referent study, **423**
Personal exposure assessment in the epidemiology of air pollutants, **458**
Primary osteoarthritis and occupations: a national cross sectional survey of 10 412 symptomatic patients, **882**
Smoking and occupation from the European Community Respiratory Health Survey, **643**
Update of predictions of mortality from pleural mesothelioma in the Netherlands, **50**
- epilogue**, Epilogue, **i105**
- ergonomics**, Work related and individual predictors for incident neck pain among office employees working with video display units, **475**
- ESR (electron spin resonance)**, Temporal variation of hydroxyl radical generation and 8-hydroxy-2'-deoxyguanosine formation by coarse and fine particulate matter, **315**
- estimating equations**, Generalised estimating equations and low back pain, **378**
- evaluation**, The effectiveness of current approaches to workplace stress management in the nursing profession: an evidence based literature review, **10**
- evidence based medicine**, Workplace counselling, **899**
- evidence based practice**, The effectiveness of current approaches to workplace stress management in the nursing profession: an evidence based literature review, **10**
- exercise**, Radiographic (ILO) readings predict arterial oxygen desaturation during exercise in subjects with asbestosis, **201**
- exercise induced bronchoconstriction**, Volcanic ash and respiratory symptoms in children on the island of Montserrat, British West Indies, **207**
- exercise test**, Cardiorespiratory fitness and risk of disability pension: a prospective population based study in Finnish men, **765**
- exhaled air**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
- expansion**, Downsizing of staff is associated with lowered medically certified sick leave in female employees, **e9**
- exposure assessment**, Effects of measurement strategy and statistical analysis on dose-response relations between physical workload and low back pain, **942**
Urine mutagenicity and lymphocyte DNA damage in fruit growers occupationally exposed to the fungicide captan, **910**
- exposure estimation**, Lymphohaematopoietic cancer mortality among workers with benzene exposure, **676**
- exposure-response**, Exposure-response relations among laboratory animal workers exposed to rats, **104**
Quantitative relations between exposure to respirable quartz and risk of silicosis, **159**
- extreme forward bending**, Occupational risk factors for symptomatic lumbar disc herniation; a case-control study, **821**
- falls**, Individual characteristics in occupational accidents due to imbalance: a case-control study in the employees of a railway company, **330**
- farming**, Animal production and wheeze in the Agricultural Health Study: interactions with atopy, asthma, and smoking, **e3**
Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men, **e11**
Mental health of British farmers, **181**
Respiratory status in dairy farmers in France; cross sectional and longitudinal analyses, **858**
- fatigue**, Acute and chronic job stressors among ambulance personnel: predictors of health symptoms, **i40**
An epidemiological approach to study fatigue in the working population: the Maastricht Cohort Study, **i32**
Assessment of fatigue among working people: a comparison of six questionnaires, **i10**
Epilogue, **i105**
Fatigue as a predictor of sickness absence: results from the Maastricht cohort study on fatigue at work, **i71**
Fatigue as a risk factor for being injured in an occupational accident: results from the Maastricht Cohort Study, **i88**
Fatigue, burnout, and chronic fatigue syndrome among employees on sick leave: do attributions make the difference?, **i26**
Fatigue in employees with diabetes: its relation with work characteristics and diabetes related burden, **i93**

- Measurement quality and validity of the "need for recovery scale", **i3**
 Need for recovery after work and the subsequent risk of cardiovascular disease in a working population, **i83**
 Need for recovery from work related fatigue and its role in the development and prediction of subjective health complaints, **i62**
 Work schedules and fatigue: a prospective cohort study, **i47**
fatigue attribution, Associations between fatigue attributions and fatigue, health, and psychosocial work characteristics: a study among employees visiting a physician with fatigue, **i99**
fetal development, Effect of trihalomethane exposure on fetal development, **173**
fibrinogen, Ischaemic heart disease among livestock and agricultural workers, **e1**
finger systolic blood pressure, Skin temperature recovery from cold provocation in workers exposed to vibration: a longitudinal study, **962**
fixed-term employment, From insecure to secure employment: changes in work, health, health related behaviours, and sickness absence, **948**
flight attendants, Prevalence of respiratory symptoms among female flight attendants and teachers, **929**
flower, Occupational asthma due to IgE mediated allergy to the flower *Moluccella laevis* (Bells of Ireland), **701**
fluazinam, Occupational asthma from fungicides fluazinam and chlorothalonil, **76**
fluid replacement, Fluid losses and hydration status of industrial workers under thermal stress working extended shifts, **90**
fluorides, World at work: Hazards and controls in aluminium potrooms, **989**
fluorochemicals, Mortality of employees of a perfluorooctanesulphonyl fluoride manufacturing facility, **722**
follow up studies, Under-ascertainment of multiple myeloma among participants in UK atmospheric atomic and nuclear weapons tests, **e18**
follow up study, Prognosis of shoulder tendonitis in repetitive work: a follow up study in a cohort of Danish industrial and service workers, **1**
forearm pain, Does computer use pose an occupational hazard for forearm pain; from the NUDATA study, **e14**
formaldehyde, Prevalence of respiratory symptoms in sheep breeders, **770**
formaldehyde exposure, DNA-protein crosslinks and p53 protein expression in relation to occupational exposure to formaldehyde, **403**
free radical, Oxidative stress: its role in air pollution and adverse health effects, **612**
functional abilities, Increased occupational physical activity does not improve physical fitness, **983**
fungicide, Occupational asthma from fungicides fluazinam and chlorothalonil, **76**
gas chromatography, Breast cancer and serum organochlorine residues, **348**
Gaza, Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery, **373**
general practitioner, Associations between fatigue attributions and fatigue, health, and psychosocial work characteristics: a study among employees visiting a physician with fatigue, **i99**
 Health problems and psychosocial work environment as predictors of long term sickness absence in employees who visited the occupational physician and/or general practitioner in relation to work: a prospective study, **295**
genetic biomarkers, Surveillance in occupational health, **705**
genetic susceptibility, Glutathione S-transferases M1-1 and T1-1 as risk modifiers for renal cell cancer associated with occupational exposure to chemicals, **789**
glutaraldehyde, Physician diagnosed asthma, respiratory symptoms, and associations with workplace tasks among radiographers in Ontario, Canada, **254**
good medical practice, Good medical practice for occupational physicians, **1**
groundwater contamination, Increased mortality odds ratio of male liver cancer in a community contaminated by chlorinated hydrocarbons in groundwater, **364**
guano, Potential effects on human health of an ammonia rich atmospheric environment in an archaeologically important cave in southeast Asia, **986**
guidelines, Occupational health guidelines for the management of low back pain: an international comparison, **618**
 The management of low back pain, **617**
H2S, Mortality and cancer morbidity in a cohort of Canadian petroleum workers, **918**
haematological effects, Follow up study of haematological effects in workers exposed to 2-methoxyethanol, **130**
haematological toxicity, Haematological effects among silk screening workers exposed to 2-ethoxy ethyl acetate, **e7**
hairdressing apprentices, Healthy worker effect and changes in respiratory symptoms and lung function in hairdressing apprentices, **831**
hand arm vibration syndrome, Hand-arm vibration syndrome in Swedish car mechanics, **287**
 Skin temperature recovery from cold provocation in workers exposed to vibration: a longitudinal study, **962**
hazard, Surveillance in occupational health, **705**
hazardous waste, Hazardous waste landfill sites and congenital anomalies, **79**
health, Organisational justice and health of employees: prospective cohort study, **27**
health complaint, Need for recovery from work related fatigue and its role in the development and prediction of subjective health complaints, **i62**
health effects, Use of personal exposure modelling in risk assessment of air pollutants, **529**
health surveillance, A mortality and morbidity study of refinery and petrochemical employees in Louisiana, **627**
 Surveillance in occupational health, **705**
healthcare personnel, From insecure to secure employment: changes in work, health, health related behaviours, and sickness absence, **948**
healthy worker effect, Mortality results for polyurethane manufacture understated, **459**
hearing loss, Agreement between hearing thresholds measured in non-soundproof work environments and a soundproof booth, **667**
 Combined effect of smoking and occupational exposure to noise on hearing loss in steel factory workers, **56**
hearing threshold, Agreement between hearing thresholds measured in non-soundproof work environments and a soundproof booth, **667**
heart rate, Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
heat stress, Fluid losses and hydration status of industrial workers under thermal stress working extended shifts, **90**
hepatitis virus infection, Synergistic effect of hepatitis virus infection and occupational exposures to vinyl chloride monomer and ethylene dichloride on serum aminotransferase activity, **774**
hepatomegaly, Periportal fibrosis and other liver ultrasonography findings in vinyl chloride workers, **60**
hexachlorobutadiene, Land contamination and urinary abnormalities: cause for concern?, **463**
hexavalent chromium, Lung cancer mortality among chromate production workers, **451**
hierarchical regression, Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men, **e11**
histopathology, Clinical and histopathological changes of the nasal mucosa induced by occupational exposure to sulphuric acid mists, **395**
Hong Kong, An outbreak of SARS among healthcare workers, **528**
household waste, Upper airway inflammation in waste handlers exposed to bioaerosols, **444**
HPA axis, Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
human health, Potential effects on human health of an ammonia rich atmospheric environment in an archaeologically important cave in southeast Asia, **986**
hydrocarbon lubricants, Mortality and cancer morbidity in a cohort of Canadian petroleum workers, **918**
hydrocarbon solvents and fuels, Mortality and cancer morbidity in a cohort of Canadian petroleum workers, **918**
hydrogen sulfide, Hydrogen sulfide: UK occupational exposure limits, **308**
hypohydration, Fluid losses and hydration status of industrial workers under thermal stress working extended shifts, **90**
hypospadias, Epidemiology, biology, and endocrine disrupters, **541**

- Risk of hypospadias in relation to maternal occupational exposure to potential endocrine disrupting chemicals, **543**
- ICAM-1**, Repeated daily exposure to 2 ppm nitrogen dioxide upregulates the expression of IL-5, IL-10, IL-13, and ICAM-1 in the bronchial epithelium of healthy human airways, **892**
- IHD**, Ischaemic heart disease among livestock and agricultural workers, **e1**
- illness**, Mental ill health in workers: observations from a few Indian populations, **225**
- importation**, The right treatment to the right patient at the right time, **235**
- incidence**, Hand-arm vibration syndrome in Swedish car mechanics, **287**
- Multiple sclerosis in nurse anaesthetists, **66**
- Reported incidence of occupational asthma in France, 1996–99: the ONAP programme, **136**
- Work related and individual predictors for incident neck pain among office employees working with video display units, **475**
- incinerator**, Risk of soft tissue sarcomas and residence in the neighbourhood of an incinerator of industrial wastes, **680**
- India**, Mental ill health in workers: observations from a few Indian populations, **225**
- indirect costs**, The economic burden of pneumoconiosis in China, **383**
- individual characteristics**, Individual characteristics in occupational accidents due to imbalance: a case-control study in the employees of a railway company, **330**
- infection**, Do junior doctors take sick leave?, **699**
- inflammation**, Ischaemic heart disease among livestock and agricultural workers, **e1**
- inflammatory potential**, Inflammatory potential of dust from schools and building related symptoms, **e5**
- inorganic lead**, Can salivary lead be used for biological monitoring of lead exposed individuals?, **696**
- inorganic substances**, Municipal waste incinerators: air and biological monitoring of workers for exposure to particles, metals, and organic compounds, **563**
- intervention**, Intervention strategies to reduce musculoskeletal injuries associated with handling patients: a systematic review, **e6**
- intrauterine growth retardation**, Effect of trihalomethane exposure on fetal development, **173**
- Inuit**, Monitoring of umbilical cord blood lead levels and sources assessment among the Inuit, **693**
- Iran**, No change in sex ratio in Ramsar (north of Iran) with high background of radiation, **146**
- ischaemic heart disease**, Ischaemic heart disease among livestock and agricultural workers, **e1**
- jet fuel**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
- job analysis**, Does psychological distress influence reporting of demands and control at work?, **887**
- job insecurity**, From insecure to secure employment: changes in work, health, health related behaviours, and sickness absence, **948**
- job satisfaction**, Sickness absence due to low back pain, **306**
- job stress**, Downsizing of staff is associated with lowered medically certified sick leave in female employees, **e9**
- Need for recovery after work and the subsequent risk of cardiovascular disease in a working population, **i83**
- job-exposure matrix**, Evaluation of the quality of coding of job episodes collected by self questionnaires among French retired men for use in a job-exposure matrix, **438**
- JP-8**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
- junior doctors**, Do junior doctors take sick leave?, **699**
- Karasek's model**, Psychosocial factors at work and self reported health: comparative results of cross sectional and prospective analyses of the French GAZEL cohort, **509**
- kidney cancer**, Glutathione S-transferases M1-1 and T1-1 as risk modifiers for renal cell cancer associated with occupational exposure to chemicals, **789**
- kidney function**, Associations of lead biomarkers with renal function in Korean lead workers, **551**
- laboratory animal**, Estimating the incidence of occupational asthma and rhinitis from laboratory animal allergens in the UK, 1999–2000, **604**
- laboratory animal allergy**, Exposure-response relations among laboratory animal workers exposed to rats, **104**
- land contamination**, Land contamination and renal dysfunction, **461**
- landfill**, Hazardous waste landfill sites and congenital anomalies, **79**
- lateral epicondylitis**, Physical and psychosocial risk factors for lateral epicondylitis: a population based case-referent study, **322**
- lead**, Monitoring of umbilical cord blood lead levels and sources assessment among the Inuit, **693**
- Neurobehavioural testing in workers occupationally exposed to lead, **145**
- lead exposure**, Time To Pregnancy and occupational lead exposure, **752**
- leukaemia**, Lymphohaematopoietic cancer mortality among workers with benzene exposure, **676**
- Lymphohaematopoietic system cancer incidence in an urban area near a coke oven plant: an ecological investigation, **187**
- lifting**, The relative importance of whole body vibration and occupational lifting as risk factors for low-back pain, **715**
- liver cancer**, Increased mortality odds ratio of male liver cancer in a community contaminated by chlorinated hydrocarbons in groundwater, **364**
- Wastewater exposure and health—a comparative study of two occupational groups, **595**
- liver ultrasonography**, Periportal fibrosis and other liver ultrasonography findings in vinyl chloride workers, **60**
- livestock workers**, Ischaemic heart disease among livestock and agricultural workers, **e1**
- Prevalence of respiratory symptoms in sheep breeders, **770**
- London**, William Harvey and air pollution, **147**
- long term sickness absence**, Reducing long term sickness absence by an activating intervention in adjustment disorders: a cluster randomised controlled design, **429**
- longitudinal study**, Course of low back pain among nurses: a longitudinal study across eight years, **497**
- Healthy worker effect and changes in respiratory symptoms and lung function in hairdressing apprentices, **831**
- High local unemployment and increased mortality in Danish adults: results from a prospective multilevel study, **e16**
- Psychosocial conditions on and off the job and psychological ill health: depressive symptoms, impaired psychological wellbeing, heavy consumption of alcohol, **489**
- longitudinal study of respiratory function**, Respiratory status in dairy farmers in France; cross sectional and longitudinal analyses, **858**
- lost time claims**, Age related differences in work injuries and permanent impairment: a comparison of workers' compensation claims among adolescents, young adults, and adults, **e10**
- low back pain**, Course of low back pain among nurses: a longitudinal study across eight years, **497**
- Occupational health guidelines for the management of low back pain: an international comparison, **618**
- LPS**, Endotoxin: is it an environmental factor in the cause of Parkinson's disease?, **378**
- lumbar disc herniation**, Occupational risk factors for symptomatic lumbar disc herniation; a case-control study, **821**
- lung cancer**, Asbestos related diseases from environmental exposure to crocidolite in Da-yao, China. I. Review of exposure and epidemiological data, **35**
- Lung cancer mortality among chromate production workers, **451**
- Lung cancer risk in workers exposed to poly (vinyl chloride) dust: a nested case-referent study, **423**
- Musicians playing wind instruments and risk of lung cancer: is there an association?, **143**
- Occupational exposures and lung cancer in New Caledonia, **584**
- Occupational silica exposure and risk of various diseases: an analysis using death certificates from 27 states of the United States, **122**
- The biological effects of coarse and fine particulate matter, **313**
- Will sewage workers with endotoxin related symptoms have the benefit of reduced lung cancer?, **144**
- lung epithelium permeability**, Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools, **385**
- lung function**, Healthy worker effect and changes in respiratory symptoms and lung function in hairdressing apprentices, **831**
- lung lining fluid**, Oxidative stress: its role in air pollution and adverse health effects, **612**
- lymphoma**, Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men, **e11**
- Lymphohaematopoietic system cancer incidence in an urban area near a coke oven plant: an ecological investigation, **187**

- magnetic field**, Residential and occupational exposure to 50 Hz magnetic fields and malignant melanoma: a population based study, **343**
- magnitude**, Dose-response patterns for vibration-induced white finger, **16**
- Malaysia**, Relation between sick leave and selected exposure variables among women semiconductor workers in Malaysia, **262**
- male reproduction**, Time To Pregnancy and occupational lead exposure, **752**
- malignant melanoma**, Residential and occupational exposure to 50 Hz magnetic fields and malignant melanoma: a population based study, **343**
- manual handling**, Intervention strategies to reduce musculoskeletal injuries associated with handling patients: a systematic review, **e6**
- measurement**, Assessment of fatigue among working people: a comparison of six questionnaires, **i10**
Measurement quality and validity of the "need for recovery scale", **i3**
- meatworkers**, Mortality and cancer incidence in a cohort of meatworkers, **e4**
- mechanic**, Hand-arm vibration syndrome in Swedish car mechanics, **287**
- mechanical factors**, Mechanical and psychosocial factors predict new onset shoulder pain: a prospective cohort study of newly employed workers, **850**
- mechanisms**, The effect of sulphurous air pollutant exposures on symptoms, lung function, exhaled nitric oxide, and nasal epithelial lining fluid antioxidant concentrations in normal and asthmatic adults, **e15**
- memory**, Memory one month after acute carbon monoxide intoxication: a prospective study, **212**
- mental health**, Ambient neighbourhood noise and children's mental health, **146**
Mental ill health in workers: observations from a few Indian populations, **225**
Quality of rehabilitation among workers with adjustment disorders according to practice guidelines: a retrospective cohort study, **i21**
The Depression Anxiety Stress Scales (DASS): detecting anxiety disorder and depression in employees absent from work because of mental health problems, **i77**
- mesothelioma**, Asbestos related diseases from environmental exposure to crocidolite in Da-yao, China. I. Review of exposure and epidemiological data, **35**
Predictions of mortality from mesothelioma, **458**
Update of predictions of mortality from pleural mesothelioma in the Netherlands, **50**
- meta-analysis**, Occupation related pesticide exposure and cancer of the prostate: a meta-analysis, **634**
- metal fume fever**, Prevalence and association of welding related systemic and respiratory symptoms in welders, **655**
- methodology**, How to conduct research on burnout: advantages and disadvantages of a unidimensional approach in burnout research, **i16**
- methylene chloride**, Mortality results for polyurethane manufacture understated, **459**
- microenvironment air sampling**, Are personal and static samples related?, **224**
- midwifery**, Occupational exposure of midwives to nitrous oxide on delivery suites, **958**
- milk proteins**, Dermatological powder as hidden cause of occupational allergy due to casein: a case report, **609**
- misclassification**, Evaluation of the quality of coding of job episodes collected by self questionnaires among French retired men for use in a job-exposure matrix, **438**
- monoterpene**, Exposure assessment of monoterpenes and styrene: a comparison of air sampling and biomonitoring, **599**
- MOR**, Increased mortality odds ratio of male liver cancer in a community contaminated by chlorinated hydrocarbons in groundwater, **364**
- morbidity**, A mortality and morbidity study of refinery and petrochemical employees in Louisiana, **627**
- mortality**, A mortality and morbidity study of refinery and petrochemical employees in Louisiana, **627**
Follow up of mortality and incidence of cancer 1952–98 in men from the UK who participated in the UK's atmospheric nuclear weapon tests and experimental programmes, **165**
High local unemployment and increased mortality in Danish adults: results from a prospective multilevel study, **e16**
Mortality and cancer incidence in a cohort of meatworkers, **e4**
- Mortality and cancer morbidity in a cohort of Canadian petroleum workers, **918**
- Mortality in workers in poultry slaughtering/processing plants: the Missouri poultry cohort study, **784**
- Mortality of employees of a perfluorooctanesulphonyl fluoride manufacturing facility, **722**
- Mortality of petroleum refinery workers, **304**
- Mortality results for polyurethane manufacture understated, **459**
- Predictions of mortality from mesothelioma, **458**
- Risk factors among elderly for short term deaths related to high levels of air pollution, **684**
- William Harvey and air pollution, **147**
- motor abilities**, Increased occupational physical activity does not improve physical fitness, **983**
- mouse device**, Does computer use pose an occupational hazard for forearm pain; from the NUDATA study, **e14**
- mucous membrane irritation**, Effects of bioaerosol polluted outdoor air on airways of residents: a cross sectional study, **336**
- multifactorial**, Physical and psychosocial risk factors for lateral epicondylitis: a population based case-referent study, **322**
- multiple myeloma**, Lymphohaematopoietic cancer mortality among workers with benzene exposure, **676**
Under-ascertainment of multiple myeloma among participants in UK atmospheric atomic and nuclear weapons tests, **e18**
- multiple sclerosis**, Multiple sclerosis in nurse anaesthetists, **66**
- musculoskeletal disease**, Early prognostic factors for duration on temporary total benefits in the first year among workers with compensated occupational soft tissue injuries, **244**
- musculoskeletal disorder**, Physical and psychosocial risk factors for lateral epicondylitis: a population based case-referent study, **322**
- musculoskeletal injuries**, Intervention strategies to reduce musculoskeletal injuries associated with handling patients: a systematic review, **e6**
- N-acetyl-β-D-glucosaminidase**, Associations of lead biomarkers with renal function in Korean lead workers, **551**
- naphthalene**, Benzene and naphthalene in air and breath as indicators of exposure to jet fuel, **969**
- nasal lavage**, Upper airway inflammation in waste handlers exposed to bioaerosols, **444**
- nasal mucosa**, Clinical and histopathological changes of the nasal mucosa induced by occupational exposure to sulphuric acid mists, **395**
- NAT2-phenotype**, Exposure related mutagens in urine of rubber workers associated with inhalable particulate and dermal exposure, **97**
- neck**, Risk factors in the onset of neck/shoulder pain in a prospective study of workers in industrial and service companies, **649**
- neck pain**, Risk factors for incident neck and shoulder pain in hospital nurses, **864**
Work related and individual predictors for incident neck pain among office employees working with video display units, **475**
- need for recovery**, Need for recovery after work and the subsequent risk of cardiovascular disease in a working population, **i83**
- neoplasia**, Availability of thyroid protective lead shields and their use by trainee orthopaedic surgeons, **381**
- neoplasm**, Cancer risk around the nuclear power plants of Trillo and Zorita (Spain), **521**
- nerve entrapment**, Does computer use pose an occupational hazard for forearm pain; from the NUDATA study, **e14**
- neurobehavioural deficit**, Neurobehavioural effects among workers occupationally exposed to organophosphorous pesticides, **279**
- neurobehavioural performance**, The impact of a week of simulated night work on sleep, circadian phase, and performance, **e13**
- neurobehavioural testing**, Neurobehavioural tests and systems to assess neurotoxic exposures in the workplace and community, **531**
Neurobehavioural testing in workers occupationally exposed to lead, **145**
- neurological symptom**, Hand-arm vibration syndrome in Swedish car mechanics, **287**
- neurotoxic exposures**, Neurobehavioural tests and systems to assess neurotoxic exposures in the workplace and community, **531**
- neurotoxicity**, Relation between colour vision loss and occupational styrene exposure, **222**
- new onset asthma**, Physician diagnosed asthma, respiratory symptoms, and associations with workplace tasks among radiographers in Ontario, Canada, **254**

- new workers**, Mechanical and psychosocial factors predict new onset shoulder pain: a prospective cohort study of newly employed workers, **850**
- newborn**, Monitoring of umbilical cord blood lead levels and sources assessment among the Inuit, **693**
Organochlorine compounds and concentrations of thyroid stimulating hormone in newborns, **301**
- Niah**, Potential effects on human health of an ammonia rich atmospheric environment in an archaeologically important cave in southeast Asia, **986**
- nickel**, Occupational contact dermatitis to nickel: experience of the British dermatologists (EPIDERM) and occupational physicians (OPRA) surveillance schemes, **954**
- nitrogen dioxide**, Are personal and static samples related?, **224**
Repeated daily exposure to 2 ppm nitrogen dioxide upregulates the expression of IL-5, IL-10, IL-13, and ICAM-1 in the bronchial epithelium of healthy human airways, **892**
- nitrogen trichloride**, Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools, **385**
- nitrous oxide**, Occupational exposure of midwives to nitrous oxide on delivery suites, **958**
- N,N-dimethylacetamide**, Biological monitoring of occupational exposure to N,N-dimethylacetamide with identification of a new metabolite, **746**
- NO₂**, Investigating the dose-response relation between air pollution and total mortality in the APHEA-2 multicity project, **977**
NO₂ as a marker of air pollution, and recurrent wheezing in children: a nested case-control study within the BAMSE birth cohort, **876**
- noise**, Agreement between hearing thresholds measured in non-soundproof work environments and a soundproof booth, **667**
Ambient neighbourhood noise and children's mental health, **146**
- nuclear reactors**, Cancer risk around the nuclear power plants of Trillo and Zorita (Spain), **521**
- nuclear veterans**, Under-ascertainment of multiple myeloma among participants in UK atmospheric atomic and nuclear weapons tests, **e18**
- nuclear weapon tests**, Follow up of mortality and incidence of cancer 1952–98 in men from the UK who participated in the UK's atmospheric nuclear weapon tests and experimental programmes, **165**
- nurses**, Course of low back pain among nurses: a longitudinal study across eight years, **497**
Risk factors for incident neck and shoulder pain in hospital nurses, **864**
- nurses aides**, Work factors as predictors of sickness absence: a three month prospective study of nurses' aides, **271**
- nursing**, The effectiveness of current approaches to workplace stress management in the nursing profession: an evidence based literature review, **10**
- occupation**, Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men, **e11**
Knee disorders in the general population and their relation to occupation, **794**
Lung cancer in heavy equipment operators and truck drivers with diesel exhaust exposure in the construction industry, **516**
Primary osteoarthritis and occupations: a national cross sectional survey of 10 412 symptomatic patients, **882**
Risk of hypospadias in relation to maternal occupational exposure to potential endocrine disrupting chemicals, **543**
Smoking and occupation from the European Community Respiratory Health Survey, **643**
- occupational accidents**, Fatigue as a risk factor for being injured in an occupational accident: results from the Maastricht Cohort Study, **i88**
Individual characteristics in occupational accidents due to imbalance: a case-control study in the employees of a railway company, **330**
- occupational allergy**, Dermatological powder as hidden cause of occupational allergy due to casein: a case report, **609**
Occupational asthma due to IgE mediated allergy to the flower *Molucella laevis* (Bells of Ireland), **701**
- occupational asthma**, Occupational asthma, **711**
Occupational asthma from fungicides fluzinam and chlorothalonil, **76**
Reported incidence of occupational asthma in France, 1996–99: the ONAP programme, **136**
The continuing challenge to reduce the burden of occupational asthma, **713**
- occupational chemical exposure**, Synergistic effect of hepatitis virus infection and occupational exposures to vinyl chloride monomer and ethylene dichloride on serum aminotransferase activity, **774**
- occupational cohort**, A mortality and morbidity study of refinery and petrochemical employees in Louisiana, **627**
- occupational contact dermatitis**, Occupational contact dermatitis to nickel: experience of the British dermatologists (EPIDERM) and occupational physicians (OPRA) surveillance schemes, **954**
- occupational disease**, Risk factors for incident neck and shoulder pain in hospital nurses, **864**
- occupational epidemiology**, Musicians playing wind instruments and risk of lung cancer: is there an association?, **143**
- Occupational exposure**, Breast cancer risk in airline cabin attendants: a nested case-control study in Iceland, **807**
Glutathione S-transferases M1-1 and T1-1 as risk modifiers for renal cell cancer associated with occupational exposure to chemicals, **789**
Municipal waste incinerators: air and biological monitoring of workers for exposure to particles, metals, and organic compounds, **563**
Occupational exposure of midwives to nitrous oxide on delivery suites, **958**
Occupational exposures and lung cancer in New Caledonia, **584**
Respiratory symptoms and cotton dust exposure; results of a 15 year follow up observation, **935**
- occupational exposure limits**, Hydrogen sulfide: UK occupational exposure limits, **308**
- occupational health**, Endotoxin: is it an environmental factor in the cause of Parkinson's disease?, **378**
Epilogue, **i105**
From insecure to secure employment: changes in work, health, health related behaviours, and sickness absence, **948**
Occupational health guidelines for the management of low back pain: an international comparison, **618**
Workplace bullying and the risk of cardiovascular disease and depression, **779**
- occupational health services**, Quality of rehabilitation among workers with adjustment disorders according to practice guidelines; a retrospective cohort study, **i21**
The Depression Anxiety Stress Scales (DASS): detecting anxiety disorder and depression in employees absent from work because of mental health problems, **i77**
- occupational injury**, Early prognostic factors for duration on temporary total benefits in the first year among workers with compensated occupational soft tissue injuries, **244**
- occupational lead exposure**, Associations of lead biomarkers with renal function in Korean lead workers, **551**
- occupational physicians**, Associations between fatigue attributions and fatigue, health, and psychosocial work characteristics: a study among employees visiting a physician with fatigue, **i99**
Good medical practice for occupational physicians, **1**
Health problems and psychosocial work environment as predictors of long term sickness absence in employees who visited the occupational physician and/or general practitioner in relation to work: a prospective study, **295**
Quality of rehabilitation among workers with adjustment disorders according to practice guidelines; a retrospective cohort study, **i21**
- occupational rehabilitation**, The Depression Anxiety Stress Scales (DASS): detecting anxiety disorder and depression in employees absent from work because of mental health problems, **i77**
- occupational risk**, Occupations with increased risk of pancreatic cancer in the Swedish population, **570**
The relative importance of whole body vibration and occupational lifting as risk factors for low-back pain, **715**
- occupational stress**, How to conduct research on burnout: advantages and disadvantages of a unidimensional approach in burnout research, **i16**
The effectiveness of current approaches to workplace stress management in the nursing profession: an evidence based literature review, **10**
- odour annoyance**, Effects of bioaerosol polluted outdoor air on airways of residents: a cross sectional study, **336**
- ODTS**, Prevalence of respiratory symptoms in sheep breeders, **770**
- official statistics**, Coverage of work related fatalities in Australia by compensation and occupational health and safety agencies, **195**
- oncogenic**, Mortality in workers in poultry slaughtering/processing plants: the Missouri poultry cohort study, **784**

- organic dust**, Effects of bioaerosol polluted outdoor air on airways of residents: a cross sectional study, **336**
- organic solvent**, Relation between colour vision loss and occupational styrene exposure, **222**
- organic substances**, Municipal waste incinerators: air and biological monitoring of workers for exposure to particles, metals, and organic compounds, **563**
- organisational justice**, Organisational justice and health of employees: prospective cohort study, **27**
- organochlorine compound**, Organochlorine compounds and concentrations of thyroid stimulating hormone in newborns, **301**
- organochlorine residues**, Breast cancer and serum organochlorine residues, **348**
- orthopaedic**, Availability of thyroid protective lead shields and their use by trainee orthopaedic surgeons, **381**
- osteoarthritis**, Primary osteoarthritis and occupations: a national cross sectional survey of 10 412 symptomatic patients, **882**
- oxidative stress**, Chromium (VI) induced oxidative damage to DNA: increase of urinary 8-hydroxydeoxyguanosine concentrations (8-OHdG) among electroplating workers, **590**
- oxygen saturation**, Respiratory status in dairy farmers in France; cross sectional and longitudinal analyses, **858**
- oxygen uptake**, Cardiorespiratory fitness and risk of disability pension: a prospective population based study in Finnish men, **765**
- oxygenation**, Earlier study on asbestos workers, ILO scores, and oxygenation more comprehensive, **611**
- p53 protein**, DNA-protein crosslinks and p53 protein expression in relation to occupational exposure to formaldehyde, **403**
- pain**, Pain tolerance in upper limb disorders: findings from a community survey, **217**
- pancreatic cancer**, Occupations with increased risk of pancreatic cancer in the Swedish population, **570**
- panel study**, Acute effects of winter air pollution on respiratory function in schoolchildren in southern England, **82**
- paramedic personnel**, Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery, **373**
- parental occupation**, Parental occupation at periconception: findings from the United Kingdom Childhood Cancer Study, **901**
- particulate exposure**, Exposure related mutagens in urine of rubber workers associated with inhalable particulate and dermal exposure, **97**
- particulate matter**, The biological effects of coarse and fine particulate matter, **313**
- passive smoking**, Yearly measurements of blood lead in Swedish children since 1978: an update focusing on the petrol lead free period 1995–2001, **370**
- peak expiratory flow**, Acute effects of winter air pollution on respiratory function in schoolchildren in southern England, **82**
- peasant**, Asbestos related diseases from environmental exposure to crocidolite in Da-yao, China. I. Review of exposure and epidemiological data, **35**
- pension**, Cardiorespiratory fitness and risk of disability pension: a prospective population based study in Finnish men, **765**
- perfluorooctane sulphonate**, Mortality of employees of a perfluorooctanesulphonyl fluoride manufacturing facility, **722**
- performance**, Increased occupational physical activity does not improve physical fitness, **983**
- periportal fibrosis**, Periportal fibrosis and other liver ultrasonography findings in vinyl chloride workers, **60**
- permanent impairment**, Age related differences in work injuries and permanent impairment: a comparison of workers' compensation claims among adolescents, young adults, and adults, **e10**
- personal exposure**, Personal exposure assessment in the epidemiology of air pollutants, **458**
- personal exposure assessment**, How important is personal exposure assessment in the epidemiology of air pollutants?, **143**
- pesticide**, Integrative assessment of multiple pesticides as risk factors for non-Hodgkin's lymphoma among men, **e11**
- Neurobehavioural effects among workers occupationally exposed to organophosphorous pesticides, **279**
- Occupation related pesticide exposure and cancer of the prostate: a meta-analysis, **634**
- Urine mutagenicity and lymphocyte DNA damage in fruit growers occupationally exposed to the fungicide captan, **910**
- petrochemical**, Sex ratio at birth and exposure to petrochemicals, **704**
- petroleum coke/spent catalyst**, Mortality and cancer morbidity in a cohort of Canadian petroleum workers, **918**
- petroleum refinery**, Mortality of petroleum refinery workers, **304**
- petroleum workers**, Mortality and cancer morbidity in a cohort of Canadian petroleum workers, **918**
- physical fitness**, Increased occupational physical activity does not improve physical fitness, **983**
- physical work load**, Effects of measurement strategy and statistical analysis on dose-response relations between physical workload and low back pain, **942**
- Risk factors in the onset of neck/shoulder pain in a prospective study of workers in industrial and service companies, **649**
- Sickness absence due to low back pain, **306**
- PM (particulate matter)**, Temporal variation of hydroxyl radical generation and 8-hydroxy-2'-deoxyguanosine formation by coarse and fine particulate matter, **315**
- PM10**, Acute effects of winter air pollution on respiratory function in schoolchildren in southern England, **82**
- Use of personal exposure modelling in risk assessment of air pollutants, **529**
- PM2.5**, Use of personal exposure modelling in risk assessment of air pollutants, **529**
- pneumoconiosis**, Radiographic abnormalities among construction workers exposed to quartz containing dust, **410**
- The economic burden of pneumoconiosis in China, **383**
- pneumoproteins**, Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools, **385**
- Poisson regression**, Occupations with increased risk of pancreatic cancer in the Swedish population, **570**
- policy**, Occupational asthma, **711**
- pollution**, Land contamination and renal dysfunction, **461**
- polyurethane**, Mortality results for polyurethane manufacture understated, **459**
- polyvinyl chloride**, Periportal fibrosis and other liver ultrasonography findings in vinyl chloride workers, **60**
- polyvinyl chloride dust**, Lung cancer risk in workers exposed to poly(vinyl chloride) dust: a nested case-referent study, **423**
- population**, The relative importance of whole body vibration and occupational lifting as risk factors for low-back pain, **715**
- population study**, Cardiorespiratory fitness and risk of disability pension: a prospective population based study in Finnish men, **765**
- post-traumatic stress disorder**, Acute and chronic job stressors among ambulance personnel: predictors of health symptoms, **i40**
- postural control**, Individual characteristics in occupational accidents due to imbalance: a case-control study in the employees of a railway company, **330**
- poultry**, Animal production and wheeze in the Agricultural Health Study: interactions with atopy, asthma, and smoking, **e3**
- Mortality in workers in poultry slaughtering/processing plants: the Missouri poultry cohort study, **784**
- prediction**, Update of predictions of mortality from pleural mesothelioma in the Netherlands, **50**
- pregnancy planning bias**, Shift work and subfecundity: a causal link or an artefact?, **e12**
- prevalence**, Prevalence and association of welding related systemic and respiratory symptoms in welders, **655**
- prevention**, The continuing challenge to reduce the burden of occupational asthma, **713**
- prior wake**, The impact of a week of simulated night work on sleep, circadian phase, and performance, **e13**
- professional diver**, Should computed chest tomography be recommended in the medical certification of professional divers? A report of three cases with pulmonary air cysts, **606**
- prognosis**, Early prognostic factors for duration on temporary total benefits in the first year among workers with compensated occupational soft tissue injuries, **244**
- Prognosis of shoulder tendonitis in repetitive work: a follow up study in a cohort of Danish industrial and service workers, **1**
- proportional hazards model**, Early prognostic factors for duration on temporary total benefits in the first year among workers with compensated occupational soft tissue injuries, **244**
- prospective studies**, Fatigue as a predictor of sickness absence: results from the Maastricht cohort study on fatigue at work, **i71**
- prostate cancer**, Occupation related pesticide exposure and cancer of the prostate: a meta-analysis, **634**

- psychiatric morbidity**, Mental health of British farmers, **181**
- psychological ill health**, Psychosocial conditions on and off the job and psychological ill health: depressive symptoms, impaired psychological wellbeing, heavy consumption of alcohol, **489**
Reducing work related psychological ill health and sickness absence: a systematic literature review, **3**
- psychosocial**, Risk factors for incident neck and shoulder pain in hospital nurses, **864**
- psychosocial factors**, Mechanical and psychosocial factors predict new onset shoulder pain: a prospective cohort study of newly employed workers, **850**
- psychosocial factors at work**, Psychosocial factors at work and self reported health: comparative results of cross sectional and prospective analyses of the French GAZEL cohort, **509**
- psychosocial work characteristics**, Health problems and psychosocial work environment as predictors of long term sickness absence in employees who visited the occupational physician and/or general practitioner in relation to work: a prospective study, **295**
- psychosocial workload**, Occupational risk factors for symptomatic lumbar disc herniation; a case-control study, **821**
- pulmonary barotrauma**, Should computed chest tomography be recommended in the medical certification of professional divers? A report of three cases with pulmonary air cysts, **606**
- pulmonary function**, A longitudinal observation of early pulmonary responses to cotton dust, **115**
- quality of care**, Quality of rehabilitation among workers with adjustment disorders according to practice guidelines; a retrospective cohort study, **i21**
Return to work of cancer survivors: a prospective cohort study into the quality of rehabilitation by occupational physicians, **352**
- quartz**, Quantitative relations between exposure to respirable quartz and risk of silicosis, **159**
Radiographic abnormalities among construction workers exposed to quartz containing dust, **410**
- questionnaire**, Risk factors for cutaneous malignant melanoma among aircrews and a random sample of the population, **815**
- radiation**, Follow up of mortality and incidence of cancer 1952–98 in men from the UK who participated in the UK's atmospheric nuclear weapon tests and experimental programmes, **165**
How much does the environment contribute to cancer?, **150**
- radiographer**, Physician diagnosed asthma, respiratory symptoms, and associations with workplace tasks among radiographers in Ontario, Canada, **254**
- radiology**, Radiographic (ILO) readings predict arterial oxygen desaturation during exercise in subjects with asbestosis, **201**
- randomised controlled clinical trial**, Effects of a home exercise programme on shoulder pain and functional status in construction workers, **841**
- Ransar**, No change in sex ratio in Ramsar (north of Iran) with high background of radiation, **146**
- Raynaud's phenomenon**, Dose-response patterns for vibration-induced white finger, **16**
Skin temperature recovery from cold provocation in workers exposed to vibration: a longitudinal study, **962**
- recovery**, Fatigue as a risk factor for being injured in an occupational accident: results from the Maastricht Cohort Study, **i88**
Measurement quality and validity of the "need for recovery scale", **i3**
Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery, **373**
Need for recovery from work related fatigue and its role in the development and prediction of subjective health complaints, **i62**
- redox chemistry**, Oxidative stress: its role in air pollution and adverse health effects, **612**
- regional pain**, Risk factors in the onset of neck/shoulder pain in a prospective study of workers in industrial and service companies, **649**
- regression analysis**, Evaluation of decompression safety in an occupational diving group using self reported diving exposure and health status, **418**
- remote sensing**, Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
- renal dysfunction**, Land contamination and renal dysfunction, **461**
- reproductive history**, Breast cancer risk in airline cabin attendants: a nested case-control study in Iceland, **807**
- respiratory effects**, Respiratory effects of sulphur dioxide: a hierarchical multicity analysis in the APHEA 2 study, **e2**
Respiratory effects of volcanic emissions, **529**
- respiratory function tests**, Gas stove use and respiratory health among adults with asthma in NHANES III, **759**
- respiratory symptoms**, A longitudinal observation of early pulmonary responses to cotton dust, **115**
Animal production and wheeze in the Agricultural Health Study: interactions with atopy, asthma, and smoking, **e3**
Prevalence and association of welding related systemic and respiratory symptoms in welders, **655**
Prevalence of respiratory symptoms among female flight attendants and teachers, **929**
Prevalence of sensitisation to cellulase and xylanase in bakery workers, **802**
Respiratory symptoms and cotton dust exposure; results of a 15 year follow up observation, **935**
- retinol binding protein**, Associations of lead biomarkers with renal function in Korean lead workers, **551**
- return to work**, Quality of rehabilitation among workers with adjustment disorders according to practice guidelines; a retrospective cohort study, **i21**
- rheumatoid arthritis**, Occupational silica exposure and risk of various diseases: an analysis using death certificates from 27 states of the United States, **122**
- rhinitis**, Estimating the incidence of occupational asthma and rhinitis from laboratory animal allergens in the UK, 1999–2000, **604**
Occupational asthma due to IgE mediated allergy to the flower *Molucella laevis* (Bells of Ireland), **701**
- risk assessment**, Asbestos related diseases from environmental exposure to crocidolite in Da-yao, China. I. Review of exposure and epidemiological data, **35**
Use of personal exposure modelling in risk assessment of air pollutants, **529**
- risk factors**, Ambiguous relation between physical workload and low back pain: a twin control study, **109**
Occupational risk factors for symptomatic lumbar disc herniation; a case-control study, **821**
- rosters**, The impact of roster changes on absenteeism and incident frequency in an Australian coal mine, **43**
- rotator cuff**, Prognosis of shoulder tendonitis in repetitive work: a follow up study in a cohort of Danish industrial and service workers, **1**
- rubber workers**, Exposure related mutagens in urine of rubber workers associated with inhalable particulate and dermal exposure, **97**
- safety**, The impact of roster changes on absenteeism and incident frequency in an Australian coal mine, **43**
- salivary biomarker**, Can salivary lead be used for biological monitoring of lead exposed individuals?, **696**
- SAM axis**, Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
- SARS**, An outbreak of SARS among healthcare workers, **528**
SARS and occupational health in the air, **539**
- self reported health**, Health problems and psychosocial work environment as predictors of long term sickness absence in employees who visited the occupational physician and/or general practitioner in relation to work: a prospective study, **295**
Psychosocial factors at work and self reported health: comparative results of cross sectional and prospective analyses of the French GAZEL cohort, **509**
- semiconductor industry**, Relation between sick leave and selected exposure variables among women semiconductor workers in Malaysia, **262**
- sensitisation**, Prevalence of sensitisation to cellulase and xylanase in bakery workers, **802**
- sensitisers**, Exposure-response relations among laboratory animal workers exposed to rats, **104**
- sentinel health event**, Surveillance in occupational health, **705**
- serum aminotransferase**, Synergistic effect of hepatitis virus infection and occupational exposures to vinyl chloride monomer and ethylene dichloride on serum aminotransferase activity, **774**
- severity**, Age related differences in work injuries and permanent impairment: a comparison of workers' compensation claims among adolescents, young adults, and adults, **e10**
- sewage**, Wastewater exposure and health—a comparative study of two occupational groups, **595**

- sewage worker**, Will sewage workers with endotoxin related symptoms have the benefit of reduced lung cancer?, **144**
- sex ratio**, No change in sex ratio in Ramsar (north of Iran) with high background of radiation, **146**
Sex ratio at birth and exposure to petrochemicals, **704**
- Shanghai**, Relation between colour vision loss and occupational styrene exposure, **222**
- shield**, Availability of thyroid protective lead shields and their use by trainee orthopaedic surgeons, **381**
- shift work**, Shift work and subfecundity: a causal link or an artefact?, **e12**
Work schedules and fatigue: a prospective cohort study, **i47**
The amount of sleep obtained by locomotive engineers: effects of break duration and time of break onset, **e17**
The impact of a week of simulated night work on sleep, circadian phase, and performance, **e13**
The impact of roster changes on absenteeism and incident frequency in an Australian coal mine, **43**
- shoulder exercise**, Effects of a home exercise programme on shoulder pain and functional status in construction workers, **841**
- shoulder impingement**, Effects of a home exercise programme on shoulder pain and functional status in construction workers, **841**
- shoulder pain**, Mechanical and psychosocial factors predict new onset shoulder pain: a prospective cohort study of newly employed workers, **850**
- shoulders**, Risk factors in the onset of neck/shoulder pain in a prospective study of workers in industrial and service companies, **649**
- sick leave**, Do junior doctors take sick leave?, **699**
Downsizing of staff is associated with lowered medically certified sick leave in female employees, **e9**
Relation between sick leave and selected exposure variables among women semiconductor workers in Malaysia, **262**
Work factors as predictors of sickness absence: a three month prospective study of nurses' aides, **271**
- sickness absence**, Fatigue as a predictor of sickness absence: results from the Maastricht cohort study on fatigue at work, **i71**
Health problems and psychosocial work environment as predictors of long term sickness absence in employees who visited the occupational physician and/or general practitioner in relation to work: a prospective study, **295**
Need for recovery from work related fatigue and its role in the development and prediction of subjective health complaints, **i62**
Reducing work related psychological ill health and sickness absence: a systematic literature review, **3**
Sickness absence due to low back pain, **306**
The Depression Anxiety Stress Scales (DASS): detecting anxiety disorder and depression in employees absent from work because of mental health problems, **i77**
- silica**, Occupational silica exposure and risk of various diseases: an analysis using death certificates from 27 states of the United States, **122**
Quantitative relations between exposure to respirable quartz and risk of silicosis, **159**
The long and winding road from silica exposure to silicosis and other health effects, **157**
- silicosis**, Occupational silica exposure and risk of various diseases: an analysis using death certificates from 27 states of the United States, **122**
Quantitative relations between exposure to respirable quartz and risk of silicosis, **159**
The long and winding road from silica exposure to silicosis and other health effects, **157**
- Singapore**, Reported incidence of occupational asthma in France, 1996–99: the ONAP programme, **136**
- skin aberration**, Exposure related mutagens in urine of rubber workers associated with inhalable particulate and dermal exposure, **97**
- sleep debt**, The impact of a week of simulated night work on sleep, circadian phase, and performance, **e13**
- sleepiness**, Driving impairment due to sleepiness is exacerbated by low alcohol intake, **689**
- small airways disease**, Chronic obstructive pulmonary disease due to occupational exposure to silica dust: a review of epidemiological and pathological evidence, **237**
- small areas**, Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
- small for gestational age**, Effect of trihalomethane exposure on fetal development, **173**
- smog**, The biological effects of coarse and fine particulate matter, **313**
- smoking**, Smoking and occupation from the European Community Respiratory Health Survey, **643**
- social class**, Downsizing of staff is associated with lowered medically certified sick leave in female employees, **e9**
- social network**, Psychosocial conditions on and off the job and psychological ill health: depressive symptoms, impaired psychological wellbeing, heavy consumption of alcohol, **489**
- social support**, Physical and psychosocial risk factors for lateral epicondylitis: a population based case-referent study, **322**
Psychosocial conditions on and off the job and psychological ill health: depressive symptoms, impaired psychological wellbeing, heavy consumption of alcohol, **489**
- soft tissue sarcoma**, Cancer risk for chemical workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin, **672**
Risk of soft tissue sarcomas and residence in the neighbourhood of an incinerator of industrial wastes, **680**
- spatial data analysis**, Geographical pattern of brain cancer incidence in the Navarre and Basque Country regions of Spain, **504**
- steatosis**, Periportal fibrosis and other liver ultrasonography findings in vinyl chloride workers, **60**
- strategy**, Occupational asthma, **711**
- stress**, Increased plasma homocysteine levels in shift working bus drivers, **662**
Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
- stress related disorders**, The Depression Anxiety Stress Scales (DASS): detecting anxiety disorder and depression in employees absent from work because of mental health problems, **i77**
- stress test**, Physiological differences between burnout patients and healthy controls: blood pressure, heart rate, and cortisol responses, **i54**
- styrene**, Exposure assessment of monoterpenes and styrene: a comparison of air sampling and biomonitoring, **599**
- subgroup**, Ambiguous relation between physical workload and low back pain: a twin control study, **109**
- substantia nigra**, Endotoxin: is it an environmental factor in the cause of Parkinson's disease?, **378**
- sulphur dioxide**, Respiratory effects of sulphur dioxide: a hierarchical multicity analysis in the APHEA 2 study, **e2**
- sulphuric acid**, Clinical and histopathological changes of the nasal mucosa induced by occupational exposure to sulphuric acid mists, **395**
The effect of sulphurous air pollutant exposures on symptoms, lung function, exhaled nitric oxide, and nasal epithelial lining fluid antioxidant concentrations in normal and asthmatic adults, **e15**
- sunlight**, Risk factors for cutaneous malignant melanoma among aircrews and a random sample of the population, **815**
- supervisor**, The effect on ambulatory blood pressure of working under favourably and unfavourably perceived supervisors, **468**
- surveillance**, Reported incidence of occupational asthma in France, 1996–99: the ONAP programme, **136**
The continuing challenge to reduce the burden of occupational asthma, **713**
- surveillance schemes**, Occupational contact dermatitis to nickel: experience of the British dermatologists (EPIDERM) and occupational physicians (OPRA) surveillance schemes, **954**
- susceptibility**, Risk factors among elderly for short term deaths related to high levels of air pollution, **684**
- sweat rate**, Fluid losses and hydration status of industrial workers under thermal stress working extended shifts, **90**
- swimming**, Lung hyperpermeability and asthma prevalence in schoolchildren: unexpected associations with the attendance at indoor chlorinated swimming pools, **385**
- systematic review**, Intervention strategies to reduce musculoskeletal injuries associated with handling patients: a systematic review, **e6**
The effectiveness of current approaches to workplace stress management in the nursing profession: an evidence based literature review, **10**
- TCDD**, Cancer risk for chemical workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin, **672**
- teachers**, Prevalence of respiratory symptoms among female flight attendants and teachers, **929**

- tennis elbow**, Physical and psychosocial risk factors for lateral epicondylitis: a population based case-referent study, **322**
- Th2 response**, Repeated daily exposure to 2 ppm nitrogen dioxide upregulates the expression of IL-5, IL-10, IL-13, and ICAM-1 in the bronchial epithelium of healthy human airways, **892**
- thyroid**, Availability of thyroid protective lead shields and their use by trainee orthopaedic surgeons, **381**
- thyroid stimulating hormone**, Organochlorine compounds and concentrations of thyroid stimulating hormone in newborns, **301**
- time of day**, The amount of sleep obtained by locomotive engineers: effects of break duration and time of break onset, **e17**
- time to pregnancy**, Shift work and subfecundity: a causal link or an artefact?, **e12**
- Time To Pregnancy and occupational lead exposure, **752**
- timed allen test**, Hand-arm vibration syndrome in Swedish car mechanics, **287**
- TNT**, Exposure to nitroaromatic explosives and health effects during disposal of military waste, **483**
- tobacco**, One year effectiveness of an individualised smoking cessation intervention at the workplace: a randomised controlled trial, **358**
- Smoking and occupation from the European Community Respiratory Health Survey, **643**
- total mortality**, Investigating the dose-response relation between air pollution and total mortality in the APHEA-2 multicity project, **977**
- total sleep time**, The amount of sleep obtained by locomotive engineers: effects of break duration and time of break onset, **e17**
- traffic noise**, Health status as a potential effect modifier of the relation between noise annoyance and incidence of ischaemic heart disease, **739**
- trihalomethane**, Effect of trihalomethane exposure on fetal development, **173**
- trinitrotoluene**, Exposure to nitroaromatic explosives and health effects during disposal of military waste, **483**
- twin study**, Ambiguous relation between physical workload and low back pain: a twin control study, **109**
- UK**, Occupational asthma, **711**
- UK-POEM**, Urine mutagenicity and lymphocyte DNA damage in fruit growers occupationally exposed to the fungicide captan, **910**
- unemployment**, High local unemployment and increased mortality in Danish adults; results from a prospective multilevel study, **e16**
- upper limb**, Pain tolerance in upper limb disorders: findings from a community survey, **217**
- urinary mutagenicity**, Exposure related mutagens in urine of rubber workers associated with inhalable particulate and dermal exposure, **97**
- urine**, Chromium (VI) induced oxidative damage to DNA: increase of urinary 8-hydroxydeoxyguanosine concentrations (8-OHdG) among electroplating workers, **590**
- urine mutagenicity**, Urine mutagenicity and lymphocyte DNA damage in fruit growers occupationally exposed to the fungicide captan, **910**
- UV radiation**, Cancer incidence in airline cabin crew, **805**
- Cancer incidence in airline cabin crew: experience from Sweden, **810**
- validity**, Measurement quality and validity of the "need for recovery scale", **i3**
- variance component**, Exposure assessment of monoterpenes and styrene: a comparison of air sampling and biomonitoring, **599**
- vasospasm**, Skin temperature recovery from cold provocation in workers exposed to vibration: a longitudinal study, **962**
- VDU**, Does computer use pose an occupational hazard for forearm pain; from the NUDATA study, **e14**
- Work related and individual predictors for incident neck pain among office employees working with video display units, **475**
- vibration**, The relative importance of whole body vibration and occupational lifting as risk factors for low-back pain, **715**
- vibration frequency**, Dose-response patterns for vibration-induced white finger, **16**
- vibration-induced white finger**, Dose-response patterns for vibration-induced white finger, **16**
- video observations**, Risk factors in the onset of neck/shoulder pain in a prospective study of workers in industrial and service companies, **649**
- vinyl chloride exposure**, Periportal fibrosis and other liver ultrasonography findings in vinyl chloride workers, **60**
- virus**, Mortality in workers in poultry slaughtering/processing plants: the Missouri poultry cohort study, **784**
- vocational rehabilitation**, Return to work of cancer survivors: a prospective cohort study into the quality of rehabilitation by occupational physicians, **352**
- volcanic eruption**, Respiratory effects of volcanic emissions, **529**
- Volcanic ash and respiratory symptoms in children on the island of Montserrat, British West Indies, **207**
- wafer polishing**, Relation between sick leave and selected exposure variables among women semiconductor workers in Malaysia, **262**
- waste incinerator**, Municipal waste incinerators: air and biological monitoring of workers for exposure to particles, metals, and organic compounds, **563**
- water**, Reported incidence of occupational asthma in France, 1996–99: the ONAP programme, **136**
- weight carrying/lifting**, Occupational risk factors for symptomatic lumbar disc herniation; a case-control study, **821**
- welding**, Prevalence and association of welding related systemic and respiratory symptoms in welders, **655**
- West Bank**, Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery, **373**
- William Harvey**, William Harvey and air pollution, **147**
- work**, Coverage of work related fatalities in Australia by compensation and occupational health and safety agencies, **195**
- Fatigue, burnout, and chronic fatigue syndrome among employees on sick leave: do attributions make the difference?, **i26**
- Measurement quality and validity of the "need for recovery scale", **i3**
- The right treatment to the right patient at the right time, **235**
- work conditions**, Psychosocial conditions on and off the job and psychological ill health: depressive symptoms, impaired psychological wellbeing, heavy consumption of alcohol, **489**
- work disability**, Cardiorespiratory fitness and risk of disability pension: a prospective population based study in Finnish men, **765**
- work related respiratory symptoms**, Physician diagnosed asthma, respiratory symptoms, and associations with workplace tasks among radiographers in Ontario, Canada, **254**
- work stress**, Medical staff in emergency situations: severity of patient status predicts stress hormone reactivity and recovery, **373**
- Reducing work related psychological ill health and sickness absence: a systematic literature review, **3**
- The effect on ambulatory blood pressure of working under favourably and unfavourably perceived supervisors, **468**
- work stressors**, Fatigue in employees with diabetes: its relation with work characteristics and diabetes related burden, **i93**
- worker**, Mental ill health in workers: observations from a few Indian populations, **225**
- Neurobehavioural effects among workers occupationally exposed to organophosphorous pesticides, **279**
- workers' compensation**, Early prognostic factors for duration on temporary total benefits in the first year among workers with compensated occupational soft tissue injuries, **244**
- working population**, An epidemiological approach to study fatigue in the working population: the Maastricht Cohort Study, **i32**
- Associations between fatigue attributions and fatigue, health, and psychosocial work characteristics: a study among employees visiting a physician with fatigue, **i99**
- working sample**, Assessment of fatigue among working people: a comparison of six questionnaires, **i10**
- workload**, Ambiguous relation between physical workload and low back pain: a twin control study, **109**
- Increased occupational physical activity does not improve physical fitness, **983**
- workplace**, Competence at the workplace, **225**
- workplace bullying**, Workplace bullying and the risk of cardiovascular disease and depression, **779**
- workplace counselling**, Workplace counselling, **899**
- workplace stress**, Workplace counselling, **899**
- worksite**, One year effectiveness of an individualised smoking cessation intervention at the workplace: a randomised controlled trial, **358**
- xray**, Availability of thyroid protective lead shields and their use by trainee orthopaedic surgeons, **381**
- xylanase**, Prevalence of sensitisation to cellulase and xylanase in bakery workers, **802**
- young workers**, Age related differences in work injuries and permanent impairment: a comparison of workers' compensation claims among adolescents, young adults, and adults, **e10**