

P46

ASSESSING EXPOSURES TO CLEANING AND DISINFECTING CHEMICALS FOR AN EPIDEMIOLOGIC STUDY OF ASTHMA IN HEALTHCARE OCCUPATIONS

Mohammed Abbas Virji, Ryan LeBouf, Rena Saito, Aleksandr Stefaniak, Marcia Stanton, Paul Henneberger *NIOSH Division of Respiratory Disease Studies, Morgantown, USA*

10.1136/oemed-2011-100382.260

Objectives Cleaning and disinfectant products have been identified as important risk factors for asthma, and are used extensively in healthcare; however, quantitative exposure measurements

of these etiologic agents are not well characterised. The objective of this study was to quantitatively characterise exposure to cleaning and disinfecting compounds, or their surrogates, and to develop a job/task-exposure matrix (J/TEM) for an epidemiologic study of asthma in healthcare occupations.

Methods Exposure assessments were conducted at four hospitals targeting six healthcare occupations: nursing/nursing aides, respiratory therapists, dental assistants/technicians, medical and clinical laboratory technicians, housekeepers, and disinfecting and sterilising technicians. Area and personal air monitoring was conducted for time-integrated volatile organic compounds (VOCs) and continuous measurements of total VOCs. Tasks performed, products used, manner of application, tools, controls and personal protective equipment use were logged in a task diary at 5-min intervals. A range of VOCs were quantified including chlorinated hydrocarbons, ketones, terpenes, acrylics, alcohols, aromatic and aliphatic hydrocarbons, most of which were in the 1 to 10 parts-per-billion range except for alcohols and m,p-xylene (parts-per-million concentrations).

Results The chemical agents and levels varied by occupation and form the basis for developing an asthma-specific J/TEM for healthcare occupations. The jobs/tasks form the job axis while the chemicals/ classes of chemicals form the exposure axis with measured concentrations as cells of the J/TEM.

Conclusions These results will be incorporated into a survey questionnaire to elicit worker-specific information on jobs, tasks, and products used, and together, will be used to develop exposure estimates for the epidemiologic study participants.