

from pleura cancer and melanoma. This cohort study is the most informative ever conducted in France among nuclear workers.

## 0278 THE PESTIMAT PROGRAM: DEVELOPMENT OF A CROP EXPOSURE MATRIX FOR PESTICIDE EXPOSURE ASSESSMENT IN AGRICULTURE

<sup>1,2</sup>Isabelle Baldi, <sup>1,2</sup>Camille Carles, <sup>1</sup>Anne Cantagrel, <sup>3,4</sup>Yannick Lecluse, <sup>3,4</sup>Elodie Niez, <sup>5</sup>Pascal Fabbro-Peray, <sup>3,4</sup>Pierre Lebaillay. <sup>1</sup>Univ. Bordeaux, ISPED, Laboratoire Santé Travail Environnement, Centre INSERM U 897 Epidemiologie-Biostatistique, F-33000 Bordeaux, France; <sup>2</sup>CHU de Bordeaux, Service de Médecine Du Travail, F-33000 Bordeaux, France; <sup>3</sup>INSERM, UMR 1086- Cancers Et Préventions, F-14000 Caen, France; <sup>4</sup>Centre François Baclesse, F-14000 Caen, France; <sup>5</sup>DIM, Hôpital G Doumergue, F-34000 Montpellier, France

10.1136/oemed-2014-102362.108

**Objectives** Pesticides include ~1000 chemicals with various toxicological properties. Pesticide exposure assessment represents a crucial methodological issue for epidemiological studies. A reconstitution of pesticides used in entire professional careers based only on the memory is questionable. PESTIMAT is a crop exposure matrix, which intends to reconstitute parameters of pesticide exposure in France.

**Method** PESTIMAT is made of tables crossing crops and active ingredients (a.i.) by year from 1950 to 2010 and providing the following exposure parameters: 1) probability corresponding to the proportion of farmers having used the a.i. (in%), 2) frequency expressed as the number of treatment days, 3) intensity documented by the application rate of the a.i. (in kg/ha). Parameters were obtained by the combination of six complementary sources: i) registrations and withdrawals from the Agriculture Ministry, ii) information from ACTA\* on products marketed iii) agricultural recommendations by the Plant Health Protection body, iv) treatment calendars provided by farmers, v) information from professional associations of farmers, vi) data from the industry

**Results** To date, 529 a.i. usable between 1950 and 2010 on 4 crops are included in PESTIMAT, corresponding to 160 fungicides (30%), 160 herbicides (30%) and 209 insecticides (40%). The total numbers were comparable on wheat/barley (N = 293), vineyards (N = 280), and apple-growing (N = 267) but lower on corn (N = 196). We will present an illustration with data obtained for dithiocarbamates fungicides, and examples of use in epidemiological studies, e.g. PHYTONER and AGRICAN cohorts.

**Conclusions** The multisource information available in the PESTIMAT matrix should enable to assess associations between pesticides and health at the a.i. level.

## 0282 A CROSS-SECTIONAL STUDY OF MARKERS OF EARLY IMMUNOLOGICAL AND CARDIOVASCULAR HEALTH EFFECTS AMONG A POPULATION EXPOSED TO CARBON NANOTUBES: THE CANTES STUDY

<sup>1</sup>Roel Vermeulen, <sup>2</sup>Anjoeka Pronk, <sup>1</sup>Jelle Vlaanderen, <sup>3</sup>Dean Hosgood, <sup>4</sup>Nat Rothman, <sup>4</sup>Allan Hildesheim, <sup>4</sup>Debra Silverman, <sup>1</sup>Annemarie Melis, <sup>2</sup>Suzanne Spaan, <sup>2</sup>Eef Voogd, <sup>5</sup>Peter Hoet, <sup>5</sup>Lode Godderis, <sup>4</sup>Qing Lan. <sup>1</sup>IRAS Utrecht University, Utrecht, The Netherlands; <sup>2</sup>TNO, Zeist, The Netherlands; <sup>3</sup>Albert Einstein College of Medicine, New York, NY, USA; <sup>4</sup>National Cancer Institute, Bethesda, MD, USA; <sup>5</sup>University of Leuven, Leuven, Belgium

10.1136/oemed-2014-102362.109

**Objectives** To assess the association between occupational exposure to carbon nanotubes (CNTs) and early immunological and cardiovascular health effects.

**Method** We collected biological samples (blood, urine, buccal and nasal epithelium cells) from a group of workers in a CNT production factory and approximately age and sex matched unexposed individuals working in a different workplace. Based on an initial assessment of exposure data collected in the weeks before biological sampling we further classified CNT production workers as operators (high probability of exposure) and R&D workers (lower probability of exposure). We analysed blood samples for a panel of inflammatory markers.

**Results** We included 8 operators (median age 34.8), 16 R&D workers (median age 32.2), and 43 controls (median age 30.6). Preliminary analyses provide some evidence for an association between CNT exposure and selected cytokines. We observed a similar pattern when we restricted our analyses to non-smoking men. Concentrations of cytokines were exposure dependently upregulated with higher levels among operators than R&D workers; both higher than unexposed workers (p for trend <0.05).

**Conclusions** We observed some indications of early immunological health effects in a pilot study conducted among workers exposed to CNTs. Further analyses are planned, including assessment of lung function, heart rate variability, oxidative stress, and blood coagulation markers. Extensive exposure measurements were conducted in the CNT production factory as well and additional analyses will use these individual exposure measurements to more thoroughly explore exposure dependent effects.

## 0285 DEVELOPMENT OF A SOURCE-BASED APPROACH TO ASSESSING OCCUPATIONAL EXPOSURE TO ELECTROMAGNETIC FIELDS IN THE INTEROCC STUDY

<sup>1</sup>Javier Vila, <sup>2</sup>Joseph D Bowman, <sup>3</sup>Laurel Kincl, <sup>2</sup>Dave L Conover, <sup>4</sup>Martie van Tongeren, <sup>1</sup>Jordi Figuerola, <sup>3</sup>Lesley Richardson, <sup>1</sup>Elisabeth Cardis, <sup>6</sup>on behalf of the INTEROCC Study Group. <sup>1</sup>Centre for Research in Environmental Health (CREAL), Barcelona, Spain; <sup>2</sup>The National Institute for Occupational Safety and Health (NIOSH), Cincinnati, OH, USA; <sup>3</sup>Oregon State University (OSU), Corvallis, OR, USA; <sup>4</sup>Institute of Occupational Medicine (IOM), Edinburgh, UK; <sup>5</sup>University of Montreal Hospital Research Centre (CRCHUM), Montreal, Canada; <sup>6</sup>Institute of Occupational Medicine (IOM), Edinburgh, Spain

10.1136/oemed-2014-102362.110

**Objectives** Exposure to electromagnetic fields (EMF) has become ubiquitous in modern life and concern has increased regarding possible associated health effects. To date, assessment of occupational exposure has relied on job-exposure matrices, with exposure estimates for very broad occupational categories. To move EMF research forward, a new approach was necessary. A source-based strategy, incorporating detailed information on tasks, equipment used and work organisation could allow a more individualised exposure assessment.

**Method** Information on occupational histories and sources of EMF was collected as part of the INTERPHONE-INTEROCC study, providing an opportunity to assess occupational EMF exposure by assigning exposure to each source used. A source-exposure matrix (SEM) was developed based on measurements identified in the literature and estimates obtained through experts' elicitation, for sources without available measurements. This paper focuses on the SEM development methodology to ensure the quality and representativeness of the estimates.