CORRESPONDENCE

Sex ratio of offspring of men exposed to sodium borates

Editor—Whorton *et al* report 529 births (250 sons and 279 daughters) sired by men exposed to sodium borates.¹ With official data on the comparable non-exposed United States population (adjusted for maternal age, race, parity, and calendar time) these authors estimate an expected 466.6 births. They took the unusual step of contrasting the observed sex ratio (proportion male) with the numbers of the two sexes expected assuming simultaneously both: a United States live birth sex ratio of 0.512; and the comparable United States fertility rates.

They thus calculate expected values of sons and daughters that are less than both the observed values. So they are able to conclude that the unusual observed sex ratio is not due to a deficit of boys. This, they considered, would allay suspicions that borates represent a reproductive hazard to male fertility.

I should like to present an alternative argument based on their data. It would be more conventional to contrast the observed sex ratio (250 boys, 279 girls) directly with the expected sex ratio of 0.512. This gives an associated χ^2 value of 3.3. Bearing in mind the suspicion of reproductive hazard cited by Whorton *et al*,¹ and the evidence that occupational reproductive hazards to men may manifest themselves in low sex ratios of offspring,² it would seem appropriate to use a one-tailed test. (In this case $P < 0.05$.) So I would conclude that indeed there are grounds for suspicion that borates affect the male reproductive system. More data should be gathered on the sex ratios of offspring of men exposed to these materials. If these prove to be low too, then more invasive investigation (sperm counts, hormone assays) would be justified.

Lastly, the excess fertility in the exposed men was estimated by adjusting United States national fertility rates for maternal age, parity, race, and time. These men were employed in the Mojave Desert. Would not fertility be greater in such areas than in urban areas? If so, the excess fertility may be illusory.

WH JAMES

The Galton Laboratory,
Department of Genetics and Biometry,
University College London,
Wolfson House,
4 Stephenson Way,
London NW1 2HE.

1 Whorton MD, Haas JL, Trent L, Wong O. Reproductive effects of sodium borates on male employees: birth rate assessment. *Occup Environ Med* 1994;51:761-7.

2 James WH. Occupations associated with low offspring sex ratios. *Am J Ind Med* 1994; 25:607-8.

Author's reply—James' comments on our sex ratio analyses need closer scrutiny on several counts. Firstly, contrary to James' comments, in our sex ratio analysis, we did not assume comparable fertility rates between our study and the United States general

population. On the other hand, in his comparison between the observed overall sex ratio in our study and the ratio for the United States, James implicitly made the assumption of comparable fertility rates between our study and that for the United States.

The primary purpose of our study was to determine if there was a deficit or excess of births by sex rather than the sex ratio itself. As such, a direct comparison of the two sex ratios would have been appropriate only if the underlying fertility rates in our study and in the general population were comparable. As shown through the use of the standardised birth ratio, the fertility rates in our study were significantly higher than those in the general population. From the methodological point of view, the problem of directly comparing sex ratios is similar to that of applying the proportional mortality ratio (PMR) in mortality analysis. It is well known that PMR compares proportions of deaths from specific causes between two groups, and it is a valid measure for mortality risk only if the underlying death rates are comparable in the two groups.^{1,2} Similarly, a direct comparison of sex ratios determines whether there is a deficit or excess by sex only if the underlying fertility rates are comparable in the two populations being compared.

James' rationale for a one tailed test assumes an original hypothesis of a lower sex ratio (proportion male) associated with exposure to sodium borates. Based on previous data on sodium borates, we had no preconceived idea whether the sex ratio (proportion male) should be higher or lower. Thus, James' argument for a one tailed test was based on a retrospective examination of the data, which would render the conventional P value of 0.05 inappropriate, and the χ^2 of 3.3 calculated by James would not have been significant.

James implied that male reproductive hazards would most likely result in an altered sex ratio. Among the papers cited in another letter to the editor by James,³ one report by Potashnik and Yanai-Inbar⁴ was due to a chemical exposure (dibromochloropropane). Potashnik and Yanai-Inbar reported an increase in girls born to men exposed to dibromochloropropane who had recovered from either azoospermia or severe oligospermia. The mechanism for this is unknown although James believed this effect to be due to an increase in gonadotrophin concentrations.³ Men who are azoospermic from exposures to dibromochloropropane have raised gonadotrophin (FSH and LH) concentrations due to direct testicular damage.⁵ There is no evidence at the present to think that there is a causal relation between the increase in gonadotrophins as a result of direct testicular toxicity and the increase in female offspring.

James' entire proposition relied on only one statistic: the sex ratio for the entire group. He ignored detailed exposure-response analyses. As reported in our paper, the highest female sex ratios (proportion female) were observed in the lowest exposure categories (table 7), and there was no

statistical trend of sex ratio by exposure. Furthermore, the female sex ratio for participants during the period of high exposure (0.525) was practically identical to that for the rest of the participants (0.528).

James also questioned the appropriateness of comparing fertility rates of workers in our study who lived in the Mojave Desert to those in the general population. There is no evidence that people who live in the Mojave Desert communities are more fertile than the rest of the nation. Although the location of the factory is rural, most of the employees live in small, urban communities and commute to work. Furthermore, James again does not seem to consider the fact that we also compared fertility ratios internally by exposure category (tables 4 and 5).

M DONALD WHORTON
OTTO WONG
1135 Atlantic Avenue, Alameda,
California, 94501, USA

- 1 Wong O, Decoufle P. Methodological issues involving the standardized mortality and proportional mortality ratio in occupational studies. *J Occup Med* 1982;24:299-304.
- 2 Wong O, Morgan RW, Kheifets L, Larson SR. Comparison of SMR, PMR, and PCMR in a cohort of union members potentially exposed to diesel exhaust emissions. *Br J Ind Med* 1985;42:435-48.
- 3 James WH. Occupations associated with low offspring sex ratios. *Am J Ind Med* 1994;25: 607-8.
- 4 Potashnik G, Yanai-Inbar I. Dibromochloropropane (DBCP): an 8-year re-evaluation of testicular function and reproductive performance. *Fertil Steril* 1987;47:317-23.
- 5 Whorton D, Milby TH, Krauss RM, Stubbs HA. Testicular function in DBCP exposed pesticide workers. *J Occup Med* 1979;21: 161-6.

Assessment of risk of lung cancer among mild steel and stainless steel welders

Editor—I have read with great interest the paper by Sjogren *et al*.¹ The authors carried out a meta-analysis of five epidemiological studies—three case-control²⁻⁴ and two historical cohort⁵ studies—on the occurrence of lung cancer among stainless steel (SS) welders. The results clearly indicated a relation between SS welding and lung cancer, the pooled relative risk being 1.94 (95% confidence interval (95% CI) 1.28-2.92). The authors concluded that "it is time to reconsider the IARC (International Agency for Research on Cancer) statement from 1990⁷ and to separate SS welding fumes from other welding fumes".

Here I compare the risks of lung cancer of mild steel (MS) welders *v* SS welders with additional results from the studies included in the meta-analysis of Sjogren *et al*¹ and from other studies not included in this meta-analysis.⁸⁻¹²

The observed odds ratio (OR) among MS welders in the Danish case-control study was 1.65 (95% CI 1.03-2.65),³ which is similar to that of SS welders, 1.54 (95% CI 0.83-2.84). In the French cohort mortality study the standardised mortality ratio (SMR) for lung cancer was slightly higher for MS welders than for welders predominantly exposed to chromium VI—that is, 1.59 (95% CI 0.73-3.02) *v* 1.03 (95% CI 0.12-3.71).⁵

Some other studies did not detect high risks of lung cancer among SS welders. Becker *et al* followed up a cohort of SS welders compared with an internal reference group of turners. The welders had a

relative risk for lung cancer of 1.7 (95% CI 0.7–4.0).⁸ This increased value, however, was not confirmed by a second follow up as the relative risk decreased to 1.2 (95% CI 0.6–2.2).⁸ These results remained unchanged after adjustments for smoking.⁹ Moreover, with an external reference, the lung cancer SMR of SS welders was low (1.13 (95% CI 0.67–1.91)).⁹

The case-control study by Hull *et al* in Los Angeles county¹⁰ showed ORs of 0.9 (95% CI 0.5–1.8) for SS welders and 1.3 (95% CI 0.6–2.3) for SS welders predominantly exposed to hexavalent chromium and nickel (manual metal arc welders), whereas the OR was 1.6 (95% CI 0.8–3.1) for MS welders. Adjustments for smoking made little difference.¹⁰

The European mortality study included 11 092 welders from eight countries.¹¹ This large study provided no consistent difference between MS welders, for which the SMR was 1.78 (95% CI 1.27–2.43, statistically different) and predominantly SS welders whose SMR was 1.23 (95% CI 0.75–1.90). The results of this study for cancer incidence provided the same pattern as those for mortality as the standardised incidence ratios (SIRs) of lung cancer were 1.75 (95% CI 1.22–2.42) for MS welders and 1.39 (95% CI 0.74–2.38) for predominantly SS welders.¹²

In conclusion, it seems to me that, although SS welders are potentially exposed to chromium and nickel compounds,⁷ epidemiological studies focused on the risk of lung cancer of SS welders do not provide clear evidence to suggest that SS welders are at higher risk of lung cancer than MS welders. Therefore, the cause of the excess lung cancers found among MS and SS welders is an unanswered issue.⁵ A recent study by Jockel *et al* supports the hypothesis that some of the excess risk of lung cancer among welders could be due to exposure to asbestos.¹³ Further investigations that controlled for smoking, exposure to asbestos, and possible differential healthy worker effects among MS *v* SS welders,¹¹ are needed.

JEAN JACQUES MOULIN

Institut National de Recherche et de Sécurité (INRS),
Service d'Epidémiologie, BP 27, F 54500
Vandœuvre

- 1 Sjögren B, Hansen KS, Kjuus H, Persson PG. Exposure to stainless steel welding fumes and lung cancer: a meta-analysis. *Occup Environ Med* 1994;57:335–6.
- 2 Gerin M, Siemiatycki J, Richardson L, Pellerin J, Lakhani R, Dewart R. Nickel and cancer associations from a multicenter occupation exposure case-referent study: preliminary findings. *LARC Sci Publ* 1984; 53:105–15.
- 3 Hansen KS, Lauritsen J. Lung cancer mortality in stainless steel and mild steel welders—A case-referent study. *Am J Ind Med* 1995; (in press).
- 4 Kjuus H, Skjaerven R, Langard S, Lien JT, Aamodt T. A case-referent study of lung cancer, occupational exposures, and smoking. I. Comparison of title-based and exposure-based occupational information. *Scand J Work Environ Health* 1986;12:193–202.
- 5 Moulin JJ, Wild P, Haguenoer JM, Faucon D, De Gaudemaris R, Mur JM, *et al*. A mortality study among mild steel and stainless steel welders. *Br J Ind Med* 1993;50: 234–43.
- 6 Sjögren B, Gustavsson E, Hedström L. Mortality in two cohorts of welders exposed to high- and low-levels of hexavalent chromium. *Scand J Work Environ Health* 1987;13:247–51.
- 7 International Agency for Research on Cancer. Chromium, nickel and welding. *IARC Monogr Eval Carcinog Risks Hum*. Lyon: IARC, 1990.

- 8 Becker N, Claude J, Frentzel-Beyme R. Cancer risk of arc welders exposed to fumes containing chromium and nickel. *Scand J Work Environ Health* 1985;11:75–82.
- 9 Becker N, Chang-Claude J, Frentzel-Beyme R. Risk of cancer for arc welders in the Federal Republic of Germany: results of a second follow up (1983–8). *Br J Ind Med* 1991;48:675–83.
- 10 Hull CJ, Doyle E, Peters JM, Garabrant DH, Bernstein L, Preston-Martin S. Case-control study of lung cancer in Los Angeles county welders. *Am J Ind Med* 1989;16: 103–12.
- 11 Simonato L, Fletcher AC, Andersen E, Anderson K, Becker N, Chang-Claude J, *et al*. A historical prospective study of European stainless steel, mild steel, and shipyard welders. *Br J Ind Med* 1991; 48:145–54.
- 12 International Agency for Research on Cancer (IARC). Mortality and cancer incidence follow-up of an historical cohort of European welders. Lyon: IARC 1990. (IARC Internal Report 89/003.)
- 13 Jockel KH, Ahrens W, Bolm-Audorf U. Lung cancer risk and welding—Preliminary results of an ongoing case-control study. *Am J Ind Med* 1994;25:805–12.

Urinary N-acetyl-β-D-glucosaminidase (NAG) and exposure to inorganic lead

Editor—Urinary activity of N-acetyl-β-D-glucosaminidase (NAG) had been reported to be one of the earliest markers to be increased in workers exposed to lead.^{1–4} The underlying mechanism for an increase in NAG activity is yet to be elucidated. It was suggested that the increase in NAG activity may be due to stimulation of exocytosis.⁵ In a recent publication⁶ based on workers from a lead smelting plant, it was further suggested that increase in NAG activity among workers exposed to lead was due to concomitant "albeit slight cadmium exposure". Their conclusion was based on the finding that urinary cadmium concentration (CdU) was the only significant variable in explaining the variation in NAG activity through stepwise regression analysis. Furthermore, the correlation coefficient (*r*) between NAG and CdU, was 0.41 in the non-exposed and 0.35 among the exposed workers.

Our data from exposed workers in a polyvinyl chloride lead stabiliser plant did not support this hypothesis. The increases in NAG, the heat labile isoenzyme (NAG-A), and the heat stable isoenzyme (NAG-B) were all highly associated with the recent change in blood lead (PbB) concentration over the past six months (PbBΔ).⁴ Among several exposure indices derived from serial PbB concentrations, PbBΔ was the only significant variable to account for the variation in NAG, NAG-B, and NAG-A. Furthermore, CdU and blood cadmium (CdB) did not correlate well with NAG and its isoenzymes. When they were forced into

the regression model, CdU was in fact negatively correlated (table). Activity of NAG is also increased when other xenobiotics affect the proximal tubules.^{7–9} It is indeed very unlikely that NAG is a specific marker for cadmium. As inorganic lead is known to cause proximal tubular dysfunction, it is therefore not surprising to see an association between lead exposure and NAG.

Among lead smelters, it is possible that there may be significant concomitant exposure to cadmium. This is evident by the fairly wide range of CdB results (0.3 to 5.9 µg/l) and the significantly higher CdB and CdU concentrations among the workers exposed to lead.⁶ Our cohort of PVC lead stabiliser workers had never been exposed to cadmium. Their geometric mean (range) CdB and CdU were much lower than those of the lead smelters (CdB: 0.56 (0.10 to 1.92) µg/l; CdU 0.41 (0.12 to 2.07) µg/g creatinine). We have also previously reported that NAG activity was better correlated with CdB and about 40% of the workers exposed to cadmium with a CdB of less than 3.0 µg/l had increased NAG concentrations.¹⁰

Hence it is possible that where there is significant cadmium exposure, cadmium plays a greater part in increasing NAG activity than does lead, and perhaps even masks the contribution of lead. Our data show that lead does increase NAG activity in the absence of cadmium exposure.

K S CHIA

Department of Community,
Occupational and Family Medicine,
National University of Singapore and WHO
Collaborating Centre for Occupational Health

- 1 Meyer RB, Fischbein A, Rosenman K, Lerman Y, Drayer DE, Reidenberg MM. Increased urinary enzyme excretion in workers exposed to nephrotoxic chemicals. *Am J Ind Med* 1984;76:989–98.
- 2 Ong CN, Endo G, Chia KS, Phoon WO, Ong HY. Evaluation of renal function in workers with low blood lead levels. In: Foa V, Emmett EA, Maroni M, Colombi A, eds. *Occupational and environmental chemical hazards: cellular and biochemical indices for monitoring toxicity*. W Sussex: Harwood, 1987:327–333.
- 3 Endo G, Horiguchi S, Kiyota A. Urinary N-acetyl-β-D-glucosaminidase activity in lead exposed workers. *J Appl Toxicol* 1990;10: 235–8.
- 4 Chia KS, Mutti A, Tan C, Ong HY, Jeyaratnam J, Ong CN, Lee E. Urinary N-acetyl-β-D-glucosaminidase activity in workers exposed to inorganic lead. *Occup Environ Med* 1994;51:125–9.
- 5 Bernard A, Lauwerys R. Epidemiological applications of early marker of nephrotoxicity. *Toxicol Lett* 1989;46:293–306.
- 6 Roels H, Lauwerys R, Konings J, Buchet JP, Bernard A, Green S, *et al*. Renal function and hyperfiltration capacity in lead smelter workers with high bone lead. *Occup Environ Med* 1994;51:505–12.
- 7 Price RG. Urinary enzymes, nephrotoxicity and renal disease. *Toxicology* 1982;23: 99–134.

Contribution of PbBΔ, CdB, and CdU to variation in NAG*

Independent variables	Partial regression coefficients (β) (95% CI)	Unit free β	r ²
Total NAG:			
PbBΔ	4.12 (2.22 to 6.01)	0.63	0.45
CdB	0.16 (–0.08 to 0.40)	0.19	
CdU	–0.14 (–0.31 to 0.04)	–0.22	
NAG-B:			
PbBΔ	11.9 (5.96 to 16.43)	0.66	0.43
CdB	–0.45 (–0.95 to 0.05)	–0.28	
CdU	–0.03 (–0.39 to 0.34)	–0.02	

*Multiple linear regression analysis with log transformed data.