Blood lead in pregnant women in the urban slums of Lucknow, India

Shally Awasthi, Rajiv Awasthi, Vinod Kumar Pande, R C Srivastav, Howard Frumkin

Abstract

Objective—To determine the concentrations of blood lead (PbB) in pregnant women in the slums of Lucknow, north India.

Methods—Of the 203 designated municipal slums in Lucknow, 70 were randomly selected for study and a cohort of 500 pregnant women was enrolled. Each participant was interviewed with questions on possible sources of exposure to lead, surrogates of nutritional status were measured, and PbB was measured.

Results—The mean PbB was 14.3 µg/dl and 19.2% of women had PbB ≥ 20 µg/dl. PbB was not associated with age, height, weight, gestation, or history of abortions, although higher PbB was associated with higher parity. Women living in inner city neighbourhoods near heavy vehicular traffic had PbB 2.2 µg/dl higher (95% confidence interval (95% CI) 0.8 to 3.6) than those living in other neighbourhoods. The PbB was not associated with reported use of piped water or the presence of paint in homes, and increasing PbB was unexpectedly associated with decreasing use of the eye cosmetic "surma" and the duration of gestation.

Conclusions—The high PbB found in this population raises concern about fetal development and points to the urgent need to reduce exposure to lead.

(*Occup Environ Med* 1996;53:836-840)

Keywords: lead; pregnancy; fetus; neurotoxicity; India

Although lead toxicity in children and adults is well recognised, exposure to lead is of special concern during pregnancy. Lead absorbed by the pregnant mother is readily transferred to the developing fetus.¹⁻³ There is evidence from animal studies that intrauterine exposure to lead may disrupt endocrine balance during pregnancy^{4,5} and lead to abnormalities of renal structure and function,⁶⁻⁸ abnormalities of the reproductive system,^{9,10} and neurodevelopmental toxicity¹¹⁻¹³ in offspring. Human evidence corroborates these findings, linking prenatal exposure to lead with reduced birth weight and preterm delivery^{14,15} and with neurodevelopmental abnormalities in offspring.¹⁶⁻²¹ These concerns are especially salient for women and children in developing nations. Not only is exposure to lead common,

but the toxicity of lead for pregnant women and their offspring may be amplified by nutritional deficiency^{22,23} and concomitant toxic exposures²⁴ which often occur in poor nations.

The aim of this study was to find the distribution of PbB among pregnant women in Lucknow, India, a densely populated city of two million in Uttar Pradesh, northern India. The study was prompted by the fact that leaded petrol is still used in Lucknow. Also, we were aware that a traditional Indian eye cosmetic "surma" contains lead, and might pose a hazard for women who use it through exposure to lead. This study was part of a larger ongoing study of the effect of maternal exposure to lead on neonatal birth weight and other outcomes of pregnancy.

Methods

This was a cross sectional study conducted in the urban slums of Lucknow. Under the national healthcare system, "slum" is an administrative designation that, in urban areas, means a population of about 1000 people, with incomes below the national poverty level and suboptimal living conditions. India's integrated child development scheme provides primary care for children and pregnant women in each slum through an "anganwadi" centre. Pregnant women in each slum register at their local centre to receive prenatal care, and are therefore listed. Of Lucknow's 203 slums, 70 were randomly selected and visited by a research assistant at least twice (in the same order) from June 1994 to July 1995. Pregnant women were identified by research assistants from the slum health workers' registers, and were invited to participate in the study. This process continued until 500 subjects were enrolled. Potential subjects were excluded if they declined to participate or if they indicated a likelihood of moving out of the slum in the next six months.

Personal interviews were conducted to collect data on age, reproductive history, and date of last menstrual period, correct to the nearest week. Participants were also asked about other sources of potential exposure to lead, including the source of drinking water, use of surma, and the presence of lead paint at home. Maternal nutritional status was estimated by measuring weight (kg), height (cm), and haemoglobin (g/dl). For measurement of PbB, 5 ml of venous blood was taken after carefully cleaning the skin at the venepuncture site. Analysis of blood lead was performed at the laboratory of the Industrial Toxicology

Department of
Pediatrics and Clinical
Epidemiology Unit,
King George's Medical
College, Lucknow,
India

S Awasthi
V Kumar Pande

State Employees'
Insurance Scheme,
Ministry of Labour,
Government of Uttar
Pradesh, Lucknow,
India

R Awasthi

Industrial Toxicology
Research Center,
Lucknow, India
R C Srivastav

Department of
Environmental and
Occupational Health,
Rollins School of
Public Health of
Emory University,
Atlanta, Georgia, USA
H Frumkin

Correspondence to:
Dr Howard Frumkin,
Department of
Environmental and
Occupational Health, Rollins
School of Public Health of
Emory University, 1518
Clifton Road, Atlanta,
GA 30322, USA.

Accepted 29 July 1996

Concentrations of PbB among pregnant women in Lucknow.

Research Centre (ITRC) in Lucknow, India. Five ml blood was digested in $\text{HNO}_3:\text{HClO}_4$ (6:1) for 24 hours and slowly evaporated to dryness. The process was repeated three times. The analyte was then brought to a volume of 4 ml in 1% HNO_3 . Blood lead was analysed with graphite furnace atomic absorption spectroscopy (GF-AAS; SpectrAA-250 Plus, Varian, Palo Alto, California). To account for any matrix effects present, measurement was effected through the use of a four point standard addition calibration with concentrations developed from a traceable standard (National Physical Laboratory, CSIR, New Delhi). The mean of four replicate analyses were used to assess the concentration. The coefficient of variation of the four analyses was < 2% for all samples. The ITRC laboratory participates in Indian national quality control programmes, and has applied for ISO 9000 certification.

This study was approved by the ethics committee of King George's Medical College, and all subjects gave informed consent before their participation.

After the sampling was completed, we hypothesised that certain neighbourhood characteristics might be associated with variations in the concentration of PbB. We therefore divided the 70 slums into four categories, according to potential sources of environmental exposure to lead. These were (a) the inner city (with high population density and heavy stop and go vehicular traffic), (b) near industrial

sources (factories producing batteries and paint, with medium population density and light vehicular traffic) that could potentially emit lead, (c) near an industry unlikely to emit lead (a large beer distillery, with medium population density and light vehicular traffic), and (d) near roads (with low population density and heavy, fast moving vehicular traffic). Univariate analyses were conducted to characterise the distribution of PbB by demographic, clinical, and exposure related variables.

Results

From the 70 participating slums, 500 pregnant women were enrolled, an average of 7.2 women per slum. The mean PbB was 14.3 (SD 7.9) µg/dl, and the median PbB was 12.9 µg/dl. One hundred and eighty one women (36.2%) had concentrations of PbB ≤ 10 µg/dl, 223 (44.6%) had concentrations 11 to 20 µg/dl, and 96 (19.2%) had concentrations > 20 µg/dl, of whom 25 women had concentrations of ≥ 30 µg/dl. The figure shows the distribution of PbB. Table 1 shows the concentration of PbB against the distribution of various demographic, clinical, and exposure related variables. The concentrations of PbB did not vary with age, weight, height, haemoglobin, gestation, or reproductive history, except that more of the women with high concentrations of lead were multiparous.

Of the 70 participating slums, 36 (51.4%) were inner city slums, nine (11.5%) were near industries that could potentially emit lead, five (6.4%) were near an industry that did not emit lead, and 20 (28.6%) were near highways. Table 2 shows the PbB results from these four categories of neighbourhood. Women in the inner city slums had a mean PbB that was 2.2 (95% confidence interval (95% CI) 0.8 to 3.6) µg/dl higher than the combined concentration of women in the other categories.

As for other potential sources of exposure to lead, 193 subjects (38.6%) reported the use of surma, 376 subjects (75.2%) reported the use

Table 1 Distribution of demographic, clinical, and exposure related variables by concentration of blood lead

	PbB (µg/dl)			Total	P value of ANOVA
	≤ 10	11-20	> 20		
Number (%)	181 (36.2)	223 (44.6)	96 (19.2)	500	
Age (y, mean (SD))	26.4 (4.8)	26.3 (4.6)	26.9 (4.8)	26.5 (4.7)	NS
Weight (kg, mean (SD))	42.9 (6.2)	42.5 (6.5)	43.1 (6.8)	42.79 (6.45)	NS
Height (cm, mean (SD))	149.8 (4.7)	149.9 (4.9)	150.1 (4.8)	149.9 (4.8)	NS
Gestation (weeks, mean (SD))	20.2 (6.6)	19.9 (6.6)	19.5 (6.6)	19.9 (6.6)	NS
Parity (n (%)):					
0	37 (20.4)	47 (19.7)	6 (6.25)	90 (18.0)	
≥ 1	144 (79.6)	176 (78.9)	90 (93.7)	410 (82.0)	
Abortions (n (%)):					
0	156 (86.2)	193 (86.5)	84 (87.5)	433 (86.6)	
≥ 1	25 (13.8)	30 (13.5)	12 (12.5)	67 (13.4)	0.77
Trimester (n (%)):					
1	14 (7.7)	17 (7.6)	9 (9.4)	40	
2	111 (61.3)	136 (60.9)	58 (60.4)	305	
3	56 (30.9)	70 (31.4)	29 (30.2)	155	0.93
Residence (n (%))*:					
1	61 (33.7)	86 (38.6)	50 (52.1)	197	
2	29 (16.0)	53 (23.8)	12 (12.5)	94	
3	29 (16.0)	24 (10.8)	11 (11.5)	64	
4	62 (34.3)	60 (26.9)	23 (23.9)	145	0.0045
Lead paint (n (%))	58 (32.0)	47 (21.1)	23 (23.9)	128	0.06
Piped water (n (%))	131 (72.4)	172 (77.1)	73 (76.0)	376	0.39
Surma (n (%))	78 (43.1)	89 (39.9)	26 (27.1)	193	0.015

*1 = Inner city (with heavy vehicular traffic); 2 = industrial (with potential lead emissions); 3 = industrial (without potential lead emissions); 4 = near highway.

Table 2 Distribution of PbB by type of neighbourhood

Type of neighbourhood	Slums n	Women n	Concentration of PbB ($\mu\text{g/dl}$) (mean (SD))	P value (compared with inner city)
Inner city	36	197	15.65 (8.17)	
Industrial (with potential Pb emissions)	9	94	13.57 (6.40)	0.03
Industrial (without potential Pb emissions)	5	64	12.99 (7.14)	0.01
Near highway	20	145	13.64 (8.40)	0.03

of piped water, and 128 (25.6%) reported the presence of paint on surfaces in their homes. The reported use of surma decreased with increasing PbB, and the reported presence of lead paint showed a trend in the same direction; both findings were unexpected.

Discussion

This study showed increased concentrations of PbB among pregnant women in the slums of Lucknow. In a sample of 500 women, the mean PbB was $14.3 \mu\text{g/dl}$. Ninety six women (19.2%) had PbB $> 20 \mu\text{g/dl}$, of whom 25 had PbB $\geq 30 \mu\text{g/dl}$.

These results confirm previous data showing high concentrations of lead in the Indian population. In the early 1980s, mean (SD) concentrations of PbB among female teachers in various Indian cities were as follows: Ahmedabad $14.7 (7.26)$, Bangalore $19.8 (10.15)$, and Calcutta $10.8 (3.46) \mu\text{g/dl}$.²⁵ At the same time, a population study in Bombay found a mean concentration of PbB to be $35.6 \mu\text{g/dl}$.²⁶ Another recent study of pregnant women in Lucknow found a mean (SD) PbB of $19.4 (8.51) \mu\text{g/dl}$ among those with normal deliveries and $22.52 (8.76) \mu\text{g/dl}$ among those with various adverse outcomes such as preterm delivery, stillbirth, and spontaneous abortion.²⁷ The higher concentrations found in the previous study in Lucknow may be explained by the fact that it was based in hospital, and involved a population selected for high obstetric risk.

These results suggest that Indian women of child bearing age have a higher biological lead burden than their counterparts in other developing countries. Table 3 shows that mean concentrations of PbB in pregnant women in developing nations ranged between 5.5 and $15.2 \mu\text{g/dl}$, with the exception of a small study in Durban, South Africa with higher concentrations. There are several reasons why Indian women might have higher concentrations of PbB. The leading source implicated by our data was motor vehicle traffic as PbB was higher among women in inner city neighbour-

hoods, where traffic exposure is most intense. As our analysis was ecological, and individual women are not likely to spend all of their time in the neighbourhoods where they live, our results are probably biased toward the null, understating the neighborhood effect, and underestimating the impact of lead in petrol. The lead content of Indian petrol remains high, ranging up to 0.56 g/l , and unleaded petrol only reached the Indian market in 1995.³⁵ Leaded petrol is well established as an important factor in concentrations of lead in the population.

Several other environmental sources of lead may affect Indian women, including surma, contaminated water supplies, and lead paints. However, our results did not point to any of these sources, and if anything the results for lead paint and surma showed an unexpected trend away from an association. Before concluding that these sources are harmless in India, further consideration is necessary.

Surma and a related but more finely ground cosmetic kohl have been found to contain lead at concentrations as high as 92%,³⁶⁻³⁹ although specific data on surma from Lucknow are not available. Studies of children in Israel,³⁹ Saudi Arabia,^{40,41} Kuwait,³⁷ the United Kingdom,⁴²⁻⁴⁴ and the United States⁴⁵ have linked the use of these cosmetics with increased PbB, although studies of children in Delhi⁴⁶ and Glasgow⁴⁷ did not find such an association. The presumed routes of absorption in children are conjunctival (the cosmetics are applied directly to the palpebral conjunctiva) and gastrointestinal (when children rub their eyes and lick their fingers). There is also evidence that adults who use surma may develop increased PbB, based on a study of pregnant women in Saudi Arabia.⁴⁸ The negative findings for surma in our study may be explained in several ways: the lead content of local surma may be negligibly low, the impecunious subjects in this study may use surma sparingly or infrequently due to its cost, there may have been exposure misclassification due to misreporting of use of surma, or absorption of lead from local surma may be low. More detailed study of surma in India would help clarify the risk from this common cosmetic.

The negative findings for water supplies may be due to exposure misclassification by subjects unaware of their household water source. It may also indicate low consumption of tap water, or low concentrations of lead in tap water due to limited lead in Lucknow's water distribution system or water that does not dissolve lead due to hardness or alkalinity. Finally, the negative findings for paint are perhaps least surprising. Most homes in Lucknow slums, if they have any painted surfaces at all, only have paint on limited areas such as door frames. Moreover, as leaded paint is more expensive than alternatives, much of the paint used in homes is probably lead free. Therefore, the high prevalence of reported lead paint (25.6%) probably reflects misclassification.

The toxicokinetics of lead in pregnancy are complex.⁴⁹ Previous reports have noted a fall

Table 3 Previously reported PbB among pregnant women in developing nations

Location	Subjects	PbB ($\mu\text{g/dl}$) (mean (SD))	Reference (year)
Riyadh	124	5.5 (2.6)	28 (1995)
Bangkok	500	6.2 (2.0)	29 (1994)
Mexico City:			
12 weeks	250	9.0 (6.0)	30 (1993)
At term	250	8.5 (6.2)	30 (1993)
Durban	21	21.7 (2.9)	31 (1993)
Shanghai	165	14.1 (6.8)	32 (1990)
Taiwan	147	6.5 (0.2)	33 (1991)
Kuala Lumpur	114	15.2 (4.2)	34 (1985)

in PbB during the second trimester of pregnancy,⁵⁰ attributed to haemodilution caused by the expansion of blood volume.⁵¹ Others⁵² have found an increase in PbB during pregnancy, possibly due to the release of lead from bone stores.⁴⁹ In our population, the PbB did not vary with the trimester of pregnancy. This may reflect the unusually high concentrations of PbB, or it may be related to deficiencies of iron, calcium, or zinc, which facilitate intestinal absorption of lead⁵² and mobilisation of bone lead during pregnancy.⁴⁹⁻⁵³ The effect of pregnancy on lead toxicokinetics needs further investigation.

A weakness of this study is that we did not attempt to ascertain exposures to other potential sources of lead. The most important of these are direct or secondary occupational exposures. Consumer goods with paint containing lead, contaminated food, and contaminated beverages are other potential sources. Future studies should attempt to ascertain the importance of these exposures.

The blood lead concentrations in the women in this study were increased to an alarming extent. In combination with poor nutrition, infectious diseases, and other disadvantages associated with poverty, the exposure to lead threatens the normal development of children. Aggressive approaches to primary prevention are needed, including the removal of lead from petrol, an intervention known to decrease concentrations of lead in the population.⁵⁴ Further research is necessary to identify other sources of exposure to lead, such as in consumer products and cosmetics, and to eliminate these as well.

This work was supported by a grant from the Environmental and Occupational Health Initiative of the International Clinical Epidemiology Network (INCLEN), from the Rockefeller Foundation.

- 1 Buchet JP, Roels H, Hubermont G, Lauwerys R. Placental transfer of lead, mercury, cadmium, and carbon monoxide in women. II. Influence of some epidemiological factors on the frequency distributions of the biological indices in maternal and umbilical cord blood. *Environ Res* 1978;15:494-503.
- 2 Roels H, Hubermont G, Buchet JP, Lauwerys R. Placental transfer of lead, mercury, cadmium, and carbon monoxide in women. III. Factors influencing the accumulation of heavy metals in the placenta and the relationship between metal concentration in the placenta and in maternal and cord blood. *Environ Res* 1978;16:236-47.
- 3 Ong CN, Phoon WO, Law HY, Tye CY, Lim HH. Concentrations of lead in maternal blood, cord blood, and breast milk. *Arch Dis Child* 1985;60:756-9.
- 4 Foster WG. Reproductive toxicity of chronic lead exposure in the female cynomolgus monkey. *Reprod Toxicol* 1992;6:123-31.
- 5 Sierra EM, Tiffany-Castiglioni E. Effects of low-level lead exposure on hypothalamic hormones and serum progesterone levels in pregnant guinea pigs. *Toxicology* 1992;72:89-97.
- 6 Vyskocil A, Cizkova M, Tejnoroova I. Effect of prenatal and postnatal exposure to lead on kidney function in male and female rats. *J Appl Toxicol* 1995;15:327-8.
- 7 Oberley TD, Friedman AL, Moser R, Siegel FL. Effects of lead administration on developing rat kidney. II. Functional, morphologic, and immunohistochemical studies. *Toxicol Appl Pharmacol* 1995;131:94-107.
- 8 Moser R, Oberley TD, Daggett DA, Friedman AL, Johnson JA, Siegel FL. Effects of lead administration on developing rat kidney. I. Glutathione S-transferase isoenzymes. *Toxicol Appl Pharmacol* 1995;131:85-93.
- 9 Coffigny H, Thoreux-Manlay A, Pinon-Lataillade G, Monchaux G, Masse R, Soufir JC. Effects of lead poisoning of rats during pregnancy on the reproductive system and fertility of their offspring. *Hum Exp Toxicol* 1994;13:241-6.
- 10 Corpas I, Gaspar I, Martinez S, Codesal J, Candelas S, Antonio MT. Testicular alterations in rats due to gestational and early lactational administration of lead. *Reprod Toxicol* 1995;9:307-13.
- 11 Legare ME, Castiglioni AJ Jr, Rowles TK, Calvin JA, Snyder-Armstead C, Tiffany-Castiglioni E. Morphological alterations of neurons and astrocytes in guinea pigs exposed to low levels of inorganic lead. *Neurotoxicology* 1993;14:77-80.
- 12 Buchheim K, Noack S, Stoltenburg G, Lilienthal H, Winneke G. Developmental delay of astrocytes in hippocampus of rhesus monkeys reflects the effect of pre- and postnatal chronic low level lead exposure. *Neurotoxicol* 1994;15:665-9.
- 13 Tang HW, Liang YX, Hu XH. Effects of low level lead exposure on behavior of young rats. *Chung Kuo Yao Li Hsueh Pao* 1994;15:316-9.
- 14 Tabacova S, Balabaeva L. Environmental pollutants in relation to complications of pregnancy. *Environ Health Perspect* 1993;101(suppl 2):27-31.
- 15 Andrews KW, Savitz DA, Hertz-Picciotto I. Prenatal lead exposure in relation to gestational age and birth weight: a review of epidemiologic studies. *Am J Ind Med* 1994;26:13-32.
- 16 Bellinger D, Leviton A, Waternaux C, Needleman H, Rabinowitz M. Longitudinal analysis of prenatal and postnatal lead exposure and early cognitive development. *N Engl J Med* 1987;316:1037-43.
- 17 Dietrich K, Krafft KM, Bornschein RJ, Hammond PB, Berger O, Succop PA, Bier M. Low level fetal lead exposure effects on neurobehavioral development in early infancy. *Pediatrics* 1987;80:721-30.
- 18 Huel G, Tubert P, Frery N, Moreau T, Dreyfus J. Joint effect of gestational age and maternal lead exposure on psychomotor development of the child at six years. *Neurotoxicology* 1992;13:249-54.
- 19 Leviton A, Bellinger D, Allred EN, Rabinowitz M, Needleman H, Schoenbaum S. Pre- and postnatal low-level lead exposure and children's dysfunction in school. *Environ Res* 1993;60:30-43.
- 20 Bellinger D, Leviton A, Allred E, Rabinowitz M. Pre- and postnatal lead exposure and behavior problems in school-aged children. *Environ Res* 1994;66:12-30.
- 21 Rothenberg SJ, Poblano A, Garza-Morales S. Prenatal and perinatal low level lead exposure alters brainstem auditory evoked responses in infants. *Neurotoxicology* 1994;15:695-9.
- 22 West WL, Knight EM, Edwards CH, Manning M, Spurlock B, James H, et al. Maternal low level lead and pregnancy outcomes. *J Nutr* 1994;124(suppl 6):981S-6S.
- 23 Bogden JD, Kemp FW, Han S, Murphy M, Fraiman M, Czerniach D. Dietary calcium and lead interact to modify maternal blood pressure, erythropoiesis, and fetal and neonatal growth in rats during pregnancy and lactation. *J Nutr* 1995;125:990-1002.
- 24 Kristensen P, Eilertsen E, Einarsdottir E, Haugen A, Skaug V, Ovrebo S. Fertility in mice after prenatal exposure to benzo[a]pyrene and inorganic lead. *Environ Health Perspect* 1995;103:588-90.
- 25 Parikh DJ. Human exposure to lead: an overview in India. *Indian Journal of Industrial Medicine* 1990;36:63-75.
- 26 Annapurna V, Mahajan BA, Murti MV. Studies of blood lead levels in Bombay's citizens. *Indian Journal of Environmental Studies* 1985;24:239-43.
- 27 Saxena DK, Singh C, Murthy RC, Mathur N, Chandra SV. Blood and placental lead levels in an Indian city: a preliminary report. *Arch Environ Health* 1994;49:106-10.
- 28 Al-Saleh I, Khalil MA, Taylor A. Lead, erythrocyte protoporphyrin, and hematological parameters in normal maternal and umbilical cord blood from subjects of Riyadh region, Saudi Arabia. *Arch Environ Health* 1995;50:66-73.
- 29 Phuapradit W, Jetsawangsi T, Chaturachinda K, Noinongyao N. Maternal and umbilical cord blood lead levels in Ramathibodi Hospital, 1993. *J Med Assoc Thai* 1994;77:368-72.
- 30 Fuentes-Aguilar L, Soto-Mora C. Influence of lead on pregnant women in metropolitan Mexico City. *Bull Environ Contam Toxicol* 1993;50:533-9.
- 31 Chetty N, Jinabhai CC, Green-Thompson RW. Lead levels in maternal and umbilical cord blood at King Edward Hospital, Durban [letter]. *S Afr Med J* 1993;83:227.
- 32 Li PJ, Sheng YZ, Wang QY, Gu LY. Transfer of lead via placenta and milk in humans. *Industrial Health and Occupational Disease* 1990;16:207-10.
- 33 Soong Y-K, Tseng R, Liu C, Lin P-W. Lead, cadmium, arsenic, and mercury levels in maternal and fetal cord blood. *J Formos Med Assoc* 1991;90:59-65.
- 34 Aslam M, Davis SS, Healy MA. Heavy metals in some Asian medicines and cosmetics. *Public Health* 1979;93:274-84.
- 35 Thomas V, Spiro G. Emissions and exposure to metals: cadmium and lead. In: Socolow RW, ed. *Industrial ecology and global change*. New York: Cambridge University Press, 1995.
- 36 Aslam M, Healy MA, Davis SS, Ali AR. Surma and blood lead in children. *Lancet* 1980;i:658-9.
- 37 Fernando NP, Healy MA, Aslam M, Davis SS, Hussein A. Lead poisoning and traditional practices: the consequences for world health. A study in Kuwait. *Public Health* 1981;95:250-60.
- 38 Parry C, Eaton J. Kohl: a lead-hazardous eye makeup from the third world to the first world. *Environ Health Perspect* 1991;94:121-3.
- 39 Nir A, Tamir A, Zelnik N, Iancu TC. Is eye cosmetic a source of lead poisoning? *Isr J Med Sci* 1992;28:417-21.

- 40 Abdullah MA. Lead poisoning among children in Saudi Arabia. *Journal of Tropical Medical Hygiene* 1984;87: 67-70.
- 41 Alkhawajah AM. Alkohl use in Saudi Arabia. Extent of use and possible lead toxicity. *Trop Geogr Med* 1992;44:373-7.
- 42 Warley MA, Blackledge P, O'Gorman P. Lead poisoning from eye cosmetic. *BMJ* 1968;i:117.
- 43 Snodgrass GJAI, Ziderman DA, Gulati V, Richards J. Cosmetic plumbism. *BMJ* 1973;4:230.
- 44 Ali AR, Smales ORC, Aslam M. Surma and lead poisoning. *BMJ* 1978;ii:915-6.
- 45 Sprinkle RV. Leaded eye cosmetics: a cultural cause of elevated lead levels in children. *J Fam Pract* 1995;40: 358-62.
- 46 Gogte ST, Basu N, Sinclair S, Ghai OP, Bhide NK. Blood lead levels of children with pica and surma use. *Indian J Pediatr* 1991;58:513-9.
- 47 Attenburrow AA, Campbell S, Logan RW, Goel KM. Surma and blood lead levels in Asian children in Glasgow. *Lancet* 1980;i:323.
- 48 Moghraby SA, Abdullah MA, Karrar O, Akiel AS, Shawaf TA, Majid YA. Lead concentrations in maternal and cord blood in women users of surma eye cosmetics. *Ann Trop Paediatr* 1989;9:49-53.
- 49 Silbergeld EK. Lead in bone: implications for toxicology during pregnancy and lactation. *Environ Health Perspect* 1991;91:63-70.
- 50 Rothenberg SJ, Karchmer S, Schnaas L, Perroni E, Zea F, Alba JF. Changes in serial blood levels during pregnancy. *Environ Health Perspect* 1994;102:876-80.
- 51 Mahaffey KR. Biokinetics of lead during pregnancy. *Fundam Appl Toxicol* 1991;16:15-6.
- 52 Jacobssen LB, Soderberg HA, Nordberg GF, Ekesrydh S, Englyst V. Biological monitoring of arsenic, lead and cadmium in occupationally and environmentally exposed pregnant women. *Scand J Work Environ Health* 1993;19: 50-3.
- 53 Fullmer CS. Intestinal interaction of lead and calcium. *Neurotoxicol* 1992;13:799-808.
- 54 Pirkle JL, Brody DJ, Gunter EW, Kramer RA, Paschal DC, Flegal KM, Matte TD. The decline in blood lead levels in the United States. *JAMA* 1994;272:284-91.

Rejected manuscripts

From February 1994, authors whose submitted articles are rejected will be advised of the decision and one copy of the article, together with any reviewers' comments, will

be returned to them. The *Journal* will destroy remaining copies of the article but correspondence and reviewers' comments will be kept.