ORIGINAL ARTICLE

Impact of ambient air pollution on birth weight in Sydney, Australia

T Mannes, B Jalaludin, G Morgan, D Lincoln, V Sheppard, S Corbett

Occup Environ Med 2005;62:524–530. doi: 10.1136/oem.2004.014282

See end of article for authors' affiliations

Correspondence to:
Ms T F Mannes, NSW
Public Health Officer
Training Program, New
South Wales Health
Department, LMB 961
North Sydney, Australia;
paman@doh.health.nsw.
gov.au

Accepted
24 February 2005

Background: Studies in Asia, Europe, and the Americas have provided evidence that ambient air pollution may have an adverse effect on birth weight, although results are not consistent.

Methods: Average exposure during pregnancy to five common air pollutants was estimated for births in metropolitan Sydney between 1998 and 2000. The effects of pollutant exposure in the first, second, and third trimesters of pregnancy on risk of "small for gestational age" (SGA), and of pollutant exposure during pregnancy on birth weight were examined.

Results: There were 138 056 singleton births in Sydney between 1998 and 2000; 9.7% of babies (13 402) were classified as SGA. Air pollution levels in Sydney were found to be quite low. In linear regression models carbon monoxide and nitrogen dioxide concentrations in the second and third trimesters had a statistically significant adverse effect on birth weight. For a 1 part per million increase in mean carbon monoxide levels a reduction of 7 (95% CI –5 to 19) to 29 (95% CI 7 to 51) grams in birth weight was estimated. For a 1 part per billion increase in mean nitrogen dioxide levels a reduction of 1 (95% CI 0 to 2) to 34 (95% CI 24 to 43) grams in birth weight was estimated. Particulate matter (diameter less than ten microns) in the second trimester had a small statistically significant adverse effect on birth weight. For a 1 microgram per cubic metre increase in mean particulate matter levels a reduction of 4 grams (95% CI 3 to 6) in birth weight was estimated.

Conclusion: These findings of an association between carbon monoxide, nitrogen dioxide, and particulate matter, and reduction in birth weight should be corroborated by further study.

Epidemiological studies addressing the relation between ambient air pollution and fetal development are accumulating worldwide. Studies conducted in China,¹ the Czech Republic,^{2,3} Korea,⁴ the UK,⁵ Brazil,⁶ and North America^{7–11} have examined the link between ambient air pollution levels during pregnancy and reduction in birth weight or intrauterine growth retardation (IUGR). There is evidence to suggest that air pollutant exposure during pregnancy has an adverse impact on birth weight, although the findings are inconsistent; the effect of individual pollutants and the period(s) during pregnancy when pollutant levels are likely to have most impact on birth weight is not clear.

It is useful, then, to examine the impact of ambient air pollution and birth weight in a variety of different sites to clarify the nature of the relation. We examined this relation in Sydney as there is complete, routinely collected data available. In the present study we evaluated the effect of prenatal exposure (in early, mid, and late pregnancy) to five common urban air pollutants: particulate matter (PM₁₀ and PM_{2.5}), carbon monoxide (CO), nitrogen dioxide (NO₂), and ozone (O₃) from routine air monitoring in the Sydney metropolitan area on birth weight.

METHODS

Information on all births in metropolitan Sydney between 1 January 1998 and 31 December 2000 was obtained from the Midwives Data Collection (MDC) at the New South Wales Department of Health. The MDC is a population based surveillance system covering all live births and stillbirths of at least 20 weeks gestation and at least 400 grams birth weight. Birth data includes maternal demographic factors (age, smoking status, country of birth, postcode of residence at time of delivery), pregnancy factors (date of the last menstrual period, gestational hypertension and diabetes,

parity, time of first antenatal visit to a healthcare provider), details about the delivery (type of delivery), and infant factors (birth weight, gestational age). Other known causes of low birth weight including multiple births, hypertension of pregnancy, and gestational diabetes were excluded.

We obtained daily air pollution data from the New South Wales Environment Protection Authority (EPA) and daily meteorological data from the Australian Bureau of Meteorology. Air pollution data included concentrations of PM₁₀, PM_{2.5}, NO₂, O₃, and CO recorded at 14 monitoring stations in metropolitan Sydney. Monitoring stations were excluded if less than 80% of readings were available for each pollutant. Pollutant concentrations were analysed as continuous variables.

For each birth, exposure to each air pollutant during gestation was estimated by calculating the average of each pollutant over 30 days (last month) and 90 days (third trimester) before birth, the mid 90 days of gestation (second trimester), and 90 days (first trimester) after the estimated date of conception. Pollutant concentrations from monitoring stations were averaged to provide an estimate for the whole of metropolitan Sydney, using an approach similar to the APHEA2 (Short-term Effects of Air Pollution and Health: A European Approach) studies.^{12–14} The process was repeated

Abbreviations: ABS, Australian Bureau of Statistics; APHEA, Short-term Effects of Air Pollution and Health: A European Approach; CI, confidence interval; EPA, Environment Protection Authority; IRSD, Index of Relative Socioeconomic Disadvantage; IUGR, intrauterine growth retardation; LBW, low birth weight; LMP, last menstrual period; MDC, midwives data collection; NSW, New South Wales; PM_{2.5}, particulate matter less than 2.5 microns diameter; PM₁₀, particulate matter less than 10 microns diameter; ppb, parts per billion (10^{–9}); ppm, parts per million (10^{–6}); SD, standard deviation; SE, standard error; SES, socioeconomic status; SGA, small for gestational age; OR, odds ratio; Temp, temperature; TSP, total suspended particulates

Table 1 Characteristics of all singleton births in Sydney from 1998 to 2000 by "small for gestational age"

Variable	Small for gestational age (n = 13 402)		Not small for gestational age (n = 138 056)		Adjusted OR (95% CI)
	No.	%	No.	%	
Maternal age (years)					
<24	2762	20.6	19460	14.1	1.02 (0.97 to 1.08)
25–34*	8340	62.2	93828	68.0	1.00
35–44	2285	17.0	24639	17.8	1.01 (0.95 to 1.01)
>45	15	0.1	129	0.1	1.55 (0.84 to 2.84)
Maternal smoker					
Yes	2659	19.8	14254	10.3	1.97 (1.87 to 2.08)
No	10741	80.1	110375	79.9	
Sex of infant					
Female	6456	48.2	60381	43.7	1.01 (0.97 to 1.05)
Male	6946	51.8	64273	46.6	
Maternal Aboriginality					
Yes	117	0.9	806	0.6	1.03 (0.83 to 1.28)
No	13275	99.1	123758	89.6	
Gestation at first antenatal visit					
>20 weeks	1803	13.5	13854	10.0	1.12 (1.05 to 1.18)
≤20 weeks	11486	85.7	110014	79.7	
Previous pregnancy					
No	7427	55.4	53102	38.5	0.57 (0.54 to 0.59)
Yes	5973	44.6	71517	51.8	
IRSD quintile					
5 (least disadvantaged; high SES)*	3843	28.7	58638	42.5	1.00
4	2857	21.3	27395	19.8	1.24 (1.17 to 1.13)
3	2374	17.7	19997	14.5	1.38 (1.30 to 1.46)
2	1684	12.6	13826	10.0	1.50 (1.41 to 1.61)
1 (most disadvantaged; low SES)	2644	19.7	18200	13.2	1.73 (1.63 to 1.83)
Season of birth					
Summer (Dec–Feb)*	3265	24.4	43727	31.7	1.00
Autumn (Mar–May)	3456	25.8	31109	22.5	1.05 (0.99 to 1.11)
Winter (Jun–Aug)	3375	25.2	31469	22.8	1.00 (0.95 to 1.06)
Spring (Sep–Nov)	3306	24.7	31751	23.0	0.99 (0.94 to 1.05)

*Referent group.

OR, odds ratio adjusted for other covariates; SD, standard deviation; SES, socioeconomic status.

Note that in Australia, autumn is in March, April, May; winter is in June, July, August; spring is in September, October, November; and summer is in December, January, February.

matching air pollution concentrations from each eligible monitoring station and births to mothers residing in postcodes within 5 km of the monitoring station.

The following covariates were included in regression models: sex of child, maternal age (in year groupings), gestational age (only included in linear regression models), maternal smoking (yes/no), gestational age at first antenatal visit (≤ 20 weeks or >20 weeks), maternal indigenous status (whether mother identifies as being Aboriginal or Torres Strait Islander), whether first pregnancy, season of birth, and socioeconomic status (SES). SES was measured using the Index of Relative Socioeconomic Disadvantage (IRSD) of postcode of maternal residence. The Australian Bureau of Statistics (ABS) constructs the IRSD to classify geographical areas on the basis of social and economic information collected in the population census. Each postcode in New South Wales (population of approximately 1000–5000 people) is assigned an IRSD index. We then ranked the postcodes and divided the list into quintiles.

A variable "small for gestational age" (SGA) was calculated based on the Australian national birth weight centile for gestational age 1991–94.¹⁵ SGA was defined as greater than two standard deviations below the mean birth weight according to gestational age as SGA. SGA was analysed as a categorical variable in logistic regression models using the SAS System for Windows v8.02. Linear regression models were also developed using birth weight as a continuous variable. Following the development of the basic model, air pollutants were then added to the model to determine the association between ambient air pollutants and birth weight. Single and multi-pollutant pollutant models were assessed and we investigated interactions between air pollution

variables and covariates. The impact of pollutant exposures in other pregnancy periods on key findings was also analysed.

RESULTS

There were 138 056 singleton births in Sydney between 1998 and 2000; 9.7% of babies (13 402) were classified as SGA. Mean (SD) birth weight for babies born in Sydney between 1998 and 2000 was 3418 (531) grams. After adjusting for other maternal and infant characteristics, SGA was significantly associated with maternal smoking, gestational age greater than 20 weeks at first antenatal visit, and first pregnancy (table 1). All SES variables exhibit a statistically significant association with SGA. Compared to the high SES category, there was increasing risk of SGA with decreasing SES.

Pollutant concentrations in Sydney from 1 April 1997 to 31 December 2000 are presented in table 2. A correlation matrix of air pollution variables is presented in table 3 to show correlations between pollutants. Correlation coefficients above 0.8 were observed between PM_{10} and $PM_{2.5}$. Correlation coefficients were also calculated to determine the correlations between air monitoring stations. Correlation coefficients between the 14 monitoring stations ranged from 0.68 to 0.85 for CO. Correlation coefficients between the 14 monitoring stations ranged from 0.46 to 0.85 for NO_2 ; from 0.53 to 0.94 for O_3 ; from 0.67 to 0.91 for PM_{10} ; and from 0.66 to 0.93 for $PM_{2.5}$. All pollutants differed according to season ($p < 0.01$) and post hoc analysis revealed that all seasons were different to all others ($p < 0.05$). Concentrations of CO, NO_2 , and $PM_{2.5}$ were highest in winter and lowest in summer. Concentrations of PM_{10} and O_3 were highest in summer and lowest in winter.

Table 2 Daily average pollutant concentrations for air monitoring stations in Sydney, 1 April 1997–31 December 2000

	Australian National Standard	Mean (SD)	Minimum	25th centile	Median	75th centile	Maximum
Pollutant							
PM ₁₀ (µg/m ³)							
24 hour av	50 µg/m ³	16.8 (7.1)	3.8	12.3	15.7	19.9	104.0
PM _{2.5} (µg/m ³)							
24 hour av	25 µg/m ³ *	9.4 (5.1)	2.4	6.5	8.4	11.2	82.1
CO (ppm)							
8 hour av	9.0 ppm	0.8 (0.7)	0.0	0.4	0.6	1.1	4.6
O ₃ (ppb)							
1 hour max	100 ppb	31.6 (14.6)	3.2	22.7	27.7	36.3	126.7
NO ₂ (ppb)							
1 hour max	120 ppb	23.2 (7.4)	56.2	18.0	23.0	27.5	59.4

*Advisory reporting standard from 2003.

ppb, parts per billion (10⁻⁹); ppm, parts per million (10⁻⁶); CO, carbon monoxide, 8 hour average in ppm; O₃, ozone, 1 hour max in ppb; PM₁₀, particulate matter less than 10 microns, 24 hour average in µg/m³; PM_{2.5}, particulate matter less than 2.5 microns, 24 hour average in µg/m³; NO₂, nitrogen dioxide, 24 hour average in ppb.

Citywide average air pollutant concentrations in the last month, third trimester, and first trimester of pregnancy had no statistically significant effect on SGA after adjusting for infant and maternal characteristics (table 4). Concentrations of PM₁₀, PM_{2.5}, and O₃ in the second trimester of pregnancy had a small but statistically significant adverse effect on SGA (OR 1.01, 95% CI 1.00 to 1.04; OR 1.03, 95% CI 1.01 to 1.05; and OR 1.01, 95% CI 1.00 to 1.01, respectively).

Eleven metropolitan air monitoring stations were eligible for inclusion in the analysis of babies born to women residing within 5 km of an air monitoring station (table 4). There were 51 460 eligible births in 1998–2000. Of these, 5985 (11.6%) were classified as SGA. NO₂ concentrations in the second and third trimesters and one month before birth had a statistically significant adverse effect on risk of SGA (ORs between 1.07 and 1.14) (table 4). The NO₂ effects were robust to controlling for exposures in other pregnancy periods (ORs between 1.1 (95% CI 1.0 to 1.2) and 1.1 (95% CI 1.0 to 1.2) for second and third trimesters respectively). Exposure to PM₁₀ in the second trimester of pregnancy had a small but statistically significant adverse effect on risk of SGA (OR 1.02, 95% CI 1.01 to 1.03) (table 4), although these findings were not robust to analysis when controlling for exposures to PM₁₀ in other periods of pregnancy (OR 1.01, 95% CI 0.99 to 1.02).

In multivariate linear regression models, the effect of pollutant concentrations on birth weight (in grams) was examined for all births in Sydney during the study period and for babies born to women residing within 5 km of an air monitoring station (table 5). Citywide average levels of PM₁₀ and PM_{2.5} in the second trimester and last month of pregnancy had a small but statistically significant adverse effect on birth weight. When analysing only babies born to

women residing within 5 km of an air monitoring station, we observed a small adverse effect of PM₁₀ exposure during the second trimester on birth weight (regression coefficient −4.3, 95% CI −5.8 to −2.8). Thus, for every 1 µg/m³ increase in 24 hour average PM₁₀ during the second trimester of pregnancy, a 4 gram reduction in birth weight was estimated. This finding persisted after controlling for exposures in other pregnancy periods and analysis in multi-pollutant models (table 6).

Citywide average levels of CO in the last month of pregnancy had a statistically significant adverse effect on birth weight, whereas the 5 km results reveal an adverse effect of CO exposure on birth weight for the second and third trimesters of pregnancy. Linear regression coefficients for the 5 km analysis ranged from −23 (95% CI −44.6 to −1.2) to −29 (95% CI −51.0 to −6.8); thus, for every 1 part per million increase in 8 hour average CO during pregnancy, a 23–29 gram reduction in birth weight was estimated. Analysis of second trimester CO findings in two and three pollutant models and controlling for exposures in other pregnancy periods are presented in table 6. CO findings in the second trimester did not remain statistically significant when analysed in multi-pollutant models.

Citywide average levels of NO₂ in the first and third trimesters of pregnancy resulted in a small but statistically significant adverse effect on birth weight, whereas the 5 km results reveal an adverse effect of NO₂ exposure on birth weight for all periods of pregnancy. Linear regression coefficients for the 5 km analysis ranged from −20 (95% CI −27.8 to −11.5) to −34 (95% CI −43.4 to −24.3). NO₂ findings in the second trimester remained when analysed in two pollutant models, but did not remain when analysed in the four pollutant model (table 6).

Table 3 Matrix of Pearson correlation coefficients of pollutant concentrations (averaged across all air monitoring stations) in Sydney, April 1997–December 2000, and birth outcomes for term births in Sydney, 1998–2000

Pollutant	Birth weight	CO	NO ₂	O ₃	PM ₁₀	PM _{2.5}	Temp
Birth weight	1.00						
CO	0.00	1.00					
NO ₂	0.01	0.57	1.00				
O ₃	0.01	−0.20	0.29	1.00			
PM ₁₀	0.01	0.26	0.47	0.52	1.00		
PM _{2.5}	0.01	0.53	0.66	0.36	0.81	1.00	
Temp	0.00	−0.42	−0.16	0.60	0.38	0.05	1.00

Temp, temperature; ppb, parts per billion (10⁻⁹); ppm, parts per million (10⁻⁶); CO, carbon monoxide, 8 hour average in ppm; O₃, ozone, 1 hour max in ppb; PM₁₀, particulate matter less than 10 microns, 24 hour average in µg/m³; PM_{2.5}, particulate matter less than 2.5 microns, 24 hour average in µg/m³; NO₂, nitrogen dioxide, 24 hour average in ppb.

Table 4 Association (OR point estimate, 95% CI) between pollutant concentrations in Sydney and risk of small for gestational age (SGA) for infants born in Sydney between 1998 and 2000 by exposure period									
Pollutant	No. of infants (SGA/total)	Exposure period							
		One month before birth		Third trimester		Second trimester		First trimester	
		OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
All births									
PM ₁₀	13402/138056	1.01	1.00 to 1.03	1.00	0.99 to 1.013	1.01	1.00 to 1.04	1.00	0.98 to 1.02
PM _{2.5}	13402/138056	1.01	0.99 to 1.03	0.99	0.97 to 1.02	1.03	1.01 to 1.05	0.99	0.97 to 1.01
CO	13402/138056	1.06	0.98 to 1.16	1.01	0.91 to 1.11	0.99	0.90 to 1.10	0.95	0.88 to 1.04
NO ₂	13402/138056	1.00	1.00 to 1.01	1.01	1.00 to 1.02	1.00	0.99 to 1.01	1.00	0.99 to 1.01
O ₃	13402/138056	1.00	0.99 to 1.01	1.00	1.00 to 1.01	1.01	1.00 to 1.01	1.00	1.00 to 1.01
5 km births									
PM ₁₀	5391/44891	1.00	0.99 to 1.02	1.01	0.99 to 1.02	1.02	1.01 to 1.03	1.01	0.99 to 1.02
PM _{2.5}	1595/13855	1.01	0.97 to 1.04	1.00	0.95 to 1.05	1.00	0.96 to 1.05	0.99	0.94 to 1.04
CO	2892/22684	1.10	0.96 to 1.27	1.05	0.90 to 1.23	1.06	0.90 to 1.25	0.99	0.86 to 1.14
NO ₂	5985/51460	1.07	1.00 to 1.14	1.13	1.05 to 1.21	1.14	1.07 to 1.22	1.06	0.99 to 1.14
O ₃	5460/45730	1.01	0.97 to 1.06	1.01	0.96 to 1.07	1.00	0.95 to 1.06	0.99	0.93 to 1.03
OR estimates adjusted for: maternal age, maternal smoking, indigenous status, SES, gestational age at first antenatal visit, season of birth, and parity. OR per 1 unit increase in pollutant concentration. ppb, parts per billion (10 ⁻⁹); ppm, parts per million (10 ⁻⁶). Units and averaging periods: PM ₁₀ (µg/m ³): 24 hour av PM _{2.5} (µg/m ³): 24 hour av CO (ppm): 8 hour av O ₃ (ppb): 1 hour max NO ₂ (ppb): 1 hour max									

DISCUSSION

We examined associations between birth weight and exposure to air pollution concentrations at various stages of pregnancy across all of Sydney and for those infants born to women residing within 5 km of air monitoring stations. We showed a small effect of PM₁₀ concentration in the second trimester of pregnancy, on birth weight in both linear and logistic regression models. We also showed that CO and NO₂ concentrations in the second and third trimesters of pregnancy (most pronounced in the second trimester), had a statistically significant effect on birth weight. The effect of NO₂ concentrations was only seen when examining births to women residing within 5 km of an air monitoring station. The effect of CO concentrations was only seen in the linear regression analysis and was most pronounced when examining births to women residing within 5 km of an air monitoring station.

There are a number of strengths to our study. We were able to investigate the association between low birth weight and pollutant concentrations for a large number of births in metropolitan Sydney between 1998 and 2000, and we were able to test this association using information on infants born to women residing within 5 km of an air monitoring station, again for a large number of births. We were also able to analyse the effect of five common air pollutants (PM₁₀, PM_{2.5}, CO, NO₂, and O₃), as these pollutants are routinely monitored at many air monitoring stations in metropolitan Sydney. One strength of this study is that we have been able to control for maternal smoking, indigenous status, gestational age at first antenatal visit, and parity among other potential confounders.

Although our analysis controlled for a number of important potential confounders, we did not have the information to adjust for some known risk factors for low birth weight; for example, maternal nutrition, maternal occupation, or pre-pregnancy maternal weight. Also, residual confounding by SES cannot be ruled out. Our SES estimate is a group level variable assigned to postcode of maternal residence at time of birth.

The most important source of bias in our study is due to measurement of exposure. By using air pollutant concentrations averaged across Sydney, we assume that ambient pollutant concentrations represent an individual's actual exposure to pollutants. This assumption does not account for time-activity patterns that may mediate exposure such as commuting habits, place or type of work, or time spent outdoors.

The use of citywide average exposure does not account for variations in pollutant concentrations across Sydney or even within the 5 km zone around an air monitoring station. It is likely that pollutants, particularly primary pollutants such as nitrogen dioxide and carbon monoxide, are not homogeneously distributed across Sydney. The distribution of primary pollutants will largely depend on the presence of combustion sources such as roads or industry. Thus we also examined the impact of pollutant concentrations only for births to women residing within 5 km of an air monitoring station to attempt to provide a better estimate of exposure for pollutants with large geographic variability. Analysis of the effect of CO and NO₂ shows a clear pattern of increased effect when examining only births to

Table 5 Changes in birth weight (in grams) for a 1 unit change in exposure to air pollutants (PM₁₀ and PM_{2.5} in µg/m³; CO in ppm; O₃, NO₂ in ppb) at each trimester, and last month of pregnancy; results of linear regression models adjusted for covariates

Pollutant	Exposure period				First trimester				Second trimester				Third trimester				Multiple linear regression coefficient				95% confidence limits			
	One month before birth				Multiple linear regression coefficient				Multiple linear regression coefficient				Multiple linear regression coefficient				Multiple linear regression coefficient				Multiple linear regression coefficient			
	Multiple linear regression coefficient				95% confidence limits				Multiple linear regression coefficient				Multiple linear regression coefficient				Multiple linear regression coefficient				Multiple linear regression coefficient			
All births																								
PM ₁₀	-1.21	-2.31	-0.11	-0.95	-2.30	-0.40	-2.05	-3.36	-0.74	-0.14	-1.37	1.09	-0.95	-2.30	-0.40	-2.05	-3.36	-0.74	-0.14	-1.37	1.09			
PM _{2.5}	-2.48	-4.58	-0.38	-0.98	-3.74	1.78	-4.10	-6.79	-1.41	0.36	-2.29	3.01	-0.98	-3.74	1.78	-4.10	-6.79	-1.41	0.36	-2.29	3.01			
CO	-15.28	-25.59	-4.97	-6.63	-18.57	5.31	-10.72	-23.09	1.65	1.86	-8.31	12.03	-6.63	-18.57	5.31	-10.72	-23.09	1.65	1.86	-8.31	12.03			
NO ₂	-0.76	-1.72	0.20	-1.48	-2.70	-0.26	-0.95	-2.07	0.17	-1.07	-2.07	-0.07	-1.48	-2.70	-0.26	-0.95	-2.07	0.17	-1.07	-2.07	-0.07			
O ₃	-0.11	-0.56	0.34	-0.45	-1.08	0.18	-0.75	-1.38	-0.12	-0.09	-0.66	0.48	-0.45	-1.08	0.18	-0.75	-1.38	-0.12	-0.09	-0.66	0.48			
5 km births																								
PM ₁₀	-2.98	-4.25	-1.71	-3.84	-5.35	-2.33	-4.28	-5.79	-2.77	-2.57	-4.04	-1.10	-3.84	-5.35	-2.33	-4.28	-5.79	-2.77	-2.57	-4.04	-1.10			
PM _{2.5}	-2.70	-6.80	1.40	-2.83	-9.00	3.34	-1.54	-4.59	7.67	1.89	-1.99	5.77	-2.83	-9.00	3.34	-1.54	-4.59	7.67	1.89	-1.99	5.77			
CO	-10.41	-30.03	9.21	-22.88	-44.58	-1.18	-28.87	-50.98	-6.76	-8.96	-28.60	10.68	-22.88	-44.58	-1.18	-28.87	-50.98	-6.76	-8.96	-28.60	10.68			
NO ₂	-19.68	-27.83	-11.53	-32.17	-41.81	-22.53	-33.85	-43.38	-24.32	-26.22	-35.41	-17.03	-32.17	-41.81	-22.53	-33.85	-43.38	-24.32	-26.22	-35.41	-17.03			
O ₃	-0.65	-6.51	5.21	-2.00	-9.64	5.64	8.77	1.05	16.49	8.99	-0.59	18.57	-2.00	-9.64	5.64	8.77	1.05	16.49	8.99	-0.59	18.57			

ppb, parts per billion (10⁻⁹); ppm, parts per million (10⁻⁶).

Units and averaging periods:

PM₁₀ (µg/m³): 24 hour avPM_{2.5} (µg/m³): 24 hour av

CO (ppm): 8 hour av

O₃ (ppb): 1 hour maxNO₂ (ppb): 1 hour max

women residing within 5 km of an air monitoring station. Ozone and PM_{2.5} are secondary pollutants and thus should be more homogeneously distributed throughout metropolitan Sydney. The effects of concentrations of PM_{2.5} and O₃ change little when analysing only births to women residing within 5 km of an air monitoring station.

Our findings of an adverse effect of particulates in the second trimester and of NO₂ and CO at various stages of pregnancy on birth weight have been replicated in previous studies. In a meta-analysis of studies examining the impact of particulate exposure on birth weight, Glinianaia and colleagues¹⁶ concluded that in those studies where an effect of particulates in birth outcomes has been shown, there is considerable variability in the stage of pregnancy that the impact occurs.¹⁶ The same can be said of those studies examining the impact of other pollutants on birth weight.

Chen and colleagues⁹ examined the impact of ambient PM₁₀ and other pollutant concentrations on birth weight in Northern Nevada (USA) between 1991 and 1999, and found that exposure to PM₁₀ in the third trimester of pregnancy was negatively associated with birth weight in Nevada.⁹ Gouveia and colleagues⁶ examined the impact of O₃, CO, and PM₁₀ concentrations on birth weight in Sao Paulo, Brazil in 1997, and found that first trimester exposure to CO and PM₁₀ had an adverse affect on birth weight. No association between O₃ concentrations in any trimester and birth weight was observed.

IUGR, defined as birth weight below the 10th centile of recorded birth weight, was associated with exposure to PM_{2.5} and PM₁₀ concentrations in the first trimester of pregnancy in Northern Bohemia.³ In the Czech Republic, IUGR was not associated with exposure to sulphur dioxide (SO₂) and total suspended particulate (TSP) concentrations during pregnancy.² Bobak² also reported that exposure to SO₂, TSP, and oxides of nitrogen (NO_x) during pregnancy had no affect on risk of low birth weight, after adjusting for gestational age (as well as other maternal characteristics). Ha and colleagues⁴ examined the effect of exposure to CO, NO₂, TSP, and SO₂ in the first trimester of pregnancy on low birth weight in Seoul, South Korea. For each inter-quartile increase in pollutant the relative risk of low birth weight was 1.08 for CO, 1.07 for NO₂, 1.06 for SO₂, and 1.04 for TSPs.⁴ No effect of exposure to these pollutants in the third trimester of pregnancy was noted.

Liu and colleagues¹¹ examined the effect of SO₂, NO₂, CO, and O₃ concentrations on IUGR, low birth weight, and preterm birth in Vancouver, Canada. Low birth weight was associated with exposure to SO₂ during the first month of pregnancy (OR 1.1 for a 5 ppb increase). IUGR was associated with exposure to SO₂ (OR 1.1 for a 5 ppb increase), NO₂ (OR 1.1 for a 10 ppb increase), and CO (OR 1.1 for a 1 ppm increase) in the first month of pregnancy.

Last trimester exposure to CO in Los Angeles, USA was reported to result in an increased risk of low birth weight in term births after adjusting for maternal characteristics including commuting habits (OR = 1.22).⁷ Maisonet and colleagues¹⁰ reported an increased risk of term low birth weight due to exposure to CO and SO₂ in all trimesters of pregnancy in six cities in North Eastern United States. No effect of exposure to PM₁₀ was noted.¹⁰ Concentrations of TSP and SO₂ in the last trimester of pregnancy were

In conclusion, we observed that CO and NO₂ concentrations, particularly in the second trimester of pregnancy, have an adverse effect on birth weight. We also observed that PM_{2.5} and PM₁₀ concentrations in the second trimester of pregnancy have a small adverse effect on birth weight. While the number of studies in this area is accumulating, a

Table 6 Key second trimester findings of multivariate linear regression analysis (multiple linear regression coefficients (SE)) using multi-pollutant models and controlling for pollutant exposures in other pregnancy periods for babies born to women residing within 5 km of an air monitoring station; birth weight is expressed in grams

Single pollutant model	Two pollutant models (PM ₁₀ and CO)		Two pollutant models (PM ₁₀ and NO ₂)		Two pollutant models (CO and NO ₂)		Two pollutant models (PM ₁₀ and O ₃)		Four pollutant model		Controlling for exposures in other pregnancy periods							
	95% confidence limits		95% confidence limits		95% confidence limits		95% confidence limits		95% confidence limits		95% confidence limits							
PM ₁₀	-4.28	-5.79	-2.77	-3.72	-6.29	-1.15	-2.65	-4.32	-0.98	-5.47	-7.06	-3.88	-3.27	7.05	0.51	-3.03	-4.85	-1.21
NO ₂	-33.85	-43.38	-24.32			-26.21		-37.46	-14.96	-7.73	-45.75		-9.91	-35.10	1.528	-25.26	-35.63	-14.89
CO	-28.87	-50.98	-6.76	-27.31	-55.30	0.68		-20.17		2.78	-43.12		-25.97	-54.15	2.21	-24.38	-47.33	-1.43
O ₃	8.77	1.05	16.49							18.28	10.03	26.53	-14.81	-33.65	4.03	8.84	1.10	16.58

ppb, parts per billion (10⁻⁹); ppm, parts per million (10⁻⁶)

Units and averaging periods:
PM₁₀ (µg/m³), 24 hour av
CO (ppm), 8 hour av
O₂ (ppb), 1 hour max

consistent pattern of pollutants and exposure times is not emerging. Further research is required to corroborate our findings.

ACKNOWLEDGEMENTS

Environmental Health Branch, NSW Health, and Health Research Foundation, South Western Sydney Area Health Service provided funding for this project. Sharyn Lymer assisted with SAS code. NSW Environment Protection Authority and Australian Bureau of Meteorology provided pollutant and meteorological data. The Australian Research Council and collaborators on the ARC SPIRT project "The Assessment of the impact of air pollution on daily mortality and morbidity in Australian cities using a protocol based on international benchmarking" provided valuable assistance with this project.

Authors' affiliations

T Mannes, NSW Public Health Officer Training Program, New South Wales Health Department, Australia
B Jalaludin, Epidemiology Unit, South Western Sydney Area Health Service, Australia and School of Public Health and Community Medicine, University of New South Wales, Australia
G Morgan, Northern Rivers Department of Rural Health, Sydney University, Australia
D Lincoln, NSW Biostatistical Officer Training Program, New South Wales Health Department, Australia
V Sheppeard, **S Corbett**, Environmental Health Branch, New South Wales Health Department, Australia

Competing interests: none

REFERENCES

- 1 Wang X, Ding H, Ryan L, et al. Association between air pollution and low birth weight: a community based study. *Environ Health Perspect* 1997;**105**:105–15.
- 2 Bobak M. Outdoor air pollution, low birth weight and prematurity. *Environ Health Perspect* 2000;**108**:173–6.
- 3 Dejmek J, Selevan SG, Benes I, et al. Fetal growth and maternal exposure to particulate matter during pregnancy. *Environ Health Perspect* 1999;**107**:475–80.
- 4 Ha EH, Hong YC, Lee BE, et al. Is air pollution a risk factor for low birth weight in Seoul? *Epidemiology* 2001;**12**:643–8.
- 5 Bobak M, Richards M, Wadsworth M. Air pollution and birth weight in Britain in 1946. *Epidemiology* 2001;**16**:358–9.
- 6 Gouveia N, Bremner SA, Noveas HMD. Association between ambient air pollution and birth weight in Sao Paulo, Brazil. *J Epidemiol Community Health* 2004;**58**:11–17.
- 7 Ritz B, Yu F. The effect of ambient carbon monoxide on low birth weight among children born in Southern California between 1989 and 1993. *Environ Health Perspect* 1999;**107**:17–25.
- 8 Rogers JF, Thompson SJ, Addy CL, et al. Association of very low birth weight with exposures to environmental sulfur dioxide and total suspended particulates. *Am J Epidemiol* 2000;**151**:602–13.
- 9 Chen L, Yang W, Jennison BJ, et al. Air pollution and birth weight in Northern Nevada, 1991–1999. *Inhal Toxicol* 2002;**14**:141–57.
- 10 Maisonet M, Bush T, Correa A, et al. Relation between ambient air pollution and low birth weight in the North-Eastern United States. *Environ Health Perspect* 2001;**109**(suppl 3):351–6.
- 11 Liu S, Krewski D, Shi Y, et al. Association between gaseous ambient air pollutants and adverse pregnancy outcomes in Vancouver, Canada. *Environ Health Perspect* 2003;**111**:1773–8.
- 12 Atkinson RW, Anderson HR, Sunyer J, et al. Acute effects of particulate air pollution on respiratory admissions: results from APHEA 2 project. *Am J Respir Crit Care Med* 2001;**164**(10 pt 1):1860–6.
- 13 Katsouyanni K, Touloumi G, Samoli E, et al. Confounding and effect modification in the short-term effects of ambient particles on total mortality: results from 29 European cities within the APHEA2 project. *Epidemiology* 2001;**12**:521–31.
- 14 Le Tertre A, Medina S, Samoli E, et al. Short-term effects of particulate air pollution on cardiovascular diseases in eight European cities. *J Epidemiol Community Health* 2002;**56**:773–9.
- 15 Roberts CL, Lancaster PAL. Australian national birth weight percentiles by gestational age. *Med J Aust* 1999;**170**:114–18.
- 16 Glinianaia SV, Rankin J, Bell R, et al. Particulate air pollution and fetal health: a systematic review of the epidemiologic evidence. *Epidemiology* 2004;**15**:36–45.
- 17 Jedrychowski W, Bendkowska I, Flak E, et al. Estimated risk for altered fetal growth resulting from exposure to fine particles during pregnancy: an epidemiologic prospective cohort study in Poland. *Environ Health Perspect* 2004;**112**:1398–402.
- 18 Peters A, Doering A, Wichmann HE, et al. Increased plasma viscosity during an air pollution episode: a link to mortality. *Lancet* 1997;**349**:1582–7.
- 19 Bouthillier L, Vincent R, Goegan P, et al. Acute effects of inhaled urban particles and ozone: lung morphology, macrophage activity, and plasma endothelin-1. *Am J Pathol* 1998;**153**:1873–84.
- 20 Knottnerus JA, Delgado LR, Knipschild PG, et al. Haematologic parameters and pregnancy outcome: a prospective cohort study in the third trimester. *J Clin Epidemiol* 1990;**43**:461–6.
- 21 Lin CC, Santolaya-Forgas J. Current concepts of fetal growth restriction: Part 1: Causes, classification and pathophysiology. *Obstet Gynaecol* 1998;**92**:1044–55.
- 22 New South Wales Environment Protection Authority. *State of the environment 2000*. Sydney: NSW Environment Protection Authority, December 2000. Available: <http://www.epa.nsw.gov.au/index.htm> [accessed July, 2003].