111

LYMPHOMA RISK AMONG ANIMAL BREEDERS

Ileana D'Andrea,¹ Maria Grazia Ennas,¹ Giannina Satta,¹ Mariagrazia Zucca,¹ Tinucia Nonne,¹ Andrea 't Mannetje,² Nikolaus Becker,³ Silvia De Sanjosé,⁴ Lenka Foretova,⁵ Anthony Staines,⁶ Marc Maynadié,⁷ Alexandra Nieters,³ Paul Brennan,⁸ Paolo Boffetta,⁹ Michele Meloni,¹ Pierluigi Cocco¹ ¹University of Cagliari, Cagliari, Italy; ²Massey University, Wellington, New Zealand; ³German Cancer Research Center, Heidelberg, Germany; ⁴Catalan Institute of Oncology, Barcelona, Spain; ⁵Department of Cancer Epidemiology and Genetics, Brno, Czech Republic; ⁶Dublin City University, Dublin, Ireland; ⁷Dijon University Hospital, Dijon, France; ⁸IARC, Lyon, France; ⁹Mount Sinai School of Medicine, New York, USA

10.1136/oemed-2011-100382.111

Objectives Occupational contact with breeding animals might be implicated in lymphoma aetiology.

Methods In 1998–2003, 2337 incident lymphoma cases and 2434 controls participated in the EPILYMPH case-control study in six European countries. A detailed occupational history was collected in cases and controls, including species of breeding animals, their approximate number, and circumstances of contact. We conducted a preliminary analysis on ever exposed to contact with breeding animals, and we stratified the analysis by age at first exposure, whether before or after 12. The OR and its 95% CI was calculated with unconditional logistic

regression for all lymphomas, and its major subtypes, adjusting by age, gender, and education.

Results Lymphoma risk (all subtypes combined) did not increase among those exposed to contact with breeding animals (OR = 1.0, 95% CI 0.8 to 1.2). Risk of DLBCL was significantly lower among subjects employed in poultry farms (OR = 0.6, 95% CI 0.4 to 1.0). This inverse association was observed among subjects who started exposure before or at age 12 (OR = 0.5, 95% CI 0.2 to 1.1), but not later.

Conclusions Early occupational contact with poultry might be associated with a decrease in risk of specific lymphoma subtypes.