Conclusions In the present study, a high prevalence of UI symptoms was found in soccer athletes, with moderate reliability between the pad test and the ICIQ-SF; mild UI was found in both groups. The athletes had the highest scores on the General Health, Emotions and Sleep/Energy domains, which are considered to have the highest impact on quality of life.

0017 LIFE QUALITY IN WORKERS WITH RELATED MUSCULOSKELETAL DISORDERS

¹Andressa Soares, ¹Fernanda Caetano, ¹Joyce Braga, ²Antonio Fernandes, ^{1,2}Eduardo Filoni. ¹Cruzeiro Do Sul University, São Paulo, São Paulo, Brazil; ²University of Mogi Das Cruzes, Mogi Das Cruzes, São Paulo, Brazil

10.1136/oemed-2014-102362.185

Objectives Analyse the quality of life of workers of administrative department who carry out repetitive functions in a medical certification central.

Method This study was carried out in a transversal analytical approaching. The sample was constituted by 62 workers of the administrative sector of a medical certification central in the state of São Paulo. As an instrument of gathering the data the used questionnaires were: Anamnesis Questionnaire with personal and professional data, Nordic Questionnaire, Job stress scale, SF-36 Quality of Life Questionnaire. The characterisation of the sample was obtained through descriptive statistics, where test t and correlation of Pearson test were applied.

Results a According to the results of the analysis of the Nordic questionnaire, 33, 87% mentioned feeling pain in the wrists and hands. In the correlation made among the data of SF-36 and Nordic questionnaire, it was showed that they presented complaints about the Nordic, minor was the score in the variables of SF-36: limitation by physical aspects, pain, vitality, emotional aspects and mental health.

Conclusions It is concluded that the most affected segments were wrists and hands, with 33, 87%, followed by anklebone and feet with 16, 13%.

0020 COMPARATIVE STUDY ABOUT QUALITY OF LIFE BETWEEN WOMEN PRACTICING DANCE AND SEDENTARY WOMEN

¹Marijalma Araújo, ¹Eduardo Filoni, ¹Antonio Fernandes, ²Aline Pansani. ¹University of Mogi Das Cruzes, Mogi Das Cruzes, São Paulo, Brazil; ²Federal University of São Paulo, São Paulo, São Paulo, Brazil

10.1136/oemed-2014-102362.186

Objectives This study aimed to compare Quality of life (QOL) of women who practice Jazz compared to sedentary women, in the Sports Centre-414 SESI in Sao Paulo - SP

Method We carried out a cross-sectional study, with women between 50 and 85 years old, divided into two groups: Activity Group (GA, n = 14) and Sedentary Group (GS, n = 14). QOL was analysed by the SF-36.

Results The GA had a significant improve in 6 domains of SF-36 compared to GS (p < 0.05).

Conclusions The results of this study suggest an improvement in the QOL of practitioners of jazz, especially in the physical and emotional aspects, compared to sedentary women.

0024 EFFECTS OF CHRONIC LOW-DOSE EXPOSURE TO IONISING RADIATION ON PHYSICIAN MICRO-VASCULAR STRUCTURE REVEALED BY NAIL FOLD CAPILLAROSCOPY

¹Pascal Wild, ²Christine Gauron, ³Christian Derock, ³Karine Champion, ³Pascal Cohen, ⁴Caroline Menez, ⁵Anne-Sophie Tellart, ⁶Hélène Thiel, ¹Michel Grzebyk, ⁷Laurent Donadille, ⁸Raffaele Pennarola, ⁹Dominique Choudat. ¹INRS, Vandoeuvre, France; ²INRS, Paris, France; ³APHP, Paris, France; ⁴University Hospital, Grenoble, France; ⁵University Hospital, Lille, France; ⁶University Hospital, Clermont-Ferrand, France; ⁷IRSN, Fontenay Aux Roses, France; ⁸Naples University, Naples, Italy; ⁹University Paris Descartes, Paris, France

10.1136/oemed-2014-102362.187

Objectives The aim of this study was to assess the long term pre-clinical effects of low-dose radiation on the micro-vascular structure among interventional physicians whose hands are exposed to ionising radiation in their daily practice.

Method The study, approved by the ethics committee, included 186 radiation-exposed (surgeons, cardiologists and radiologists) and 35 unexposed physicians. The study participants filled in a questionnaire describing their present and past daily practice from which the present and cumulative radiation exposures were estimated. Their dermal microcirculation was assessed by capillaroscopy of the nailfold of 8 fingers -thumbs excluded- of each study participant. Two quantitative scores characterising respectively extravasation and morphological abnormalities were obtained based on post-hoc codings of seven semi-quantitative indices by five capillaroscopists. These assessments were randomised and were blind to the exposure. The effect of the radiation exposure on both features was modelled using multilevel proportional odds regression adjusted for potential confounders.

Results The exposure intensity of each act was greatest among surgeons but with a lesser number of acts. The mean estimated cumulative exposure was largest among radiologist and cardiologists.

No effect of the exposure could be detected on the extravasation score. The morphological anomaly score increased significantly with duration of exposure and cumulative exposure indices among surgeons and interventional radiologists contrary to cardiologists among which no effect could be detected.

Conclusions Capillaroscopy showed significant effects of chronic low-dose exposure to ionising radiation on micro-vascular structure among exposed physicians but its interest for individual monitoring has not been demonstrated.

0025 ASSESSMENT AND ESTIMATES OF EXPOSURE TO SYNTHETIC AMORPHOUS SILICA AT FIVE GERMAN PRODUCTION PLANTS

^{1,2}Peter Morfeld, ³Dirk Taeger, ⁴Heike Mitura, ⁵Axel Bosch, ⁶Adrian Nordone, ⁷Reinhard Vormberg, ⁸Robert McCunney, ³Rolf Merget. ¹Institute for Occupational Epidemiology and Risk Assessment (IERA) of Evonik Industries, Essen, Germany; ²Institute and Policlinic for Occupational Medicine, Environmental Medicine and Prevention Research, Köln, Germany; ³Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Bochum, Germany; ⁴AQura GmbH, Hanau, Germany; ⁵Wacker Chemie AG, Burghausen, Germany; ⁶Cabot Corporation, Boston, USA; ⁷Evonik Industries AG, Hanau, Germany; ⁸Massachusetts Institute of Technology (MIT), Boston, USA

10.1136/oemed-2014-102362.188

Objectives Synthetic amorphous silicas (SAS) are nanostructured polymorphs of silicon dioxide. We compared two different exposure assessments.

Method This study estimated cumulative exposure to inhalable SAS dust in 484 male workers from five German SAS producing plants. Two procedures (P1, P2) were applied. P1 was based on