OPEN ACCESS

ORIGINAL ARTICLE

Short-term metal particulate exposures decrease cardiac acceleration and deceleration capacities in welders: a repeated-measures panel study

Peter E Umukoro,¹ Jennifer M Cavallari,^{1,2} Shona C Fang,¹ Chensheng Lu,¹ Xihong Lin,³ Murray A Mittleman,^{4,5} David C Christiani^{1,4,6}

¹Department of Environmental Health, Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA

²Department of Community Medicine and Health Care, University of Connecticut Health Center, Farmington, Connecticut, USA

³Department of Biostatistics, Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA

⁴Department of Epidemiology, Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA

⁵Cardiovascular Epidemiology Research Unit, Beth Israel Deaconess Medical Center/Harvard Medical School, Boston, Massachusetts, USA

⁶Pulmonary and Critical Care Unit, Massachusetts General Hospital/Harvard Medical School, Boston, Massachusetts, USA

Correspondence to

Dr David C Christiani, Harvard T.H. Chan School of Public Health, Department of Environmental Health, Boston, MA 02115, USA; dchris@hsph.harvard.edu

Received 30 April 2015

Revised 24 September 2015

Accepted 20 October 2015

Published Online First

7 December 2015

Open Access
Scan to access more
free content

CrossMark

To cite: Umukoro PE, Cavallari JM, Fang SC, et al. *Occup Environ Med* 2016;**73**:91–96.

ABSTRACT

Objective Acceleration (AC) and deceleration (DC) capacities measure heart rate variability during speeding up and slowing down of the heart, respectively. We investigated associations between AC and DC with occupational short-term metal PM_{2.5} exposures.

Methods A panel of 48 male welders had particulate matter less than 2.5 microns in diameter (PM_{2.5}) exposure measurements over 4–6 h repeated over 5 sampling periods between January 2010 and June 2012. We simultaneously obtained continuous recordings of digital ECG using a Holter monitor. We analysed ECG data in the time domain to obtain hourly AC and DC. Linear mixed models were used to assess the associations between hourly PM_{2.5} exposure and each of hourly AC and DC, controlling for age, smoking status, active smoking, exposure to secondhand smoke, season/time of day when ECG reading was obtained and baseline AC or DC. We also ran lagged exposure response models for each successive hour up to 3 h after onset of exposure.

Results Mean (SD) shift PM_{2.5} exposure during welding was 0.47 (0.43) mg/m³. Significant exposure–response associations were found for AC and DC with increased PM_{2.5} exposure. In our adjusted models without any lag between exposure and response, a 1 mg/m³ increase of PM_{2.5} was associated with a decrease of 1.46 (95% CI 1.00 to 1.92) ms in AC and a decrease of 1.00 (95% CI 0.53 to 1.46) ms in DC. The effect of PM_{2.5} on AC and DC was maximal immediately postexposure and lasted 1 h following exposure.

Conclusions There are short-term effects of metal particulates on AC and DC.

INTRODUCTION

Short-term and long-term particulate exposures have been shown to have adverse effect on cardiovascular outcomes.^{1–4} One of the mechanisms involved is through affectation of the autonomic nervous system, which has been measured traditionally using heart rate variability (HRV).^{5–7} Air pollution studies have consistently shown a decrease in HRV with exposure to PM_{2.5}.^{5–9} A decrease in HRV has also been linked to an increase in adverse cardiovascular outcomes.^{10–13}

Whereas there is ubiquitous exposure to particulates in ambient air, occupational exposure to particulates is usually greater. For example, welders have been shown to have been exposed to about 24 times

What this paper adds

- ▶ Heart rate variability (HRV) has been the widely used outcome in cardiac autonomic research but has been critiqued for its potential of misclassification with changing heart rates.
- ▶ Short-term metal PM_{2.5} exposure decreases HRV, but the specific effects on accelerations and decelerations are unknown.
- ▶ This paper shows that there are decreases in both cardiac accelerations and decelerations.
- ▶ The decreases in acceleration and deceleration capacities persist up to 1 h even after exposure ceases.
- ▶ Targeted interventions should aim at reducing particulate exposure and mitigating effects on both cardiac acceleration and deceleration capacities.

the level of ambient PM_{2.5} exposure levels.^{14–15} Furthermore, an exposure-related decrease in HRV has been demonstrated among these welders.¹⁶

Although previous research suggests that there is a significant exposure–response relationship with increasing exposures to metal PM_{2.5} producing increasing cardiac autonomic dysfunction (reduced HRV), this was evaluated using HRV as the index of autonomic dysfunction, which has limitations. HRV fails to account for the heart rate, and is therefore prone to misclassification.¹⁷ Current mechanistic research using HRV cannot clarify if a decrease in HRV means a decrease in accelerations and/or decrease in decelerations of the heart rate.

Acceleration capacity (AC) is a measure of the responsiveness of the heart to speed up when stimulated, and it is known to be under both autonomic (sympathetic) and non-autonomic control.^{11–12} Deceleration capacity (DC), on the other hand, describes the behaviour of the heart when the heart rate is slowing, and it reflects a measure of parasympathetic modulation of the heart.¹⁸ While there have been no population-level studies yet to measure the normal range levels of AC and DC in the healthy general population, most studies of patients with postmyocardial infarction have reported baseline levels of magnitude 7.0–8.0 ms for both AC and DC in their respective scales.^{17–19–21}

In 2006, Bauer *et al*¹⁷ described the phase-rectified signal averaging (PRSA) method for calculating the cardiac AC and DC, which are measures of the responsiveness of the heart, like HRV.²¹ These have the advantage over HRV of parsing the Holter data into accelerations and decelerations while also accounting for the heart rate, and have been demonstrated to be more predictive of morbidity and mortality among patients with postmyocardial infarction than traditional HRV.¹⁷ Therefore, using sensitive indices—AC and DC—this study aims to investigate the potential for cardiac autonomic dysfunction from metal PM_{2.5} exposure. We hypothesised that metal-rich PM_{2.5} exposure would decrease both AC and DC.

METHODS

Participant recruitment

We recruited a convenience sample of 72 male boilermakers based on outreach to the union membership with an overall participation rate of 93% during five sampling periods between January 2010 and June 2012 from the boilermaker union in Quincy, Massachusetts. These boilermakers were part of an ongoing 'Harvard Boilermakers' Cohort' initiated in 1999 to study the cardiopulmonary effects of particulates.²² The 'Harvard Boilermakers' Cohort' was a coalition of different smaller panel studies that were conducted at different sampling periods to answer specific research questions including the effect of secondhand smoke and metal particulates on cardiopulmonary effects. Participants were mostly monitored on consecutive non-welding and welding days during each sampling period. Although we had recruited 72 boilermakers, we restricted our study to 52 participants who were only monitored on welding days. We were only able to record simultaneous PM_{2.5} and digital ECG on non-welding days for 20 participants, poor-quality ECG data in 2 participants, and failed ECG data retrieval in 2 other participants. We analysed data from 48 boilermakers (our final sample) on only welding days for our study. They constituted 67% of the participants in the existing cohort whom we were able to obtain PM_{2.5} exposure from during welding shift, as well as record continuous digital ECG recording during welding shifts recruited within sampling periods between 2010 and 2012. We conducted our study during winter or summer when 75% (36) of the study participants had not actively welded 2 weeks prior to our data collection. The other 25% (12) had welded between 3 and 10 days before our sampling (weld) day. The Institutional Review Board at the Harvard T. H. Chan School of Public Health approved the study protocol, and informed consent was obtained from each study participant.

Data collection

We collected continuous PM_{2.5} exposure and continuous ECG data of study participants at a union welding school. The welding school was designed for training apprentices and had booths where boilermakers practised welding (mainly stick-metal and gas-metal arc welding), cutting and grinding. The school was a ventilation-controlled and temperature-controlled environment, and there were no other sources of PM_{2.5} except from the welding fumes, tobacco smoke and ambient indoor levels. Welding fumes contain fine particles and vaporised metals from the metal alloy being welded and the electrodes used.^{5 23} All boilermakers used metal arc welding (stick or gas) using base metals of mild steel with electrodes composed mainly of iron with manganese.¹⁴

We also collected medical history and medication use information, demographics, lifestyle information, including smoking,

typical diet and occupational history, using self-administered questionnaire. Participants were asked to report any of the following heart and blood vessel problems, diagnosed by a physician: hypertension, use of blood pressure medications such as β -blockers or ACE inhibitors, congestive heart failure, myocardial infarction, angina, arrhythmia, heart/chest surgery, or otherwise non-classified heart problems—not diagnosed by a physician.

PM_{2.5} assessment

We measured PM_{2.5} concentrations during welding shifts of study participants at the union welding school using personal DustTrak Aerosol Monitor (TSI, Inc, St. Paul, Minnesota, USA). The DustTrak monitor was strapped to the participant's shoulder close to their breathing zone. DustTrak has a PM_{2.5} inlet impactor to measure continuously and record at 1 min intervals average concentrations of fine particulates during the welding shifts. The continuous DustTrak readings of PM_{2.5} had been validated compared with gravimetric methods in welders.²⁴ We calculated mean hourly concentrations of PM_{2.5} exposure during each work shift for each participant.

ECG recording and processing

Study participants wore a standard five lead ECG Holter monitor after a 30 min rest period in the morning on arrival at the union hall. The rest period was allowed for so that we could record their unbiased resting ECG free from acute changes resulting from commuting to the study site. To ensure that the leads of the ECG were well secured and remained secured on the chest of participants, we shaved their skin if necessary, cleaned with an alcohol wipe after slightly abrading the skin, and research staff checked them intermittently.⁵ The participants had this monitor worn throughout the welding shift, and up to 3 h following welding fume exposure. The digital recordings were then downloaded and sent to the Cardiovascular Epidemiology Research Unit (CVERU) of Beth Israel Deaconess Medical Center (Boston, Massachusetts, USA) for processing and analysis. Holter recordings were uploaded into the GE MARS ECG analysis system, which automatically scans recordings for areas of noise and groups heartbeats as normal or arrhythmic. Trained technicians blinded to the exposure status of the participant from whom the ECG reading was obtained verified the automated scans as correct or changed them to the appropriate designation. The data were then exported for analysis using the Physionet toolkit.²⁵ To remove artefacts from the data, they used only beats with a distance between two consecutive R waves on the electrocardiogram (RR) interval within 5% difference of adjacent beats. They used an automated process described by Bauer *et al*¹⁷ to create 5 min segments with anchors for the PRSA method of computing the AC and DC. In brief, to compute the DC, this involves identifying heartbeat intervals longer than the preceding interval as anchors (for AC, beats shorter than preceding beats were anchors). Overlapping segments of interval data were then automatically generated from the ECG such that all segments are aligned at the anchors in the centre and averaged. The PRSA method then quantifies the signals within aligned segments using the Haar wavelet analysis with a scale of 2 by a computer processing of the ECG with visual and digital outputs. Thus, AC and DC were calculated separately as a quarter of the difference between two sums, that is, the sum of the averaged anchor points RR intervals (X_0) with the succeeding RR intervals (X_1) and the sum of the two averaged RR intervals preceding anchor points (X_{-2} , X_{-1}).²¹

AC and DC

Using the digital ECG data in the time domain, we computed the average AC and average DC for each simultaneous hour of PM_{2.5} exposure and monitoring by taking the mean of the 12 adjacent 5 min segments of the ECG within each hour of the day using the automated output.

Data analysis

We calculated summary measures of potential covariates, and percentiles of our exposure and outcome to further understand their distribution. Potential covariates that we considered include: age, race, smoking status (smoker/non-smoker), actively smoking during work (yes/no) and secondhand smoke exposure (exposed/not exposed), time of day when ECG was obtained, season of study, previous weld exposure (last weld day), chronic effects of welding (years of boilermaker), actively welding (yes/no), presence of heart problems, and baseline cardiac autonomic function (baseline AC or DC for AC and DC models, respectively). Secondhand smoke exposure was defined as being exposed to smoke from other nearby workers smoking during our sampling. We then explored the inter-relationships between them by using Spearman's correlations for continuous variables and t tests for binary variables. In order not to have missed any inter-relationships between potential covariates with our exposure/outcome, we used $\alpha=0.10$ level for these correlations and t tests between covariates and outcome/exposure.

For our model without any lag between exposure and response, we used linear mixed models to assess the associations between hourly PM_{2.5} as a continuous measure and simultaneously measured hourly AC, and a separate model for PM_{2.5} and DC. Furthermore, we used a backward method of model selection using a p value of 0.2 each for staying in the model to select our final model. We compared models using the log likelihood ratio test. We considered controlling for age, smoking status, last weld day (unmeasured acute weld exposure), number of years as a boilermaker (chronic effects), baseline AC or DC (for AC or DC models respectively), and time of day and season when ECG reading was obtained. Our final model included age, baseline AC or DC, smoking status, active smoking, secondhand smoke exposure, time of day and season of ECG. For the lagged models, we used linear mixed models to assess the associations between hourly PM_{2.5} and lagged hourly AC or DC at 1, 2 and 3 h lags. A lag was defined as the time period between exposure (metal-rich PM_{2.5}) and its response (AC or DC). Therefore, a '1 hour' lag model was evaluating the effect of PM_{2.5} at a time point on AC or DC level '1 hour' later. Statistical significance was assessed at $\alpha=0.05$ level in two-sided tests for our final model. All analyses were performed using PROC MIXED in SAS V9.4 (Cary, North Carolina, USA).

RESULTS

We successfully collected 892 person-hours of weld day PM_{2.5} and 1392 person-hours of weld day ECG from 48 participants during a median of three work shifts per participant of 4–6 h duration, all males with a mean age of 40 years and had been boilermakers for a median of 4 years (range 0.25–21 years). The study population included 19 (40%) smokers (table 1).

Five of these 48 participants reported heart problems and possible heart-related problems. Of these five, one reported arrhythmia, one reported sinus (sinoatrial node) problems, one reported angina in the past year, one reported mitral valve prolapse with murmur and one reported hypertension for which he was taking a β -blocker. No other participant reported the use of

Table 1 Demographics and characteristics for the 48 study participants

Individual characteristics	N	Per cent
Male	48	100
Caucasian	42	88
Smoking status (current smoker)*	19	40
Other characteristics		
Time of ECG (AM)†	18	38
Season of ECG (Winter)‡	33	69
Heart problems§	5	10
	Mean	SD
Age at start of study 2010 (years)	40	12
Number of years as a boilermaker (years)	9	9
Last weld day before study (days)	44	51

*Current smokers versus non-smokers and previous smokers.

†Morning (AM) versus afternoon (PM).

‡Winter versus summer.

§Heart problems include reported previous history of arrhythmia, cardiac sinus problems or palpitations.

β -blockers or ACE inhibitor drug use. Less than half (38%) of the baseline ECG were taken in the morning, and each participant had PM_{2.5} measurements taken in 2–5 typical work shifts of 4–6 h.

The mean number of years as a boilermaker was 9 years (range 1–35 years). The mean PM_{2.5} during the work shift for participants measured was 0.47 mg/m³ (range 0.01–1.40 mg/m³). The mean (range) baseline AC was –7.1 ms (–2.1 to –22.6) on the negative scale (table 2). The mean (range) baseline DC was 8.8 ms (3.3–22.4).

Spearman's correlations coefficients (and p value) for correlation between potential covariates and PM_{2.5} or baseline measures of AC and DC were mostly statistically non-significant. However, season of the year and active smoking were correlated with AC and/or DC, but not with PM_{2.5}. Among the other covariates considered, age was neither correlated with PM_{2.5} exposure nor with the baseline AC or DC. The number of years as a boilermaker and last weld day were only moderately correlated with PM_{2.5} exposure but not with baseline AC or DC. There were no differences in categories of other covariates in terms of measures of shift PM_{2.5}, baseline AC and DC.

Table 2 Baseline and hourly levels of metal PM_{2.5} (mg/m³), acceleration (AC) and deceleration (DC) capacities of the heart (ms)

	Baseline Mean (SD)	Hourly average Mean (SD)*	Hourly average minus baseline Mean	p Value
Exposure				
PM _{2.5} (mg/m ³)†	0.04 (0.3)	0.50 (0.4)	0.46	<0.01
Outcome				
AC (ms)‡	–7.1 (3.8)	–6.3 (3.1)	0.8	0.25
DC (ms)	8.8 (3.4)	7.8 (3.8)	–1.0	0.17

Bold values indicate significant associations (p<0.05).

*This represents the mean of the hourly averages (hours 1, 2 and 3) from the onset of welding.

†PM_{2.5} ambient levels before shift and average hourly levels during work shift are reported as baseline and hourly average, respectively.

‡AC is measured on a negative scale. N=48 participants.

The linear mixed models analyses for our model without any lag revealed that PM_{2.5} levels were associated with a decrease in AC and DC after adjusting for age, smoking status, active smoking, secondhand smoke exposure and baseline AC or DC (tables 3 and 4).

When we introduced lags by the hour between PM_{2.5} exposure and AC or DC, there were associations between metal PM_{2.5} exposure and lagged responses in AC and DC at 1 h postexposure. These were consistent with or without adjustment for PM_{2.5} exposure in the previous hour(s). When we excluded the five participants with heart problems, the results were consistent. A sensitivity analysis that excluded the 12 participants who had welded within 2 weeks before our study showed qualitatively similar results (table 5).

DISCUSSION

The goal of this study was to investigate the associations between AC and DC with acute metal-rich PM_{2.5} exposures. Consistent with our hypothesis, metallic PM_{2.5} exposure was associated with AC and DC. We found significant exposure-response relationships between increasing short-term metallic PM_{2.5} exposure producing decreasing AC and DC. These associations were qualitatively consistent with or without adjustment for PM_{2.5} exposure in the previous hour(s). There may therefore be a reduction in the capacity of the heart to accelerate and decelerate over time with acute insults from these exposures. These data imply that there may be changes in parasympathetic control, sympathetic modulation and/or non-autonomic control of the heart with acute exposure to particulates. Few studies have documented associations between particulate exposures and HRV in the short term.^{2 5 7 9 26} Yet, fewer studies have observed associations between short-term fine particulate exposure and DC.¹⁸ This study is the first to our knowledge to demonstrate declines in AC following acute metal-rich particulate exposures among healthy welders.

We cannot directly compare our study results with those of prior studies due to the differences in exposure characteristics and study population. We did find declines in DC slightly greater than declines reported following short-term PM_{2.5} exposures among post-myocardial infarction patients.¹⁸ In addition, we observed declines in AC among welders who are not acutely

Table 4 Main effect of PM_{2.5} (mg/m³) on acceleration capacity (AC; ms) and deceleration capacity (DC; ms) with adjustment for hourly pre-exposures from the linear mixed effect response-lagged models with 48 participants

Parameter	AC* (95% CI)	DC (95% CI)
Models		
No lag	1.46 (1.00 to 1.92)	−1.00 (−0.53 to −1.46)
1 h lag†	0.81 (0.30 to 1.32)	−0.70 (−0.14 to −1.26)
2 h lag†	1.00 (0.13 to 1.87)	−0.45 (0.52 to −1.42)
3 h lag†	0.41 (−1.65 to 2.46)	−0.12 (2.48 to −2.71)

Bold values indicate significant associations (p<0.05). Models are adjusted for age, smoking status, actively smoking, secondhand smoke exposure, time of day and season of ECG reading, and baseline AC or DC.

*AC is measured on a negative scale; therefore, a 1 mg/m³ increase in PM_{2.5} results in a decrease in AC by 1.46 ms (on a negative scale) for the association without any lag.

†Lagged models adjust for exposure time points before the lags. For example, a 1 h lag model also adjusts for PM_{2.5} measurements in the preceding hour before the exposure time point of interest.

ischaemic which has never been reported. This result highlights the possibility of differences in effect among our study population (active healthy welders) compared with other studies (patients with postmyocardial infarction) that were mostly conducted in clinical settings. Furthermore, the different constituents of the particulate exposures may have varying effects on the electrical activity of the heart. While organic and elemental carbons have been implicated as suspects, metals have also been shown to play a role.^{6 14 18 27} Welders are exposed to metal-rich fumes, and this may be responsible for the effects on AC we found in this study.

Only five of the study participants reported known or possible heart problems. There was no difference in quality of results in the sensitivity analyses that excluded these five participants. This suggests that the effects of particulates in chronic ischaemic heart conditions and healthy persons are qualitatively similar.

We found significant effects with both the AC and DC, as well as lagged responses between PM_{2.5} and DC. We would

Table 3 Main effect of PM_{2.5} (mg/m³) on acceleration capacity (AC; ms) and deceleration capacity (DC; ms) without adjustment for hourly pre-exposures from the linear mixed effect response-lagged models with 48 participants

Parameter	AC* (95% CI)	DC (95% CI)
Models		
No lag	1.46 (1.00 to 1.92)	−1.00 (−0.53 to −1.46)
1 h lag†	0.73 (0.26 to 1.20)	−0.40 (0.08 to −0.87)
2 h lag†	−0.29 (−0.83 to 0.24)	0.65 (1.22 to 0.07)
3 h lag†	−0.34 (−0.92 to 0.25)	0.52 (1.15 to −0.12)

Bold values indicate significant associations (p<0.05). Models are adjusted for age, smoker status, actively smoking, secondhand smoke exposure, time of day and season of ECG reading, and baseline AC or DC.

*AC is measured on a negative scale; therefore, a 1 mg/m³ increase in PM_{2.5} results in a decrease in AC by 1.46 ms (on a negative scale) for the association without any lag.

†Lagged models adjust for exposure time points before the lags. For example, a 1 hour lag model also adjusts for PM_{2.5} measurements in the preceding hour before the exposure time point of interest.

Table 5 Main effect of PM_{2.5} (mg/m³) on acceleration capacity (AC; ms) and deceleration capacity (DC; ms) with adjustment for hourly pre-exposures from the linear mixed effect response-lagged models with 36 participants who had not welded 2 weeks before the study

Parameter	AC* (95% CI)	DC (95% CI)
Models		
No lag	1.54 (0.96 to 2.32)	−1.18 (−0.60 to −1.79)
1 h lag†	0.97 (0.13 to 1.81)	−0.89 (−0.09 to −1.70)
2 h lag†	0.66 (−0.94 to 1.87)	−0.37 (1.09 to −1.83)
3 h lag†	0.29 (−2.58 to 2.60)	−0.18 (2.57 to −2.15)

Bold values indicate significant associations (p<0.05). Models are adjusted for age, smoking status, actively smoking, secondhand smoke exposure, time of day and season of ECG reading, and baseline AC or DC.

*AC is measured on a negative scale; therefore, a 1 mg/m³ increase in PM_{2.5} results in a decrease in AC by 1.54 ms (on a negative scale) for the association without any lag.

†Lagged models adjust for exposure time points before the lags. For example, a 1 hour lag model also adjusts for PM_{2.5} measurements in the preceding hour before the exposure time point of interest.

therefore hypothesise that during acute exposure to particulates, there is a blunting of the autonomic response of the heart to the activation of both parasympathetic and sympathetic nervous system influences to regulate the response of the heart, as well as the interplay of non-autonomic control (cytokines and inflammatory mediators) which may be playing a major role in the lagged responses of AC and DC to exposure of metal particulates. We found the effects on AC and DC lasted up to 1–2 h postexposure. We doubt the sustenance of autonomic effects up to this time, and we would hypothesise that non-autonomic influences may begin to play a role at this time. If this hypothesis is correct, then our results are not completely consistent with previous knowledge that AC is influenced by both autonomic control and inflammatory stimuli, and that DC which is mainly under autonomic control,¹⁷ since we found that sustained effects of PM_{2.5} exposure affect both AC and DC. This discrepancy may be explained by AC and DC being antagonistically opposed to each other. There may be greater sensitivity on AC by acute insults as the effect on AC remains until 2 h following the insult.

DC has been shown to be predictor of mortality following acute myocardial infarction, even among patients with preserved AC, as there is a rapid decline in the parasympathetic innervation of the heart following ischaemia.^{17–28} We did observe declines in the DC among these healthy welders as only one of them had had an ischaemic event within the last year of the study. However, the clinical usefulness of a decline in the DC of the heart of healthy persons is unknown. Even more so, the significance of declines in AC in both ischaemic/non-ischaemic hearts is uncertain. This may be due to the multifactorial influences on the AC.

Short-term metal particulates have effect on the AC and DC lasting 1 h postexposure. This is similar to the early phase of the multiphasic response of HRV to particulates which showed a sustained effect on HRV up to 3 h.⁸ However, our study results did not show a sustained effect on AC and DC up to 3 h postexposure. This may be as a result of misclassification of HRV with respect to accelerations and decelerations. Additionally, the declines in HRV (15 ms per 1 mg/m³ PM_{2.5}) observed in similar study population were 10-fold greater than the declines observed in AC and DC.¹⁶ Thus, HRV may be differentially misclassifying effect of PM_{2.5} exposure on cardiac automaticity with respect to both accelerations and decelerations.

There was no difference between mean hourly AC and DC compared with baseline. This may be the result of averaging over few hours after PM_{2.5} exposure ceases to have effect on AC and DC. This result suggests that the heart may be adapting its response to the continuous exposure of fine particulate. This is further supported by the intermediate decline in AC at the '1 h lag' compared with the 'no lag' and '2 h lag' indicating some attempt of the heart at recovery (table 4). Excluding study participants who had welded within 2 weeks before our study did not significantly alter the results of our study. This suggests that there may not be a significant cardiac sensitisation on re-exposure to metallic PM_{2.5} following a washout period.

Based on our study results, reducing exposure would be a major goal of health promotion for these workers. The use of personal protective equipment such as a respirator has been shown to reduce exposures to particulates and improve HRV.²⁹ Our study has its strengths and limitations. Importantly, we captured ECG tracings digitally that were parsed into phases of accelerations and decelerations accounting for differences in heart rate. Our results are therefore not confounded by heart rate. We also adjusted for confounding by potential

confounders. We used repeated continuous shift PM_{2.5} measured using DustTrak Aerosol Monitor. We were also able to account for previous welding exposure by restricting most of our study population to those who had not welded 2 weeks prior to our study, and considered adjusting for prior uncaptured PM_{2.5} exposure using 'last weld day' in our models. PM_{2.5} exposure measurements were obtained in comparable work settings at the union hall⁵ and may reflect what their typical exposures would have been if they were followed up at their work places, which would be challenging to do. There is a potential for selection bias in our study as we were only able to retrieve weld day data for 67% of our participants. However, there were no differences in baseline characteristics of the welders whom we were not able to retrieve their records and those in our study. We also followed up study participants for only 3 h following onset of exposure. The specific component of PM_{2.5} exposure causing the declines in AC and DC would require further study. Therefore, the results of this study will only be generalisable to similar occupational settings with rich metallic PM_{2.5} exposure.

In conclusion, metal-rich occupational PM_{2.5} exposure is associated with a reduction in AC and DC lasting up to 1 h after exposure has ceased. This suggests that there may be more pathways other than a direct autonomic effect of particulates on AC and DC that sustain the acute cardiac response following metallic PM_{2.5} exposure up to 1 h following exposure.^{15–30} An investigation of the non-autonomic control of cardiac AC and DC is needed. The specific component of PM_{2.5} exposure causing the declines in AC and DC would require further study. In addition, targeted interventions should aim at reducing particulate exposure and mitigating effects on both cardiac AC and DC.

Acknowledgements This research was supported by the National Institute of Environmental Health Sciences (NIEHS) grant numbers R01ES009860, P50ES00002, ES009860 and ES00002. The authors thank the participants and the leadership of Local 29 of the International Brotherhood of Boilermakers, Iron Ship Builders, Blacksmiths, Forgers and Helpers in Quincy, Massachusetts. They also thank Georg Schmidt for his expert contribution.

Contributors DCC, JMC and SCF drafted data collection tools. PEU, along with a team of others supervised by DCC collected data. MAM supervised the algorithm that generated AC and DC from the ECGs of participants. PEU, XL, CL and DCC designed research methods. PEU wrote the statistical analysis plan under supervision of XL. In addition, PEU cleaned and analysed the data, and drafted and revised the paper with inputs from all co-authors and supervision from DCC, CL and XL. DCC is the guarantor.

Funding Harvard ERC-NIOSH grant 5T42OH008416.

Competing interests None declared.

Patient consent Obtained.

Ethics approval Institutional Review Board, Harvard T. H. Chan School of Public Health.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Brunekeef B, Beelen R, Hoek G, *et al.* Effects of long-term exposure to traffic-related air pollution on respiratory and cardiovascular mortality in the Netherlands: the NLCS-AIR study. *Res Rep Health Eff Inst* 2009;5–71.
- 2 Rich DQ, Zareba W, Beckett W, *et al.* Are ambient ultrafine, accumulation mode, and fine particles associated with adverse cardiac responses in patients undergoing cardiac rehabilitation? *Environ Health Perspect* 2012;120:1162–9.

- 3 Lepeule J, Laden F, Dockery D, *et al.* Chronic exposure to fine particles and mortality: an extended follow-up of the Harvard Six Cities study from 1974 to 2009. *Environ Health Perspect* 2012;120:965–70.
- 4 Madrigano J, Kloog I, Goldberg R, *et al.* Long-term exposure to PM_{2.5} and incidence of acute myocardial infarction. *Environ Health Perspect* 2013;121:192–6.
- 5 Cavallari JM, Eisen EA, Chen JC, *et al.* Night heart rate variability and particulate exposures among boilermaker construction workers. *Environ Health Perspect* 2007;115:1046–51.
- 6 Magari SR, Schwartz J, Williams PL, *et al.* The association of particulate air metal concentrations with heart rate variability. *Environ Health Perspect* 2002;110:875–80.
- 7 Zanobetti A, Gold DR, Stone PH, *et al.* Reduction in heart rate variability with traffic and air pollution in patients with coronary artery disease. *Environ Health Perspect* 2010;118:324–30.
- 8 Cavallari JM, Fang SC, Eisen EA, *et al.* Time course of heart rate variability decline following particulate matter exposures in an occupational cohort. *Inhal Toxicol* 2008;20:415–22.
- 9 Vallejo M, Ruiz S, Hermosillo AG, *et al.* Ambient fine particles modify heart rate variability in young healthy adults. *J Expo Sci Environ Epidemiol* 2006;16:125–30.
- 10 Anastasiou-Nana MI, Terrovitis JV, Athanasoulis T, *et al.* Prognostic value of iodine-123-metaiodobenzylguanidine myocardial uptake and heart rate variability in chronic congestive heart failure secondary to ischemic or idiopathic dilated cardiomyopathy. *Am J Cardiol* 2005;96:427–31.
- 11 Erdogan A, Coch M, Bilgin M, *et al.* Prognostic value of heart rate variability after acute myocardial infarction in the era of immediate reperfusion. *Herzschrittmacherther Elektrophysiol* 2008;19:161–8.
- 12 Thayer JF, Yamamoto SS, Brosschot JF. The relationship of autonomic imbalance, heart rate variability and cardiovascular disease risk factors. *Int J Cardiol* 2010;141:122–31.
- 13 Perkiomaki J, Ukkola O, Kiviniemi A, *et al.* Heart rate variability findings as a predictor of atrial fibrillation in middle-aged population. *J Cardiovasc Electrophysiol* 2014;25:719–24.
- 14 Cavallari JM, Eisen EA, Fang SC, *et al.* PM_{2.5} metal exposures and nocturnal heart rate variability: a panel study of boilermaker construction workers. *Environ Health* 2008;7:36.
- 15 Kile ML, Fang S, Baccarelli AA, *et al.* A panel study of occupational exposure to fine particulate matter and changes in DNA methylation over a single workday and years worked in boilermaker welders. *Environ Health* 2013;12:47.
- 16 Cavallari JM, Fang SC, Mittleman MA, *et al.* Circadian variation of heart rate variability among welders. *Occup Environ Med* 2010;67:717–19.
- 17 Bauer A, Kantelhardt JW, Barthel P, *et al.* Deceleration capacity of heart rate as a predictor of mortality after myocardial infarction: cohort study. *Lancet* 2006;367:1674–81.
- 18 Schneider A, Hampel R, Ibaldo-Mulli A, *et al.* Changes in deceleration capacity of heart rate and heart rate variability induced by ambient air pollution in individuals with coronary artery disease. *Part Fibre Toxicol* 2010;7:29.
- 19 Bauer A, Barthel P, Schneider R, *et al.* Improved Stratification of Autonomic Regulation for risk prediction in post-infarction patients with preserved left ventricular function (ISAR-Risk). *Eur Heart J* 2009;30:576–83.
- 20 Eick C, Rizas KD, Zuern CS, *et al.* Automated assessment of cardiac autonomic function by means of deceleration capacity from noisy, nonstationary ECG signals: validation study. *Ann Noninvasive Electrocardiol* 2014;19:122–8.
- 21 Kantelhardt JW, Bauer A, Schumann AY, *et al.* Phase-rectified signal averaging for the detection of quasi-periodicities and the prediction of cardiovascular risk. *Chaos* 2007;17:015112.
- 22 Hauser R, Eisen EA, Pothier L, *et al.* A prospective study of lung function among boilermaker construction workers exposed to combustion particulates. *Am J Ind Med* 2001;39:454–62.
- 23 Fang SC, Eisen EA, Cavallari JM, *et al.* Acute changes in vascular function among welders exposed to metal-rich particulate matter. *Epidemiology* 2008;19:217–25.
- 24 Kim JY, Magari SR, Herrick RF, *et al.* Comparison of fine particle measurements from a direct-reading instrument and a gravimetric sampling method. *J Occup Environ Hyg* 2004;1:707–15.
- 25 Goldberger AL, Amaral LA, Glass L, *et al.* PhysioBank, PhysioToolkit, and PhysioNet: components of a new research resource for complex physiologic signals. *Circulation* 2000;101:E215–20.
- 26 Pope CA III, Hansen ML, Long RW, *et al.* Ambient particulate air pollution, heart rate variability, and blood markers of inflammation in a panel of elderly subjects. *Environ Health Perspect* 2003;112:339–45.
- 27 Magari SR, Hauser R, Schwartz J, *et al.* Association of heart rate variability with occupational and environmental exposure to particulate air pollution. *Circulation* 2001;104:986–91.
- 28 Guzik P, Piskorski J, Barthel P, *et al.* Heart rate deceleration runs for postinfarction risk prediction. *J Electrocardiol* 2012;45:70–6.
- 29 Langrish JP, Mills NL, Chan JK, *et al.* Beneficial cardiovascular effects of reducing exposure to particulate air pollution with a simple facemask. *Part Fibre Toxicol* 2009;6:8.
- 30 Chuang KJ, Chan CC, Su TC, *et al.* The effect of urban air pollution on inflammation, oxidative stress, coagulation, and autonomic dysfunction in young adults. *Am J Respir Crit Care Med* 2007;176:370–6.