

Contents

Editorial

Feasibility of route extrapolation in risk assessment W E PEPELKO *page 649*

Possible toxic metal exposure of prehistoric bronze workers M HARPER *page 652*

Inorganic particulates in pneumoconiotic lungs of hard metal grinders J R RÜTTNER, M A SPYCHER, I STOLKIN *page 657*

Exposure to respirable coalmine dust and incidence of progressive massive fibrosis J F HURLEY, W P ALEXANDER, D J HAZELDINE, M JACOBSEN, W M MACLAREN *page 661*

Development, radiological zone patterns, and importance of diffuse pleural thickening in relation to occupational exposure to asbestos H BOHLIG, A CALAVREZOS *page 673*

Possibility of inducing glandular stomach cancer in rats exposed to asbestos F M KOGAN, N N VANCHUGOVA, V N FRASCH *page 682*

Surface character and membranolytic activity of rutile and anatase: two titanium dioxide polymorphs R P NOLAN, A M LANGER, I WEISMAN, G B HERSON *page 687*

Calculating expected deaths: a comparison of two methods A C CHOUIL *page 699*

An analysis of the validity of self reported occupational histories using a cohort of workers exposed to PCBs C R ROSENBERG, M N MULVIHILL, A FISCHBEIN, S BLUM *page 702*

Exposure to vinyl chloride monomer: report on a cohort study AGNÈS LAPLANCHE, FRANÇOISE CLAVEL, J-C CONTASSOT, CLAUDINE LANOUZIERE *page 711*

Correspondence

Update on lung disease in coal miners W K C MORGAN *page 716*

Incidence of cancer among vinylchloride and polyvinylchloride workers: further evidence for an association with malignant melanoma W F TEN BERGE *page 718*

Pulmonary fibrosis in asbestos insulation workers with lung cancer Y SUZUKI, H KIPEN, RUTH LILIS, I J SELIKOFF *page 719*

Notice *page 720*