Occupational hazards in hospitals: accidents, radiation, exposure to noxious chemicals, drug addiction and psychic problems, and assault

J J GESTAL

From the Department of Preventive and Social Medicine, Faculty of Medicine, University of Santiago, Santiago de Compostela, Spain

ABSTRACT Except for infectious diseases all the main occupational hazards affecting health workers are reviewed: accidents (explosions, fires, electrical accidents, and other sources of injury); radiation (stochastic and non-stochastic effects, protective measures, and personnel most at risk); exposure to noxious chemicals, whose effects may be either local (allergic eczema) or generalised (cancer, mutations), particular attention being paid to the hazards presented by formol, ethylene oxide, cytostatics, and anaesthetic gases; drug addiction (which is more common among health workers than the general population) and psychic problems associated with promotion, shift work, and emotional stress; and assault (various types of assault suffered by health workers, its causes, and the characterisation of the most aggressive patients).

In a previous paper attention was drawn to the current importance of occupational hazards to hospital workers.¹ These were classified into six categories (infection, accidents, radiation, exposure to noxious chemicals, drug addiction and psychic problems, and assault), and the risk of infection was discussed. In the present paper the remaining categories are considered.

Accidents

Among the possible types of accidents are included fires, explosions, electrocution, and gas leaks; undesired effects of electromagnetic fields, microwaves, lasers, and vibration; cuts, bruises, and fractures; asphyxia and burns; and the effects of noise.

EXPLOSIONS

Explosions usually occur in laboratories, operating theatres, and boiler rooms. All products that are inflammable or otherwise likely to explode should be labelled as such and stored accordingly.²

FIRE

Official regulations concerning the risk of fire in hospitals are in general adequate. Each centre should draw up fire procedures and all hospital workers

should be instructed as to what to do in case of fire. Of 300 fires in United States hospitals, 74% were due to the following four causes, human error being ultimately responsible in almost all cases: electrical faults (23%), cigarettes and matches (21%), incorrect use of anaesthetics, oxygen, and inflammable fluids (19%) and non-electrical heating (11%).

The most common direct cause of casualties associated with hospital fires is asphyxia by smoke and fumes, which accounts for 78% of deaths and 43% of non-fatal casualties.³ Fire prevention measures should aim to achieve four main objectives in cases of fire: immediate detection, rapid extinction or isolation, non-propagation of smoke and fumes, and safe evacuation of staff and inmates. To these ends attention should be paid to four basic points:

- (1) Construction—for example, evacuation routes and fire escapes.
- (2) Use of combustible materials such as plastics or inflammable fluids.
- (3) Countermeasures, including compulsory installation of fire extinguishers.
- (4) Training of personnel.

The hospital's fire procedures should be familiar to all staff so that they all know exactly what to do in case of fire. Fire procedures should furthermore be reviewed at least every six months and corrected as necessary, and fire drill should be practised in all shifts.⁴

Accepted 5 August 1986

Occupational hazards in hospitals

ELECTRICAL ACCIDENTS

Electrical accidents may cause injury both directly, due to the passage of electric current through the body producing electrocution, internal and external burns, or gaseous embolism; and indirectly, in the form of burns or asphyxia produced by electrical fires or explosions, or injuries suffered in falls after electric shock. The immediate physiological effects of electric shocks depend on the intensity of the current borne, the point of contact, and the path taken through the body. Currents near the threshold of perception (1–3 mA) are innocuous even when there is prolonged contact, but as the intensity of the current rises the tingling sensation becomes progressively more unpleasant. Sudden application of 8 mA generally stimulates a reflex response which may itself bring about injury—due, for example, to a consequent fall off a ladder—but electrification by currents of this magnitude does not normally have direct physiological consequences. Currents greater than about 10 mA produce muscular contractions that lock the hands and arms and prevent the release of the objects held. Burns may also appear. When the current borne is of more than 25–30 mA, spasm may affect the muscles of the thorax and cause “asphyxia” if artificial respiration is not rapidly applied.

Electric shock may be fatal if the current passes through the brain or the thorax, especially if the heart is affected. Since electric current takes the path of least resistance, particular danger thus attends double contacts in which the body forms a bridge between a live wire and a conductor (right hand and left foot or vice versa, right and left hands, or hand and head). Double contacts affecting a single limb are less serious.

There are two contributions to the electrical resistance of the human body: the external resistance of the surface at which contact is made and the resistance of internal tissues. The external resistance varies from one individual to another and from one part of the body to another (a calloused palm is much more resistive than the back of the same hand), and also depends greatly on the humidity of the surface (due, for example, to sweating: more than 60% of electrical accidents suffered by hospital personnel occur during the hottest five months). A skin with a resistance of several hundred thousand ohms when dry may offer just 1000 ohms resistance or less when wet. The resistance of the area of contact also decreases as the contact pressure increases. The resistance of internal tissues varies less and is estimated at between 100 and 5000 ohms.

The chief electrical hazard in hospitals is the large static charge that may accumulate in electrical apparatus. Arcing may occur between charged material and neighbouring conductors, and the human body

may receive electric discharges if insulated from earth by rubber soled footwear. In fact, if the atmosphere is dry just walking about on asphalt flooring, or even on a woollen carpet, can charge the body sufficiently as to produce small sparks that, though not normally harmful in themselves, may ignite inflammable vapours and gases, cause dust explosions, or (more commonly) startle the person in question so that other types of accidents occur. An idea of the risk of sparking may be given by the fact that an insulated human body dressed in nylon overalls can easily acquire up to 5 or 10 mJ of electrostatic energy just by sitting down on a chair covered in polyvinyl chloride and getting up again.

The control of electrostatic charge is difficult and generally entails facilitating its loss. In the case of the human body both footwear and flooring should be sufficiently conductive as to prevent the accumulation of charge. Flooring should be of wood, footwear of antistatic rubber, and polyester or polyamide clothing should be worn. The risk of sparking may also be reduced by increasing the humidity of the air and by wearing clothing with antistatic coatings (although most of the latter are lost in the wash, some have appeared in recent years which are relatively permanent). As regards other types of electrical accident, apart from the instruction of personnel in the proper use of electricity and the frequent periodic maintenance of apparatus and mains wiring, the measures to be taken to minimise risk include the earthing of all apparatus, the installation of circuit breakers, the use of low mains voltages, and the separation of mains circuits.⁵

CUTS, BRUISES, AND FRACTURES

Cuts, bruises, and fractures may occur in hospital work just as in any other human activity. Dominguez-Carmona, for example, mentions the case of a nursing nun who was killed in 1976 in the Hospital Clinico, Madrid, when a lift she was entering started up on being called from a higher floor, the second fatal accident of this type in the centre in question.⁶ The risk of accident is greatly increased by physical fatigue, which is common towards the end of the day's work but varies in intensity from one department to another and also depends on the building in which the department is housed and the type of personnel concerned. In old fashioned buildings it may be necessary to do a lot of walking from place to place, and even in well designed hospitals nurses spend most of the day on their feet, with the consequent prevalence of varicose veins and foot complaints. Long working hours, changes of shift, and the arrangements before such changes, eating hours, and coffee or tea breaks, and free time and holidays all greatly affect the fatigue and general health of nursing staff. Training and

qualifications nevertheless reduce the risk of accidents: a survey carried out in the United States showed that nurses who had received a diploma suffered fewer injuries than less qualified nursing staff. Among nurses with a diploma about 30% of all accidents were due to overeffort and 20% to falls.⁷ Accidents due to overeffort when lifting heavy weights, however, are far from being restricted to hospital nurses; domestic nurses being quite as much at risk, if not more. The risk of accidents from falling objects, burns, machines, beds, operating tables, glassware, needles, or scalpels is similar to that encountered in other occupations in which comparable equipment is used.

The nature of dental work exposes dentists to a special set of hazards. The use of pointed metal instruments carries a risk of pricks and cuts (which may also be caused by the edges of gold or steel crowns) and of face and eye wounds caused by fragments of tooth or metal when drilling. In the past, painful and deforming cervical, dorsal, and lumbar complaints were also frequent as the result of the dentist spending all day on foot leaning over the chair. Modern workposts have largely solved this problem by allowing the patient to be positioned so that the dentist may work comfortably while seated or standing.

NOISE

Noise constitutes an increasing problem in hospitals and clinics as elsewhere. No cases have been reported in dentists in which actual loss of hearing may be attributed to the use of low, medium, or high speed drills, though the latter are extremely noisy⁷; but it is less clear whether the noise made by drills may not give rise to psychological problems such as loss of concentration, character changes, or irritability.⁷ Other health workers likely to be exposed to high noise levels produced by certain of the machines they work with include traumatologists, orthopaedic surgeons, and ear, nose, and throat specialists.⁸ The harmfulness of noise depends not only on the characteristics of the noise itself (its intensity and cadence)⁹ and those of the place in which the noise is experienced but also on the susceptibility of the exposed subject,¹⁰⁻¹⁵ many cases of acoustic trauma having been reported in which the noise level was not particularly high.^{16,17} It should also be remembered that quite soft noises of certain types may be extremely unpleasant and their continual repetition may constitute a long term psychic health hazard. In the United States the recommended maximum noise levels for hospital wards are 45 dB (A) during the day and 35 dB (A) at night.¹⁸ Hospital rooms should be suitably sound proofed, noisy faults in taps, cisterns, trolleys, and so on should be repaired as quickly as possible,

soft soled footwear should be worn, and medical system alarms and telephones should be muffled. Other suggested measures for limiting noise levels in hospitals include restrictions on conversations between staff and patients with hearing difficulties and the isolation of noisy patients.¹⁹

There have been several studies of the economic effects of accidents in various Spanish hospitals.²⁰⁻²³ R Fernández-Crehuet estimated that in the five years 1978-82 the working days lost through accident in the Hospital "Reina Sofía" in Cordoba cost 16 117 629 pesetas (about £80,600).

Radiation

Serious hazards are presented to health workers by the medical use of ionising radiation. The radiation to which hospital staff may be exposed includes both photonic radiation (x rays and γ rays) and charged particle radiation (α and β rays).

x Rays are generated by conventional radio-diagnostic and x ray therapy equipment and by high energy x ray tubes. Gamma rays are produced by cobalt and caesium bombs and by radioelements encapsulated in needles, tubes, or pearls, which may also emit β rays. Finally, unencapsulated radioelements are used in solutions or colloidal suspensions administered to patients and laboratory animals or applied in radionuclear laboratories to biological material obtained from patients or animals. The use of these unencapsulated sources entails the risk of laboratory equipment, workspaces, and personnel becoming contaminated.^{24,25}

IONISING RADIATION

The effects of exposure to ionising radiation appear sooner or later after a latency period and may be either stochastic or non-stochastic. Most of the cases mentioned in textbooks of radiobiology involve either massive exposure to radiation used for military purposes or doses of the same order of magnitude as those administered to patients subjected to radiotherapy. Personnel working in properly constructed centres should never be exposed to such large doses so long as they observe the necessary precautions,²⁵ but many health workers are none the less the object of chronic low intensity radiation and as a consequence a small percentage may suffer somatic and genetic stochastic effects and possibly certain non-stochastic effects also. In 1959 it was reported that there had been 359 deaths from radiological causes throughout the world.

Non-stochastic somatic effects

Non-stochastic somatic effects may arise in the haematopoietic system, with occasional hypoplasia or

Occupational hazards in hospitals

even aplasia, and yearly or half yearly analyses of peripheral blood should be carried out to check for this condition. Dermatological effects may include brittle, cracked, or grooved nails, the disappearance of body hair and fingerprints, and chronic radiodermitis. Cataracts may develop. Irradiation of embryos or intrauterine fetuses by doses of more than 1 rem may cause abortion, fetal death, or the serious malformation of those fetuses that complete their gestation. Alterations may also be produced by unencapsulated radioisotopes able to cross the placental barrier.

Stochastic effects

Stochastic effects include the shortening of life, the induction of cancer and leukaemia, and genetic effects (an increase in the frequency of mutations).^{26 27}

Nowadays, the main radiation hazard occurs not in hospital radiological departments²⁸⁻³⁰ but in the consulting surgeries of general practitioners and small private clinics where adequate safety measures have not been put into effect, and in hospital departments other than the radiological department that may also have occasion to use radiation (operating theatres and casualty and paediatric departments, for example). Professional radiological staff are also less likely to expose themselves to radiation than other personnel such as casualty staff or paediatricians, among whom cases of radiodermitis and skin cancer still sometimes occur. The most dangerous operations for these workers include radioscopy and the use of x ray surveillance of the setting of fractures, searches for foreign bodies, and the introduction of catheters.

NON-IONISING RADIATION

Theoretically, non-ionising radiation used in hospitals may also constitute a health hazard. Such radiation includes ultraviolet light, laser beams, magnetic fields, and radiofrequencies. Ophthalmologists who work many hours a week with lasers have been reported to suffer from reduced central visual acuity, abnormal colour perception, and other defects indicative of possible macular damage,³¹ and although these claims have not been confirmed, various bodies have established norms for the use of laser equipment.³²⁻³⁴ Monitor screens have been blamed for eye strain, postural complaints, and psychological alterations leading to "technostress."³⁵⁻⁴² In general, however, the risk of health problems derived from non-ionising radiations is negligible. Perhaps the only individuals clearly put at risk by a subclass of radiations of this type are those who have had pacemakers implanted, since exposure to strong magnetic fields such as those used in nuclear magnetic resonance equipment will probably disrupt the timing of the pulses generated by the pacemaker.

Exposure to noxious chemicals

Many of the hazards faced by hospital staff consist in the wide variety of chemical substances used. Apart from cutaneous infections, dermatological effects of other infectious diseases, and infestation by parasites, and leaving aside the now quite rare occurrence of radiodermitis discussed above, the cutaneous occupational diseases suffered by hospital workers may be classified as either irritations (caused by chemical disinfectants and detergents) or allergies (caused, for example, by antibiotics, rubber, nickel, disinfectants, aldehydes, or phenothiazines). The occupational origin of dermatosis may be ascertained on the basis of the shape of the area affected, its distribution and location, the recurrent coincidence in time with particular occupations, the patch test, and the response to treatment. Skin complaints make up half the total number of occupational illnesses both among hospital workers and among the working population in general, 90% of them being cases of contact dermatitis. Some 60-80% of these cases are caused by chemical irritants (eczema due to primary irritants) and the remainder are of an allergic nature (allergic eczema).

Occupational allergic eczema is more common among nurses, surgical assistants, and other auxiliaries than among doctors because of their continual contacts with drugs, anaesthetics, and antiseptics. Resistance is reduced by repeated washing with soap and, especially, by scrubbing the hands and forearms, which destroys the skin's protective layer of fatty acids. Surgical and casualty personnel may suffer from "dry hand syndrome" due to handling plaster of Paris. Dentists and dental technicians are often affected by contact eczema caused by handling acrylic monomers, local anaesthetics, essential oils, dental mould paste, epoxy resins, and amalgams. Laboratory personnel are likewise put at risk by many of the chemicals they handle.⁴³⁻⁴⁷

Mutagenic or carcinogenic effects result from many substances used or found in hospitals. Thus beta-propiolactone is known to be carcinogenic⁴⁸⁻⁵⁰; hexachlorophene is neurotoxic when applied to the skin, as was illustrated by the neural degeneration suffered some years ago by French infants to whom it was applied in talc, and its teratogenic effects in people are also currently being investigated⁵¹⁻⁵³; sodium orthophenylphenate and glycols have been responsible for cancer of the bladder; and benzol (and xylol, toluene, and xylene contaminated with benzol^{54 55}) produces leukaemia and chromosomal aberrations. Busch and Nelson have reported cases of acute respiratory distress in histological laboratory staff exposed to xylene,⁵⁶ which may also give rise to headaches, nausea, loss of appetite, lassitude, and

impaired coordination.⁵⁷

FORMOL

Since 1894 formol has been used for terminal disinfection of hospital rooms, beds, and apparatus. It is also used as a preserving agent in pathological anatomy and in the preparation of Salk and hepatitis B vaccines. Apart from these medical uses of formol in hospitals,⁵⁸⁻⁶² there are also non-medical sources, including chipboard and plywood, carpet glues, parquet varnishes, and urea-formol foam plastics used as thermal insulators.⁶³⁻⁷⁰ The emanation of formaldehyde from these materials depends on the temperature, the humidity, the age of the materials, and the technical quality of the products.⁶⁸⁻⁷¹ A volume concentration of 40 ppm of formaldehyde is also contained in tobacco smoke,⁷²⁻⁷⁴ and an individual smoking 20 cigarettes a day may retain some 0.38 mg.

Uptake of formaldehyde occurs chiefly through the digestive and respiratory systems, absorption through the skin being negligible.⁷⁵ Small quantities are normal in the cell. Its half life in the blood is about one and a half minutes.

In recent years several comprehensive reviews of the toxicity of formaldehyde have appeared.⁷⁶⁻⁷⁹ The main effect of exposure to low concentrations is to irritate the eyes and the upper respiratory tract,⁸⁰⁻⁸⁵ though other parts of the respiratory system and the skin may also become sensitised.⁸⁶⁻⁸⁸ A concentration of 10 ppm begins to be unbearable, giving rise to burns in the nose, trachea, and eyes accompanied by intense lacrimation, coughing, palpitations, and a sensation of thoracic and cephalic oppression. Concentrations of 50-100 ppm produce pulmonary inflammation and oedema and ultimately death.⁸⁹

The North American Contact Dermatitis Group have ranked formaldehyde as tenth in a list of those chemicals causing most skin reactions.⁹⁰ Allergic dermatitis caused by contact with formaldehyde has been found in the medical sphere among nurses handling thermometers sterilised in 100 g/l formaldehyde solution,⁹¹ personnel using disinfectants containing formaldehyde,⁹² and patients whose injuries have been dressed with plaster impregnated bandages containing 100 g/l of melamine formaldehyde resin.⁹³⁻⁹⁴ In 1966 Blejer and Miller described an outbreak of contact eczema in a haemodialysis unit where a 20 g/l solution of formol was used to sterilise open tanks.⁹⁵ Specific sensitisation of the respiratory tract is also possible, though this is much less common than sensitisation of the skin.⁹⁶⁻⁹⁷ In recent years the automating of sterilisation processes, better ventilation, and greater awareness of the need to use protective systems has reduced the incidence of such cases.⁹⁸⁻⁹⁹

Formaldehyde is known to be highly reactive with nucleic acids and proteins,¹⁰⁰⁻¹⁰² and several studies

have shown its mutagenic effects on various micro-organisms,¹⁰³⁻¹⁰⁷ arthropods,¹⁰⁸⁻¹¹⁰ cell test systems,¹¹¹⁻¹¹⁶ and mammals.¹¹⁷ Since squamous cell carcinomas of the nasal passages of rats and mice exposed to 6-15 ppm of formaldehyde were observed in the United States,¹¹⁸⁻¹²⁰ there has been controversy world wide as to whether formaldehyde is or is not carcinogenic,¹²¹⁻¹²⁴ and there have been numerous studies of morbidity¹²⁵ and mortality¹²⁶⁻¹³⁵ among groups of workers subject to occupational exposure to formaldehyde. No abnormally high rates of mortality from nose or lung cancer have been observed in these groups, but there have been reports of higher than average mortality from tumours of the prostate, skin, kidney, brain, mouth, colon, bladder, and bone, and from leukaemia. Most of these studies are open to the criticism that the small number of subjects prevented significant conclusions being reached—for example, in the case of nose cancer—but a much larger study has recently been carried out by Acheson *et al.*, who found no evidence of higher than average rates of mortality from any of the tumours mentioned in earlier studies (including nose cancer) and concluded that formaldehyde is not carcinogenic in man.¹³⁶ Further research on this subject is nevertheless still needed before definitive conclusions can be arrived at.

The attitude of official bodies to formaldehyde has exhibited the same incoherence as scientific publications. In 1981 the National Institute for Occupational Safety and Health declared formaldehyde to be a possible carcinogen,¹³⁷ and in April 1982 the United States Consumer Products Safety Commission responded to numerous complaints regarding the irritant effects of formaldehyde in the respiratory tract and its possible carcinogenic potential by banning the use of urea-formol foams as insulators in the building trade.¹³⁸ A year later the ban was lifted by the Court of Appeal on the grounds of the incorrectness of the documentation presented by the CPSC regarding evidence of the health hazard posed by the chemicals given off by the foams.¹³⁹ In May 1984 the Environmental Protection Agency nevertheless included among its priorities review of the use of urea-formol resins in housing.¹⁴⁰ Similar situations have been created in other countries. A World Health Organisation working party recently recommended that because of its possible carcinogenic effects, formaldehyde concentrations at work should be reduced as much as possible, and that as a provisional limit workers should in any case not be exposed to concentrations in the air of more than 0.5 mg/m³ for a mean eight hours a day in a working week of 40 hours.⁷⁶

ETHYLENE OXIDE

The bactericidal properties of this colourless gas were

Occupational hazards in hospitals

first made known in 1929,¹⁴¹ and since 1962 it has been widely used to sterilise medical material. At present it is the disinfectant gas most widely used to sterilise plastics and other materials that cannot be subjected to heat. Its acting at room temperature also makes it economical. Its disadvantages, however, include its being explosive, which makes it necessary to mix it with inert gases; its remaining impregnated in the material being sterilised (it dissolves in rubber), which means that the material sterilised must subsequently be ventilated; and its toxicity, concentrations of 50 000–100 000 ppm being lethal in a few minutes.¹⁴²

In the United States more than 2000 million kilos of ethylene oxide are produced yearly. Although less than 1% of this total production is used for industrial or medical sterilisation, this use constitutes the greatest health hazard, with an estimated 100 000 health workers at risk in the sterilisation services of United States hospitals. Whereas other industrial applications use ethylene oxide in closed circuits or outdoors, sterilisation involves open processes carried out in closed premises.¹⁴³ The areas of greatest risk are in general the sites where ethylene oxide chambers are emptied, the most dangerous moment being when the door is opened.

Exposure to ethylene oxide, usually either contact with the skin or inhalation, may produce either local or generalised effects. The exposed individual's defensive reactions are hampered by the typical lag of several hours between exposure and the appearance of symptoms, and by the fact that the gas is imperceptible to smell until quite high concentrations are present (700 ppm).¹⁴⁴ Contact with insufficiently ventilated materials containing ethylene oxide (such as gloves, face masks, overalls, tampons, endotracheal tubes, or anaesthetic masks) produces irritant cutaneous lesions,^{145–149} conjunctivitis and corneal burns,¹⁵⁰ and, after exposure to high concentrations, cataracts.^{151–153} Repeated contact may give rise to allergic sensitisation.¹⁵⁴ Generalised effects may include acute poisoning, the seriousness depending on the intensity of exposure. Mild symptoms include nausea and vomiting¹⁵⁵ and irritation of the eyes, nose, and throat, whereas in the more serious cases there may be respiratory problems (dyspnoea, cyanosis, and pulmonary oedema), which may be produced by just a few minutes exposure to concentrations of 500–700 ppm.¹⁴⁴ Other symptoms may affect the heart, the nervous system (with headaches, sleepiness, weakness, lack of coordination, and convulsions),¹⁴⁴ the blood¹⁵⁶ and anaphylaxis.¹⁵⁷ Chronic ethylene oxide poisoning may cause encephalopathy, polyneuritis, and neurovegetative lesions.¹⁵⁸ Exposure during gestation has been reported to induce premature deliveries¹⁴⁴ or abor-

tions.¹⁵⁹

There are no data suggesting that ethylene oxide has teratogenic effects in man, and the results obtained with laboratory animals are contradictory.^{160 161} As an alkylating agent,¹⁶² many studies have confirmed its mutagenic action in micro-organisms,¹⁶³ arthropods,¹⁶⁴ plants,^{165 166} and the cells of man^{167 168} and other mammals.^{169 170} The frequency of chromosomal and chromatinic anomalies in lymphocytes is also known to be higher among people chronically exposed to ethylene oxide than among the general population.^{171–175} Rats exposed to ethylene oxide atmospheres have been observed to suffer from leukaemia and mesothelioma,¹⁷¹ and subcutaneous inoculation in mice increases the incidence of fibrosarcomas.¹⁷⁶ The epidemiological studies conducted among people are nevertheless insufficient to enable definitive conclusions regarding its possible carcinogenic action, for whereas Hogstedt *et al*^{177 178} reported an increase in both general mortality and mortality from tumours (especially leukaemias), a study carried out in the United States found no such effect.¹⁷⁹ Uncertainty regarding the duration and intensity of exposure hinders the drawing of conclusions, and further epidemiological studies are required.

The degree to which an individual has been exposed to ethylene oxide may be measured biologically by its alkylating haemoglobin and binding to histidine to give N-3(2-hydroxyethyl) histidine.¹⁸⁰ Legally established tolerable limits vary from one country to another. In 1971 the American Conference of Governmental Industrial Hygienists proposed 50 ppm as the mean allowable concentration in the workplace,¹⁸¹ but information gathered concerning mutagenic and possible carcinogenic effects has led to a legal threshold limit value (TLV) of 10 ppm and a proposal in 1982 for a reduction to 1 ppm,¹⁴³ the same level proposed in 1983 by the Occupational and Safety Health Administration.¹⁸² In the USSR the limit is 0.5 ppm. After comparing the alkylating and mutagenic potential of ethylene oxide with that of ionising radiation, Calleman *et al* proposed a TLV of 0.25 ppm, which would be equivalent to the maximum allowed dose of 5 rads of radiation a year.¹⁸⁰

In the light of the data now available, it may be concluded that whenever possible sterilisation by ethylene oxide should be replaced by sterilisation by γ rays or propyl oxide.

CYTOSTATICS

Certain oncolytic agents have long been known to produce contact dermatitis and blisters, and elementary precautions have always been taken to avoid their contact with the skin of the patients to whom they are administered and the nurses responsible for

their administration. Modern studies, however, have shown that the mutagenic hazard presented by cytostatics (found by Hoffman¹⁸³ in the urine of nurses and technicians who had prepared or administered antitumour agents) is not limited to physical contact but also includes the inhalation of vapours and microdroplets given off during the preparation and administration of cytostatic solutions, and these mutagenic effects are reinforced by smoking. Immunological alterations have been observed in the lymphocytes of the peripheral blood of both patients treated with cytostatics and the staff who had prepared and administered the drugs.¹⁸⁴⁻¹⁸⁷ To eliminate these risks, cytostatics should be prepared and reconstituted in vertical laminar flow chambers using a laboratory coat, gloves, and a face mask, and the same rules should apply to the handling of the urine of patients treated with these substances.¹⁸⁸⁻¹⁹⁰

ANAESTHETIC GASES

The first reports of harmful effects of anaesthetic gases among health workers are those of Hewitt (1893), Kirschner,¹⁹¹ and Perthes (1925),¹⁹² all of whom described acute effects. Hirsch and Kappurs soon pointed out that although acute poisoning is infrequent, the likelihood of chronic poisoning cannot be neglected,¹⁹³ and cases of chronic poisoning were described by Werthmann in 1948.¹⁹⁴ Nevertheless, this problem was generally ignored until in 1967 Vaisman carried out an epidemiological study of Soviet anaesthetists,¹⁹⁵ since when a variety of disorders have been attributed to anaesthetic gases.¹⁹⁶⁻¹⁹⁸ Following Cascorbi,¹⁹⁹ we shall classify them into three main groups: changes in behaviour and the ability to perform psychometric tests, diseases of the organs for the biotransformation and excretion of xenobiotics, and faulty cell replication.

Changes in behaviour and the ability to perform psychometric tests

Bruce *et al* found that volunteers exposed to trace concentrations of anaesthetic gases (50–500 ppm of N₂O with or without 1, 10, or 15 ppm of halothane) suffered alterations of perception, cognition, and motor skills.²⁰⁰⁻²⁰² Neither Smith and Shirley,²⁰³ nor Frankhuizen *et al*,²⁰⁴ nor Venables *et al*,²⁰⁵ however, have been able to reproduce their results, which are furthermore questionable as regards their statistical significance¹⁹⁷ and the surprising finding that nitrous oxide alone was more deleterious than when accompanied by halothane.²⁰² Occupational exposure to traces of anaesthetic gases has not been observed to give rise to neuropsychiatric disorders.²⁰⁶⁻²⁰⁸ Quimby *et al* found that rats exposed to 10 ppm of halothane for eight hours a day, five days a week from birth to the age of 60 or 135 days, made

more mistakes than controls in specific tests but that their relative learning rates were similar.²⁰⁹⁻²¹⁰ Again, however, these results and their statistical significance have been questioned.¹⁹⁷ Present knowledge suggests that no acute behavioural alterations are produced by concentrations of nitrous oxide below 8–12% or by less than 0.1% of halothane.²¹¹⁻²¹³

A finding that may or may not be relevant to the toxicity of anaesthetics is that among United States doctors, whose overall suicide rate is 15% higher than that of the general population,²¹⁴⁻²¹⁵ anaesthetists are the second most suicidal subgroup (behind psychiatrists), with a rate three times that of the control group.²¹⁶⁻²¹⁷ This has been confirmed by Lew in a study of the causes of death among members of the American Society of Anesthesiologists,²¹⁸ which has named suicide as the major health risk among anaesthetists aged under 55.²¹⁹⁻²²⁰

Diseases of the organs for the transformation and excretion of xenobiotics

Hepatic diseases—Since the introduction of halothane in 1956, occasional cases of hepatic necrosis after its use as an anaesthetic have been reported,²²¹⁻²²² and repeated anaesthesia under halothane is now regarded as an aetiological factor that cannot be ignored.²²³ Subanaesthetic doses of halothane, isoflurane, diethyl ether, fluoxene, enflurane, and nitrous oxide have been found by some authors to cause hepatic alterations in animals,²²⁴⁻²²⁷ whereas others have observed no such changes even after lengthy or intense exposure²²⁸⁻²³⁰; controversy on this point continues.¹⁹⁷⁻²³¹⁻²³² In operating theatre staff temporary rises in hepatic transaminase levels have been reported,²³³ and also jaundice²³⁴ and hepatic cirrhosis,²³⁵ whereas in the United States Cohen *et al* found that the incidence of hepatic diseases was above average among female operating theatre staff²³⁶ and dentists using general anaesthesia three hours a week or more.²³⁷ An increase in the frequency of hepatic diseases has also been observed among anaesthetists in Czechoslovakia²³⁸ and England and Wales.²³⁹

Kidney disease—Certain epidemiological studies have reported a higher than average rate of kidney disease among anaesthetists, women especially.²¹⁶⁻²³⁶⁻²³⁸; Chang *et al* observed ultrastructural changes in the kidneys of laboratory animals chronically exposed to low levels of halothane²⁴⁰; and Dahlgren found moderate temporary alterations of kidney function in Swedish health workers exposed to methoxyflurane.²⁴¹

Faulty cell replication

Abortion and congenital malformation—In Vaisman's report attention was drawn to the high rate of abor-

tions among women working in operating theatres in the Soviet Union.¹⁹⁵ The same phenomenon was subsequently observed by Askrog and Harvald in Denmark²⁴² and was confirmed by Cohen *et al* in the United States and Canada,²³⁶ where the abortion rate among pregnant women working in operating theatres was 17.1% whereas that of other pregnant hospital workers was only 8.9%. Other investigators have reached similar conclusions,^{238 243 244} though there have also been contradictory reports in which no increase in the spontaneous abortion rate was observed among doctors and nurses exposed to anaesthetic gases.^{245 246} Of the various dangers attributed to chronic exposure to traces of anaesthetic gases, the risk of abortion among women working in surgical departments is the only one recognised by the United Kingdom Department of Health and Social Security.²⁴⁷

A more surprising finding has been that abortions are twice as common among the wives of United States dentists administering general anaesthesia at least three hours a week as among the wives of those who use only local anaesthesia (18% v 9%),²³⁷ though Askrog and Harvald had already reported a similar phenomenon among the wives of Danish anaesthetists.²⁴² This suggests that anaesthetic gases may affect spermatogenesis, and such alterations have indeed been observed in animals,^{248–251} but Wryobek *et al* found no changes in either the morphology or the concentration of spermatozoa produced by anaesthetists working in three San Francisco hospitals (though the modern gas extraction systems in these centres may have meant that these anaesthetists enjoyed better than average protection).²⁵²

Several studies have reported that women exposed to anaesthetic gases during gestation run a greater than average risk of giving birth to children with congenital malformations.^{236 253–256} Tomlin found that the central nervous system and musculoskeletal system were especially affected by this abnormal incidence of malformations,²⁴⁴ whereas Pharaoh *et al* observed higher than average frequencies of congenital cardiovascular malformations and stillborn births and a smaller average size among live newborn babies.²⁴⁵ Baltzar, however, found no such differences.²⁵⁷ The validity of many of these studies^{237 242 244 256 257} has in any case been questioned^{258–262} on the grounds that most were carried out using retrospective postal questionnaires. Fersting,²⁶⁰ Axelsson and Rylander,²⁶¹ and Tannenbaum and Goldberg²⁶² have been especially critical, pointing out numerous methodological flaws such as the lack of criteria for exposure or outcome, poor survey response rates, selection bias, lack of validation of outcome, recall bias, and lack of control

of potentially confounding variables.²⁶² These critics emphasise the need to carry out suitably designed prospective studies to determine whether or not exposure to trace concentrations of anaesthetic gases entails a real risk of abortion or congenital malformation.

Chronic exposure to anaesthetic concentrations of nitrous oxide or halothane has been reported to increase the resorption and congenital skeletal malformation rates among rats exposed during gestation^{263–267} or to reduce the weight and length of hamster fetuses,²⁶⁸ but no such effects have been observed at subanaesthetic levels.^{267 269–271} Neither have changes been observed in the fertility of animals exposed to subanaesthetic doses,^{248 269 272} or in exposed chick embryos.^{273–280} Mutagenic effects on bacteria have been observed by some^{281 282} but not by others.^{283 284} To sum up, there is at present no conclusive experimental evidence that chronic exposure to traces of anaesthetic gases—or even short exposure to high concentrations²⁸⁵—has any adverse effects on gestation in animals.

Malignancies—Chloroform^{286 287} and trichloroethylene²³⁸ have been found to be potentially carcinogenic (causing hepatic carcinomas and kidney tumours) for experimental animals exposed to large doses introduced by mechanisms different from those involved in the occupational exposure of man. With the exception of a study by Corbett,²⁸⁸ however, the results of which he was unable to reproduce,²⁸⁹ there is no experimental evidence that low concentrations of halothane, nitrous oxide, enflurane, isoflurane, or methoxyflurane are carcinogenic.^{290–294}

Bruce *et al* detected an abnormally high rate of mortality from lymphoid and reticuloendothelial tumours among anaesthetists during the period 1947–66,²¹⁶ but a subsequent study failed to confirm any relation²¹⁷; Cohen *et al* observed a greater incidence of cancer among female anaesthetists in their nationwide study²³⁶ but found no difference between exposed and unexposed subjects in their study of dentists²³⁷; and although Corbett *et al* reported an above average cancer rate among anaesthetic nurses in Michigan,²⁸⁸ serious doubts have been raised as to the validity of both this study and Cohen's.^{258 259} The suggestion that anaesthetic gases may act as transplacental carcinogens is not supported by the studies of Knill-Jones *et al*²⁴³ and Pharaoh *et al*²⁴⁵ in which the incidence of cancer is no greater among the children of female anaesthetists in England and Wales than among those of other women, though a difference between the two groups has been claimed by Tomlin.²⁴⁴ In view of these studies it cannot be concluded that there is a risk of cancer from chronic exposure to traces of anaesthetic gases.

Other risks, and preventive measures

Irritation of the respiratory tract has been reported after chronic inhalation of ether, and laryngitis and attacks of asthma after exposure to halothane and enflurane respectively.²⁹⁵ Personnel long exposed to nitrous oxide have been reported to suffer neuropathies²⁹⁶⁻²⁹⁸ recently shown to be caused by the ability of nitrous oxide to oxidise vitamin B₁₂, thus impeding the synthesis of methionine, folate, and thiamine.²⁹⁹⁻³⁰³

To prevent abnormalities caused by exposure to anaesthetic gases it is necessary to make anaesthetic and surgical staff aware of the risks and to urge the introduction of efficient gas extraction systems (levels of 130-7000 ppm of nitrous oxide and 10-85 ppm of halothane have been detected in the region where the anaesthetist breathes).³⁰⁴⁻³⁰⁶ Good practice codes should be adopted to reduce atmospheric pollution, routine inspections of equipment should be carried out, and routine checks on pollution levels in the affected areas and personnel should be performed. In the United States, where (without counting surgeons, dental staff, and vets, whose exposure is intermittent and variable) about 50 000 people, 35% of them women, are occupationally exposed to anaesthetic gases (anaesthetists, anaesthetists's nurses, operating theatre nurses, and technicians),³⁰⁵ hospital regulations limit pollution in operating theatres to 25 ppm of nitrous oxide and 1-2 ppm of halogenated agents.³⁰⁷

Drug addiction and psychic problems

The risk of health workers becoming addicted to drugs arises because of the stress and anxiety to which they are subjected at times and because of the ready availability of suitable drugs.³⁰⁸ It is quite common for attempts to be made to overcome the pressures of overwork, stress, and affective strain by excessive resort to stimulants, whose use may alternate with excessive consumption of coffee or tobacco and the use of sleeping pills, alcohol, or other drugs. Several studies have confirmed that alcoholism, drug addiction, and depression are more common among doctors than among the general population,³⁰⁹⁻³¹⁵ as are visits to the psychiatrist.³¹⁶ For example, of the 5925 patients admitted to the psychiatric wards of the Mayo Clinic between 1956 and 1963, 93 were doctors (one in every 64 admissions),³¹⁷ and in 59% of these cases the cause was either alcoholism (16.6%) or drug addiction (34.4%). Almost all these patients had used drink or drugs as a defence against acute anxiety. Similar findings were reported by Murray after examining admissions to psychiatric centres in Scotland³¹⁸: there were significantly higher than average rates of alcoholism, drug addiction, functional

neurotic depression, and affective psychosis among doctors. Perhaps because of the high prevalence of depression, suicides are two to three times more common among doctors—psychiatrists especially³¹⁸—than among the general population.^{313 319-321} Marital and sexual dissatisfaction is also more prevalent than among non-doctors,^{316 322 323} though the divorce rate is no higher than average.³²⁴⁻³²⁶

Work, which as a channel for the fulfilment of the individual may favour his or her psychic wellbeing, may also prove psychically deleterious owing to the individual's own traits (personality, social conflicts, etc), to the working environment (group relations), to the nature of the work itself (disagreeable or alienating jobs, overfast workplace, excessive workloads, changes of shift), or to promotion.

CHANGES OF SHIFT

Changes of shift originate personal, family, and social problems.³²⁷ The resulting lack of synchronisation between exogenous stimuli³²⁸ (the periodic variation of light, sound, heat, and atmospheric ionisation) and endogenous clocks^{329 330} gives rise to disorders due to dysrhythmia. In such cases the human organism, which is programmed for daytime activity, receives stimuli opposed to its physiological requirements, the consequence being a characteristic disorder featuring insomnia, irritability, dyspepsia, spacial and temporal disorientation, and eventual corticoadrenal exhaustion.³³¹ Night work goes against the genetic, professional, and social characteristics of the human species. Working while in a state of nocturnal "disactivation" requires double effort and hence produces greater fatigue than day work, and this fatigue is incompletely eliminated because daytime sleep is less restful than night sleep. This state of chronic fatigue is exemplified by the slowing down of motor responses to optical and acoustic stimuli and by decreased efficiency at work. Digestive dysrhythmia is also observed, and there is greater dependence on tobacco and alcohol among night workers than among day workers.

The family life of the night worker is likewise upset, since unless the hours kept by the rest of the family are made to fit in with his he himself must either fit in with theirs or live apart. Socially, night work results in greater insularity (nightworkers have fewer friends), and less participation in group activities (sports, union or political work, or cultural pursuits), though it is also true that morning and afternoon shifts likewise have their social disadvantages. Thus the morning shift allows the midday meal to be shared with the family and leaves time for other family and social activities, but since it begins so early may also prove fatiguing. The afternoon shift prevents social life, since contact with the children is limited and there are

no evenings out, but sleeping hours are normal and workers with alternating shifts regard it as allowing lost sleep to be made up. It is the night shift that causes most problems due to its consequences for family life and social activity.³³²

Emotional stress is one of the main health hazards for hospital personnel,³³³⁻³³⁵ especially for the student or young graduate nurse, and derives from contact with the dying, with seriously ill children, and with sorrowful, anxious, and suicidal patients. Working in intensive care units also gives rise to situations that, depending on the personality of the individual, may cause pathological psychic disorders.³³⁶⁻³³⁷ In this context the relation between doctor and patient is passive, with the doctor and nurse being under the constant pressure of having continually to take instant decisions that are critical for the patients' lives. This highly stressful combination of great responsibility and continuous attention to patients' needs, which is further exacerbated by a work pattern that does not allow relaxation during breaks and leisure periods, ends up by producing exhaustion and reduced operational capacity.³³⁶⁻³³⁸⁻³³⁹ Signs of stress among intensive care unit workers include psychosomatic disorders (mainly affecting the digestive system), anomalous social reactions, various types of neuroses and depression, and even dereliction of duty.³⁴⁰ Psychopathological stress likewise tends to increase when health work becomes monotonous, boring, and "meaningless" while retaining its demanding nature. In all these stressful circumstances, however, the effect on the health worker depends largely on his attitude to the situation and on his resources for coping with it.³³⁷⁻³⁴¹

Assault

Human life undoubtedly includes a degree of aggressive conduct, whether this be innate or acquired. The necessity of continual adaptation to a changing environment forces such conduct on us, especially in a society as competitive as ours. In this section, however, we shall limit our considerations to the urge to harm some person (or some object that represents or replaces that person), the harm caused being the sole aim of the aggressive act. Innate aggressive behaviour needs no stimulus, but there also exist socially acquired forms of aggression that are triggered by external stimuli such as frustration. The patient and his relatives should therefore never be frustrated, since they may become aggressive.³⁴²⁻³⁴⁵

Frustration may be produced when the behaviour of health workers is unfitting or is not what is expected, in short when they forget their role. Health care involves a social group comprising the health

worker, the patient, and the latter's relatives and friends. To paraphrase Berard and Gubler, sometimes it achieves its purpose, the curing of the disease; generally it manages to provide relief; and it always provides assuagement.³⁴⁶ Each member of the group has his own status and accompanies the execution of a particular function with a behaviour that is expected by the others. Each status and role in the group has its complement: husband and wife, father and son, health worker and patient. The roles of the health worker and the patient are defined by the ambient culture and certain aspects are even regulated by law.³⁴⁷ The role of the patient is characterised both by rights such as exemption from his responsibilities and the right to receive aid, and by obligations such as his obligation to wish for his own recovery and to cooperate in his treatment.³⁴⁸ The status and role of the health worker are characterised by impartiality (giving each patient the attention required by the gravity of his illness), by confidentiality (using the patient's body and the information he knows regarding the patient solely for the patient's cure), by affective neutrality (avoiding involvement in the patient's affective problems), by disinterested altruism, and by technical and scientific competence.⁴¹ Criticism of the health worker's behaviour will come from the patient's relatives and friends, the commonest criticism being that little time is devoted to Social Security patients.³⁴⁹ The patient should be convinced that a diagnosis has been attained or at least that everything possible is being done to attain it. The harmonious course of the relation between doctor and patient, each with his status and role, is fundamental for achieving the objective, health.³⁵⁰⁻³⁵² If disrupted the results may be pathogenic not only for the patient but, paradoxically, for the doctor also.³⁵³

For the doctor, the pathogenic consequences of a breakdown in his relation with the patient may be physical (injuries), anatomopathological (heart attacks), or functional or psychosomatic (ulcers). The sources of the patient's aggressive conduct may be classified in three categories: those arising in the context of his relation with the doctor, those arising in his own family and working environment, and a residual group of other social sources. Aggression may take the form of physical assault or verbal attacks and be received either by the health worker himself or by his family or possessions. Verbal assault is undoubtedly the most common but cases of physical attack are by no means rare and are occasionally reported by the daily newspapers.

When faced with patients known to be mentally ill, awareness of the possibility of apparently unmotivated fits of violence puts the doctor on his guard. The most dangerous are the paranoics,

especialmente los hipocóndricos convencidos que están sufriendo de una enfermedad grave porque el doctor los está tratando mal (hay usualmente otro trabajador que alienta a tales pacientes). Los erotómanos convencidos que sólo el doctor de su timidez evita declarar su amor por ellos generalmente limitarse a insultos verbales pero el doctor debe evitar estar solo con ellos. Los histéricos mentirosos que se inventan rumores y reportes de lo que creen que una vez ellos han expresado. Los excitados esquizofrénicos y los hipomaníacos también pueden ser agresivos (el último—usualmente verbalmente—al ser informado de la verdad sobre su condición, lo que a veces es más sabio mantenerlo a sí mismo). Los ataques de drogadictos y sadistas también pueden esperarse. Los ataques de pacientes sanos son mucho más peligrosos porque son inesperados; y son mucho más importantes porque su inesperabilidad significa que el doctor es inconsciente de cómo él ha frustrado al paciente.³⁵⁴

Finalmente, un tipo diferente de "ataque" al que los trabajadores de salud están expuestos es al ser demandados por mala práctica por sus pacientes.³⁵⁵⁻³⁵⁸ Aunque ninguno quisiera negar a los pacientes este derecho, debe ser señalado que el abuso generalizado de este derecho puede ser perjudicial para la atención médica al inducir a los trabajadores de salud a practicar la medicina a la defensiva, dando mayor importancia a lo que es legalmente seguro para el doctor que a lo que es beneficioso para el paciente.³⁵⁹ Paradójicamente, hay razones para pensar que el exceso de casos de mala práctica actualmente perturbando la paz de ánimo de los médicos en los Estados Unidos es en parte debido al éxito de la medicina moderna: el hecho de que el acto de curar es ahora un lugar común ha resultado en una tendencia a cualquier fracaso ser inmediatamente atribuido a negligencia del doctor en lugar de a las limitaciones intrínsecas de su arte.

References

The following are the most important of the references cited in the text. A complete list may be obtained from the author on request.

- 1 Gestal JJ. Occupational hazards in hospitals: risk of infection. *Br J Ind Med* 1987;44:435-42.
- 2 Freeman NT, Whitehead J. *Introduction to safety in the chemical laboratory*. London: Academic Press, 1982.
- 5 Link LJ. *Electrical safety in hospitals, a current review of principles and practices*. Eire, Pennsylvania: American Sterilizer Company, 1976.

- 30 Braestrup CB, Vikteriof KJ. *Manuel de radioprotection dans les hopitaux et en pratique generale*. Genève: Organisation Mondiale de la Santé, 1974:vois I, II, III.
- 34 Department of Health and Social Security. *Guidance on the safe use of lasers in medical practice*. London: HMSO, 1984.
- 37 Steward T. Problems caused by the continuous use of visual display units. *Lighting Research Technology* 1980;1:26-36.
- 44 Fisher AA. Contact dermatitis in medical and surgical personnel. In: Maitach HI, Gellin GA, eds. *Occupational and industrial dermatology*. London: Year Book Medical Publishers, 1982:219-28.
- 54 Organización Mundial de la Salud. Límites recomendados por razones de salud en la exposición profesional a determinados solventes orgánicos. *Ginebra: OMS Ser inf técn* 1982;664:9-43.
- 76 Organización Mundial de la Salud. Límites de exposición profesional que se recomiendan por razones de salud: sustancias irritantes de las vías respiratorias. *Ginebra: OMS Ser inf técn* 1984;707:39-76.
- 77 González Ferradas E. *Formaldehído: toxicología e impacto ambiental*. Madrid: Ed Mapfre, 1986.
- 78 Gibson JE. *Formaldehyde toxicity*. London: Hemisphere Publishing Corporation, McGraw-Hill International Book Company, 1983.
- 136 Acheson ED, Gardner MJ, Pannett B, Barnes HR, Osmond C, Taylor CP. Formaldehyde in the British chemical industry. *Lancet* 1984;ii:611-6.
- 143 Gennart JP, Dutrieux M, Lauwerys R. La toxicité de l'oxide d'éthylène: revue de la littérature. *Archives des Maladies Professionnelles de Médecine du Travail et de Sécurité Sociale* 1983;4:269-74.
- 189 *Riesgos y prevención en el manejo de soluciones citostáticas*. Málaga: Sociedad Española de Higiene y Medicina Preventiva Hospitalaria, 1985:2. (Serie documentos técnicos.)
- 197 Ferstanning LL. Trace concentrations of anesthetic gases: a critical review of their disease potential. *Anesth Analg* 1978;57:328-45.
- 198 Edling C. Anesthetic gases as an occupational hazard: a review. *Scand J Work Environ Health* 1980;6:85-93.
- 262 Tannenbaum TN, Goldberg RJ. Exposure to anesthetic gases and reproductive outcome: a review of the epidemiologic literature. *J Occup Med* 1985;9:659-68.
- 311 Modlin HC, Montes A. Narcotics addiction in physicians. *Am J Psychiatry* 1964;121:358-65.
- 313 A'Brook MF, Hailstone JD, McLaughlan IEJ. Psychiatric illness in the medical profession. *Br J Psychiatry* 1967;113:1013-23.
- 322 Scheiber SC. Emotional problems of physicians: I nature and extent of problems. *Ariz Med* 1977;34:323-5.
- 324 Fine C. *Married to medicine: an intimate portrait of doctors' wives*. New York: Atheneum, 1981.
- 332 Leonard R. *Le travail de nuit et les horaires alternants*. Paris: Université de Paris, 1982.
- 335 McCue JD. The effects of stress on physicians and their medical practice. *N Engl J Med* 1982;8:458-63.
- 343 Carthy JD, Ebling FJ. *Historia natural de la agresión*. México: Siglo XXI editores, 1966.
- 358 Ataz L. *Los médicos y la responsabilidad civil*. Madrid: Editorial Montecorvo, 1985.