

Asbestos related health hazards among power plant workers

Y Lerman, A Finkelstein, Y Levo, M Tupilsky, Mimi Baratz, A Solomon, G Sackstein

Studies of asbestos insulation workers have shown a higher than expected death rate due to malignant mesothelioma.¹ Although asbestos has been widely used in power plants to provide thermal insulation for steam pipes and turbines, asbestos related health effects are rarely described among power plant workers² (and J Bonnell *et al*, XVIII International Conference on Occupational Health, Brighton, 1975). We report two cases of malignant pleural mesothelioma in a clerk and an insulator worker of an electric power plant.

Case report

A 72 year old man was admitted because of left pleuritic chest pain, non-productive cough, dyspnoea, and progressive weakness. Past and family history were non-contributory. Family history was positive for bone cancer and breast cancer in two siblings. The patient was a retired employee of a power plant where he had worked for 31 years as a clerk until seven years before admission. Before his employment in the power plant he had been a construction worker for five years and a manager in the wood industry for nine years. No history of exposure to asbestos other than in the power plant was obtained.

On examination the patient was malnourished and in mild pulmonary distress. There was no chest wall tenderness or palpable masses. On percussion, the left lung field was dull and breath sounds were appreciably decreased. A chest radiograph showed a left pleural thickening with effusion, the right hemithorax was normal. A pleural tap yielded 800 cc of haemorrhagic exudative fluid. Histological examination of the fluid showed many lymphocytes, histiocytes with few mesothelial cells.

Computed tomography of the chest showed that the left hemithorax was of small volume compared to the right. There was circumferential crenated pleural encasement affecting the left lung and a left basal

pleural effusion was present. The attenuation numbers of the irregularly thickened pleura were considerably higher than the accompanying free pleural fluid. The suggested diagnosis was of extensive pleural mesothelioma. The presence of a calcified pleural plaque on the right posterior pleural surface indicated exposure to asbestos in the remote past (fig 1).

A pleural biopsy showed a thickened fibrous pleura partially covered by hyperplastic mesothelial cells and fibrin. Within the fibrous stroma were bundles of spindle cells, some with enlarged hyperchromatic nuclei. These findings were interpreted as consistent with cellular fibrous mesothelioma (fig 2). The patient died seven months later from respiratory failure.

Case report 2

A 54 year old man was admitted because of recurrent pleural effusion. He had been an insulator worker for 31 years and had worked in the same power plant as the first patient. He had no history of exposure to asbestos. On examination, the patient was well nourished and not in pulmonary distress. Expansion of the left chest was decreased. The left lung was dull to percussion and breath sounds were decreased over the same area. A chest radiograph showed a massive left pleural effusion. Pleuroscopy showed the




Figure 1 Computed tomography of chest showing encasement of left lung with thick pleural mass accompanied by pleural effusion. Calcified pleural plaque is seen on right posterior surface of pleura.

Israel Defence Forces, Medical Corps, Ichilov Medical Centre, Tel Aviv, Israel
Y Lerman, A Finkelstein, Y Levo, M Tupilsky, M Baratz, A Solomon
Israel Electric Company, Medical Division, Tel Aviv
G Sackstein




Figure 2 Bundles of spindle cells within fibrous stroma. (Haematoxylin & eosin $\times 100$).

presence of 1500 cc of greenish exudative fluid and a gelatinous mass covering the lower third of the visceral pleura, diaphragm, and pericardium. Biopsy of the pleural mass showed malignant epithelial mesothelioma with many Schauman bodies. The patient received intrapleural treatment with nitrogen mustard, thiotepa, and yttrium (Itrium) and intravenous treatment with adriamycin. An abdominal tap of ascitic fluid four months later showed malignant mesothelial cells. He died 12 months later from respiratory failure.

Discussion

Thermal power stations produce electric energy by generators. The generators are complex units consisting of two principal units, the boiler and the turbine. The heat generated by the turbines and steam pipes is one of several health hazards in power stations,³ and it is essential to provide thermal insulation.

Asbestos has been widely used for this purpose.⁴ Measurements of asbestos in air inside and outside power stations show low fibre counts but higher concentrations are found in storerooms where asbestos is deposited and handled.² The asbestos related health risk in power plants was also confirmed by the presence of ferruginous bodies in the sputum of power station workers.²

Although asbestos related health effects have been well described among insulation workers,¹ only rarely have such effects been described among power

station workers² (and J Bonnell, 1975). Bonnell reported on asbestos related disease among 77 insulation workers of an electric power station in London; eight cases of mesothelioma were described. Another survey of 55 full time power station workers in France showed that the risk of exposure to asbestos was not confined to workers engaged in lagging operations: ferruginous bodies and asbestos related abnormalities, mainly pleural thickening and calcification, were also present among workers in other job categories.² In a recent review Cammarano *et al* reported on site specific mortality among thermo-electric power plant workers in Italy.⁵ Eighteen patients with neoplasms were observed whereas only 8.32 were expected. There were five cases of lung cancer (expected 2.83) but no cases of mesothelioma.

The nature of exposure of the first patient reported here is a low level long term exposure type. Several authors have suggested that even minimal exposure, such as experienced by maintenance workers ("by-stander" exposure) or by neighbourhood or household members may induce malignant mesothelioma.^{6,7} The risk of asbestosis after this type of exposure is minimal.¹ Under such conditions of low level exposure to asbestos cases of mesothelioma serve as an index of the potential health hazards. The cases reported here emphasise the need for population based studies among power plant workers to evaluate the risk accurately.

- 1 Selikoff IJ, Lee DHK. *Asbestos and disease*. New York: Academic Press, 1978.
- 2 Hirsch A, Di Menza L, Carre A. Asbestos risk among full-time workers in an electricity-generating power station. *Ann NY Acad Sci* 1979;330:137-45.
- 3 Parmeggiani L, ed. *Encyclopaedia of occupational health and safety*. Vol 2. 3rd ed. Geneva: International Labour Office, 1983:1778-80.
- 4 Burger H. *Asbestos fundamentals. Origin, properties, mining processing, utilisations*. New York: Chemical Publishing Company, 1963.
- 5 Cammarano G, Crosignani P, Bernino F, *et al*. Additional follow-up of cancer mortality among workers in a thermo-electric power plant. *Scand J Work Environ Health* 1986;12:631-2.
- 6 Epler GR, Gerald MXP, Gansler EA, *et al*. Asbestos related disease from household exposure. *Respiration* 1980;39:229-40.
- 7 Chen W, Mottel NK. Malignant mesothelioma with minimal asbestos exposure. *Human Pathol* 1978;9:253-8.

Accepted 12 June 1989