Lung dust content in idiopathic pulmonary fibrosis: a study with scanning electron microscopy and energy dispersive x ray analysis

E Monsó, J M Tura, J Pujadas, F Morell, J Ruiz, J Morera

Abstract

Examination with an optical microscope and polarised light is not sensitive enough to detect low diameter asbestos fibres. This limitation implies that some cases of asbestosis can be erroneously diagnosed as idiopathic pulmonary fibrosis (IPF) if asbestos bodies are not found in the standard examination of abnormal tissue. To determine whether IPF is overdiagnosed, a study was carried out with scanning electron microscopy (SEM) and energy dispersive x ray analysis (EDXA) on 25 samples previously diagnosed as IPF at the standard examination. Scanning electron microscopy will show the presence of low diameter fibres in the lung without tissue destruction, and these fibres can be identified using EDXA. The quantitative and qualitative results for lung tissue from patients diagnosed as having IPF were compared with the results of the examination of 25 samples of normal lung. Most of the samples from patients diagnosed as having IPF showed only occasional inorganic particles (<10 particles/SEM field at 160 $\times$), results equivalent to the results obtained in normal lung. Two cases of IPF, however, showed innumerable asbestos fibres (>100 fibres/SEM field). One of these two patients had an antecedent of brief exposure to asbestos. No environmental antecedent was found in the second patient. Asbestosis was the final diagnosis for these two patients. The examination of inorganic particles in normal lungs showed mainly non-fibrous silicates (61.4%) and particles of heavy elements (34.9%). Only one asbestos fibre was found (0.9%). It is concluded that standard pathological techniques overdiagnose IPF in a few cases in which asbestos bodies are not found with the optical microscope.

A lung sample showing pulmonary fibrosis and inorganic particles allows the diagnosis of pneumoconiosis. Absence of inorganic particles in a fibrotic lung sample of a patient with diffuse interstitial lung disease, if no other cause is determined, usually gives a diagnosis of idiopathic pulmonary fibrosis (IPF). It is well known, however, that examination with an optical microscope and polarised light is not sensitive enough to detect low diameter inorganic particles, especially asbestos fibres. For lung samples from patients previously diagnosed by optical microscopy as having IPF, an examination using more sensitive techniques could find that some of the samples contain enough asbestos fibres to diagnose asbestosis.

Advances in mineralogical analysis have enabled its application in pneumology during the last two decades.¹⁻³ Among the techniques applicable to organic matter, scanning electron microscopy (SEM) with energy dispersive x ray analysis (EDXA) is especially useful.^{4,5} Inorganic particles in the lung are easily identified if secondary and backscattered imaging are used (spot 0.125 μm ; maximum resolution power 150 $\text{\AA}$). The atomic composition of every inorganic particle visualised can be identified using EDXA. If the area studied in all the samples is predetermined, a quantitative and qualitative examination can be done.

We studied 25 samples with a previous pathological diagnosis of IPF and 25 samples of normal lung, and determined the number of inorganic particles seen in the area visualised with SEM at 160 $\times$ (278 300 μm^2), identifying their atomic composition with EDXA.

Material and methods

Twenty five histological samples of lung tissue from patients (mean age (SD) 58.3 (13.7); 14 men, 12 women) with a previous pathological diagnosis of IPF were studied. All samples were obtained by surgical biopsy, mainly from the lingula. Twenty two had a pathological diagnosis of usual interstitial pneumonia (UIP), two had UIP with a desquamative component, and one patient had giant cell interstitial pneumonia. Exposure history indicated that nine patients might have inhaled inorganic particles (table

Serveis de Pneumologia, Hospital Germans Trias i Pujol (Badalona) i Hospital Vall d'Hebron (Barcelona). ITQT Consell Superior d'Investigacions Científiques (Barcelona), Catalonia, Spain
E Monsó, J M Tura, J Pujadas, F Morell, J Ruiz, J Morera

Table 1 *Inhalatory antecedents in patients with idiopathic pulmonary fibrosis*

Case No.	Sex	Age (y)	Antecedent*
1	M	39	No antecedent
2	W	53	No antecedent
3	M	71	Quarry 20 years
4	M	55	Cement <one year
5	W	54	No antecedent
6	W	70	No antecedent
7	M	58	No antecedent
8	W	67	Husband asbestos worker
9	W	78	No antecedent
10	M	49	Chemical industry > five years
11	W	50	Rubber factory > five years
12	W	23	No antecedent
13	M	75	No antecedent
14	W	52	No antecedent
15	M	66	No antecedent
16	M	44	No antecedent
17	M	65	No antecedent
18	M	72	No antecedent
19	M	54	Quarry > five years
20	W	31	Chemicals < one year
21	M	70	Asbestos three years
22	W	71	No antecedent
23	M	72	No antecedent
24	W	63	No antecedent
25	M	56	Asbestos 20 years

*Only antecedents that could imply inhalation of inorganic particles, even in small amounts were considered.

M = Man; W = woman.

1). Pulmonary biopsy was indicated in these patients due to a brief exposition period or because diagnoses other than pneumoconiosis were considered, which could not be confirmed without an examination of pulmonary tissue. In these patients absence of inorganic particles in the examination by optical microscopy and polarised light precluded a diagnosis of pneumoconiosis.

Twenty five samples of normal lung were also examined from patients (mean age (SD) 62.8 (14.7); 12 men, 13 women) who had died, mainly of cardiovascular causes. Patients with a history of inhalation of inorganic particles were excluded by personal history and phone call to relatives. The samples were all obtained from peripheral areas of the lung.

Samples from cases of IPF were set in paraffin and samples of normal lung were kept in formalin. Thick sections were obtained from all samples; 20 μ m from those in paraffin and 1–2 mm from those in formalin. Paraffin samples were deparaffined in two baths of xylol. All samples were dried by the critical temperature method (Polaron E300) after substitution of the tissue water with acetone by immersion in baths of increasing concentration. For the SEM examination and mineralogical analysis the samples were placed on pure carbon stubs. We ruled out contamination of the stubs and paraffin with silica or silicates.

The SEM examination (Phillips SEM 500) was carried out first at low magnification (160 $\times$) looking for a representative area of the pulmonary parenchyma and avoiding the peribronchial, perivascular,

and subpleural areas. The area selected (magnification 160 $\times$ (278 300 μ m²) was examined for inorganic particles increasing the magnification to 5000 $\times$ to identify low diameter fibres. The visualised inorganic particles were quantified and the atomic composition of each was determined using EDXA (working conditions: 25 kV; spot 0.125 μ m; detector stub distance 32.5 mm; x ray angle 20 $^\circ$), a spectrometric technique that detects all elements with an atomic number above 10.^{6,7}

The morphology and atomic composition of the particles visualised with the SEM and analysed with EDXA allows us to classify the inorganic particles as silica (detection of a peak of Si), non-fibrous silicates (Si with Mg, K, Al, Ca, or Fe), asbestos (length: width ratio > 3 $\leftarrow$ fibrous silicate with Mg or amphibole atomic composition),^{8,9} and particles of heavy elements (Fe, Pb, Ti, Ag, Sn, or Ba). The particles of elements with a possible organic origin (Ca) were not considered.

Table 2 *Inorganic particles: idiopathic pulmonary fibrosis*

Patient	No of particles	Type
1	0	—
2	1	Silicate (1)
3	0	—
4	0	—
5	6	Silicate (6)
6	7	Silicate (5) Silica (1) Fe (1)
7	7	Silicate (5) Silica (1) Al (1)
8	5	Silicate (3) Silica (1) ZnCu (1)
9	6	Silicate (2) Silica (2) Fe (1) Al (1)
10	4	Silicate (2) ZnCu (2)
11	0	—
12	5	Silicate (5)
13	0	—
14	6	Silicate (3) Ba (1) Co (2)
15	3	Silicate (1) Silica (2)
16	0	—
17	1	Silicate (1)
18	1	Silicate (1)
19	2	Silicate (1) Fe (1)
20	10	Silicate (4) Cu (6)
21	Innumerable	Fibrous silicate Fe (1)
22	Innumerable	Fibrous silicate
23	5	Silicate (5)
24	—	—
25	0	—

Figure 1 Asbestos fibres in a sample with a previous diagnosis of idiopathic pulmonary fibrosis (scanning electron microscopy 1250 $\times$).

Results

One sample of normal lung tissue and one sample of tissue from a patient with IPF were discarded from mineralogical analysis due to poor quality of the SEM imaging that impeded recognition of the inorganic particles. These samples were not considered in the results.

Most of the samples from patients diagnosed as having IPF contained only occasional inorganic particles (< 10 particles in the area studied), but two (8.3%) showed innumerable asbestos fibres (> 100 asbestos fibres in the area). One of these patients had an antecedent of a brief occupational exposure to asbestos. No relevant antecedent was found in the second patient (table 2; figures 1, 2).

The twenty four samples of normal lung analysed

showed only occasional inorganic particles in 18/24 cases (< 10 particles). The analysis of the atomic composition of the detected particles showed a high prevalence of non-fibrous silicates (67/109 particles; 61.4%) and particles of heavy elements (38/109 particles; 34.9%). One case contained 27 inorganic particles in the area studied, and one of these particles was the only asbestos fibre found in the normal group (1/109 particles; 0.9%). This patient, together with all others in the normal group, had no relevant inhalatory antecedents (table 3).

Discussion

Our results provide evidence that an examination using optical microscopy and polarised light over-diagnoses IPF. We found 2/24 cases with a previous diagnosis of IPF (8.3%) that contained innumerable asbestos fibres in the area studied, with no asbestos

Table 3 Inorganic particles in normal lung types

Patient	No of Particles	Type
26	7	Silicate (4) Fe (3)
27	6	Silicate (3) Fe (3)
28	6	Silicate (1) Silica (2) Fe (2) ZnCu (1)
29	5	Silicate (4) Silica (1)
30	5	Silicate (4) ZnCu (1)
31	27	Silicate (11) Asbestos (1) Pb (15)
32	3	Silicate (1) Al (1) Ag (1)
33	0	—
34	8	Silicate (8)
35	0	—
36	4	Silicate (4)
37	4	Silicate (4)
38	5	Silicate (2) Fe (3)
39	3	Silicate (3)
40	2	Silicate (2)
41	0	—
42	0	—
43	5	Silicate (5)
44	9	Silicate (7) Fe (2)
45	1	Silicate (1)
46	7	Silicate (3) Al (1) Cu (1) Fe (2)
47	1	Fe (1)
48	0	—
49	1	Fe (1)
50	—	—

Figure 2 Energy dispersive x ray analysis of an asbestos fibre.

bodies. The final diagnosis for these cases must be asbestosis.¹⁰

It has been previously suggested that IPF, or more probably a subgroup of it, could in fact be pneumoconiosis not diagnosed by standard techniques,^{5 11-18} Until now this opinion has been poorly founded on original work with no control group.¹⁵ Most of the asbestos fibres are below the 0.2 µm resolution of the optical microscope,¹⁹ and will not be detected with a conventional examination.²⁰ Asbestos bodies, however, are usually easily identified and allow the diagnosis of asbestosis (two or more asbestos bodies in a pulmonary fibrosis lung sample).²¹ There exists a relation between the number of asbestos bodies and the number of asbestos fibres in the lung,^{22 23} but this relation is extremely variable.²⁴⁻²⁶ It is clear, however, that the lung content of asbestos fibres is always much higher than the lung content of asbestos bodies.^{24 27} This situation allows that a patient with asbestosis, sampled in an area that does not show asbestos bodies, was erroneously diagnosed as having IPF after examination with optical microscopy and polarised light.^{28 29} A more sensitive analysis will correct the diagnosis.

The SEM examination of a lung sample in cases of IPF combined with the determination of the atomic composition of the inorganic particles using EDXA is easier than a mineralogical analysis using destructive techniques³⁰⁻³² and can be used as a complementary study in cases of a questionable diagnosis. Moreover, surface SEM examination offers the advantage that it can be used on samples obtained by lung biopsy that cannot be used for mineralogical analysis because there is not enough tissue available (such as our IPF lung samples).

Inorganic particles in normal lungs have not been previously studied in the western Mediterranean population. Our study shows that the particles found are mainly non-fibrous silicates (61.4%), and, rarely, silica (2.75%). We also found a high proportion of particles of heavy elements (34.9%).

It is well known that particles of silica and non-fibrous silicates can be found in normal lungs.³³ Berry *et al* studied 40 normal lungs and found Fe, Ti, and Al silicates with amorphous structure.¹¹ Brambrilla *et al*, examining the lungs of animals of San Diego zoo, found 95% of silicates, 5% of silica, and 5% of particles of heavy elements. The silicates were identified by x ray diffraction as mica.³⁴

Our results allow us to conclude that standard techniques overdiagnose IPF. The use of more sensitive, non-destructive techniques such as SEM and EDXA in diagnosis of IPF could show that some of the lung samples with a previous diagnosis of IPF are really asbestosis.

The study was supported by Institut d'Estudis Catalans, SEPAR and FiSss 90/0431.

Requests for reprints to: Dr Eduard Monsó, Servei de Pneumologia, Hospital Germans Trias i Pujol, Ap correus 72, 08916 Badalona, Catalonia, Spain.

- DeVuyst P, Jedwab J. L'analyse minéralogique en pathologie respiratoire. *Rev Med Brux* 1981;2:115-9.
- DeVuyst P, Dumortier P, Rickaert F, Van de Weyer R, Lenclud C, Yernault JC. Occupational lung fibrosis in an aluminium polisher. *Eur J Respir Dis* 1986;68:131-40.
- DeVuyst P, Van de Weyer R, DeCoster A, *et al*. Dental technician's pneumoconiosis. A report of two cases. *Am Rev Respir Dis* 1986;133:316-20.
- Morgenroth K, Blaschke R, Schlake W. Energy dispersive X ray analysis of semi-thin section in the scanning transmission. *Beiträge zur Pathologie* 1973;150:406-11.
- Monsó E, Tura JM, Marsal M, Morell F, Pujadas J, Morera J. Mineralogical microanalysis of idiopathic pulmonary fibrosis. *Arch Environ Health* 1990;45:185-9.
- Ghadially FN. The technique and scope of electron-probe x-ray analysis in pathology. *Pathology* 1979;11:95-110.
- Davis JMG, Gylseth B, Morgan A. Assessment of mineral fibres from human lung tissue. *Thorax* 1986;41:167-75.
- Mossman BT, Gee JBL. Asbestos-related diseases. *New Engl J Med* 1989;320:1721-30.
- Murphy RL, Becklake MR, Brooks SM, *et al*. The diagnosis of non-malignant diseases related to asbestos. *Am Rev Respir Dis* 1986;134:363-8.
- Churg A. Nonneoplastic diseases caused by asbestos. In: Churg A, Green FHY, eds. Pathology of occupational lung disease. New York: Igaku-Shoin, 1988:213-78.
- Berry JP, Henoc P, Galle P, Pariente R. Pulmonary mineral dust. A study of ninety patients by electron microscopy, electron microanalysis and electron microdiffraction. *Am J Pathol* 1976;83:427-38.
- Rüttner JR, Spycher MA, Sticher H. The detection of etiologic agents in interstitial pulmonary fibrosis. *Hum Pathol* 1973;4:497-512.
- Miller A, Langer AM, Teirstein AS, Selikoff IJ. Nonspecific interstitial pulmonary fibrosis. Association with asbestos fibers detected by electron microscopy. *N Engl J Med* 1975;292:91-3.
- Pariente R, Berry JP, Galle P, Cayrol E, Brouet G. A study of pulmonary dust deposits using the electron microscope in conjunction with the electron sound analyser. *Thorax* 1972;27:80-2.
- Abraham JL, Hertzberg MA. Inorganic particulates associated with desquamative interstitial pneumonia. *Chest* 1981;80:67-70.
- Abraham JL, Spragg RG. Documentation of environmental exposure using open biopsy, transbronchial biopsy and bronchoalveolar lavage in giant cell interstitial pneumonia (GIP). *Am Rev Respir Dis* 1979;119:197.
- Davison AG, Haslam PL, Corrin B, *et al*. Interstitial lung disease and asthma in hard metal workers: bronchoalveolar lavage, ultrastructural, and analytical findings and results of bronchial provocation tests. *Thorax* 1983;38:119-28.
- Stettler LE, Groth DH, Mackay GR. Identification of stainless steel welding fume particulates in human lung and environmental samples using electron probe microanalysis. *Am Ind Hyg Assoc J* 1977;38:76-86.
- Churg A. Non asbestos pulmonary mineral fibers in the general population. *Environ Res* 1983;31:189-200.
- Dodson RF, Hurst GA, Williams MG, Corn C, Greenberg SD. Comparison of light and electron microscopy for defining occupational asbestos exposure in transbronchial lung biopsies. *Chest* 1988;94:366-70.
- Craighead J, Abraham J, Churg A, *et al*. Pathology standards for the diagnosis of asbestos related diseases. *Arch Pathol Lab Med* 1982;106:543-97.
- Whitwell F, Scott J, Grimshaw M. Relationship between occupations and asbestos fibre content of the lungs in patients with pleural mesothelioma, lung cancer and other diseases. *Thorax* 1977;32:377-86.
- Ashcroft T, Heppleston AG. The optical and electron microscopic determination of pulmonary asbestos fibre concentration, and its relation to the human pathological reaction. *J Clin Pathol* 1973;26:224-34.
- Morgan A, Holmes A. Concentrations and dimensions of coated and uncoated asbestos fibres in the human lung. *Br J Ind Med* 1980;37:25-32.

- 25 Kobayashi H, Watanabe H, Zhang WM, Ohnishi Y. A quantitative and histological study of pulmonary effects of asbestos exposure in general autopsied lungs. *Acta Pathol Jpn* 1986;**36**:1781-91.
- 26 Case BW, Sebastien P. Environmental and occupational exposures to chrysotile asbestos: a comparative microanalytic study. *Arch Environ Health* 1987;**42**:185-91.
- 27 Pooley FD, Ranson DL. Comparison of the results of asbestos fibre dust counts in lung tissue obtained by analytical electron microscopy and light microscopy. *J Clin Pathol* 1986;**39**:313-7.
- 28 Dodson RF, Williams Jr MG, O'Sullivan MF, Corn CJ, Greenberg SD, Hurst GA. A comparison of the ferruginous body and uncoated fiber content in the lungs of former asbestos workers. *Am Rev Respir Dis* 1985;**132**:143-7.
- 29 Churg A. The diagnosis of asbestosis. *Hum Pathol* 1989;**20**:97-9.
- 30 DeNee PB. Identification and analysis of particles in biological tissue using SEM and related techniques. In: O'Hare AMF, Johari O, Becker RP, eds. *SEM/1976/III. SEM Inc.* IL60666. Chicago: ITT Research Institute IL60666, 1976:461-8.
- 31 Vallyathan NV, Green FHY, Craighead JE. Recent advances in the study of mineral pneumoconiosis. *Pathol Annu* 1980;**15**:77-104.
- 32 Henderson WJ, Melville-Jones C, Wilson DW, Griffiths K. Oxygen incineration and electron microscope x-ray microanalysis of mineral particles in biological tissues. *J Histochem Cytochem* 1978;**26**:1087-93.
- 33 Ziskind M, Jones RN, Weill H. Silicosis. *Am Rev Respir Dis* 1976;**113**:643-65.
- 34 Brambilla C, Abraham J, Brambilla E, Bemrschke K, Bloor C. Comparative pathology of silicate pneumoconiosis. *Am J Pathol* 1979;**96**:149-70.

Accepted 12 November 1990

Destruction of manuscripts

From 1 July 1985 articles submitted for publication will not be returned. Authors whose papers are rejected will be advised of the decision and the manuscripts will be kept under security for three months to deal with any inquiries and then destroyed.