PostScript

LETTERS

Occupational exposure of midwives to nitrous oxide on delivery suites

In our opinion, the article "Occupational exposure of midwives to nitrous oxide on delivery suites"¹ is in need of some remarks.

In the paper a serious problem seems to be the presence of nitrous oxide in samples collected at the beginning of the shift.

Many years ago, when N₂O in urine was first evaluated, we frequently observed "uncommon" concentration of N₂O in urine of exposed and unexposed subjects. The phenomenon was kept under control and disappeared when urine samples were treated with a small quantity of H₂SO₄ (0.2 ml). For this reason, we suggested the following:² "... Approximately 10 ml of urine were collected from all the subjects at the end of the exposure period in 120 ml gastight glass vials with airtight plugs. Caps were rapidly replaced in the vials to prevent any significant loss of N₂O. The vials contained 0.2 ml sulfuric acid in order to avoid the in vitro production of N₂O (probably due to microflora activity).³ ...".

Another point we consider very important is that the subjects must void the bladder rapidly in areas known to be free of nitrous oxide, otherwise a significant contamination of samples can occur.

In conclusion, we think that among the simple precautions that should be taken to avoid significant errors (avoiding collection of urine samples in places contaminated with N₂O, carrying out collection rapidly, and using airtight collection vials in order to avoid any major loss of dissolved anaesthetic), one point should be emphasised in view of its importance: storage of urine before analysis can produce an endogenous formation of N₂O originating from the oxidation processes of the nitrogen compounds present in biological liquids. Experiments performed to study this phenomenon have shown that the process is inhibited if the urine is kept acid. If, as a precaution, a few drops of strong acid are added to each collection vial before urine samples are collected, neoformation of nitrous oxide will be avoided and the urine samples may then be stored as long as required prior to the analysis.

M Imbriani

Università degli Studi di Pavia Dipartimento di Medicina Preventiva, Occupazionale e di Comunità, Pavia, Italy

S Ghittori, I Maestri

Labs-Meia, Fondazione Salvatore Maugeri IRCCS, via Ferrata, 8, Pavia, Italy

Correspondence to: Dr S Ghittori; sghittori@fsm.it

doi: 10.1136/oem.2003.012534

References

- 1 Henderson KA, Matthews IP, Adisesh A, *et al.* Occupational exposure of midwives to nitrous oxide on delivery suites. *Occup Environ Med* 2003;60:958-61.
- 2 Imbriani M, Ghittori S, Pezzagno G, *et al.* Nitrous oxide in urine as biological index of exposure in operating room personnel. *Appl Ind Hyg* 1988;3:223-7.
- 3 World Health Organisation (WHO). *Health hazards from nitrates in drinking water* Environmental Health 1. Report on a WHO meeting. Copenhagen, 1985:33-4.

Author's reply

Professor Imbriani and colleagues report experiments which showed that endogenous formation of N₂O was inhibited if urine is kept acid. The convenience of adding 0.2 ml of sulphuric acid to vials recommends its routine use in practice and we do not disagree with this recommendation.

The likelihood that the pre-shift urine measurements which we reported arise from this phenomenon rather than other factors should be judged in the light of the following considerations:

- All pre-shift urine samples were collected in areas free of nitrous oxide.
- The period between sample collection and deposit in a freezer was approximately the same for each sample. Despite this 24 midwives had zero N₂O in their pre-shift samples and 22 had non-zero values, of whom 12 had very high values.
- The period between deposit in a freezer and analysis varied between samples but biological activity should not occur in the freezer.

I Matthews

Department of Epidemiology, Statistics and Public Health, University of Wales College of Medicine, Health Park, Cardiff CF14 4XN, UK; matthewsip@cf.ac.uk

The evidence for workplace counselling is in Medline

Henderson *et al* point out the increasing approval of counselling as an effective intervention to treat or prevent the effects of stress at work by British judges, although they could use expert advice on this matter.¹ In reaction to this development, they pose the rhetorical question: where to find evidence on the effectiveness of counselling. In stead of answering this question they grasp the opportunity to criticise the report of the British Association for Counselling.² I totally agree with their criticism of the report. It is of low quality and does not provide reliable evidence on the effectiveness of counselling. However, I was surprised by the fact that the authors did not present reliable evidence that does exist on the topic. The question cannot be left unanswered. We gave an answer to an almost similar question in our article on evidence based medicine. We showed the feasibility of searching for evidence in Medline for practitioners of occupational

health.³ We elaborated an example of a teacher with symptoms of burnout who wanted to know the best treatment for his condition. Our search resulted in at least one good review and one meta-analysis.^{4,5} The meta-analysis by van der Klink *et al* firmly concludes: "stress management interventions are effective and cognitive-behavioural interventions are more effective than the other intervention types". This is in line with the earlier findings of the review by Murphy that we found as well.

From the authors' editorial it can be inferred that they favour interventions such as a reduction of working hours or increasing staff numbers, more than counselling. This does sound sympathetic to me as well and it is in line with the principle of hierarchy of controls, which states that primary prevention is to be preferred to, for example, personal protective equipment.⁶ However, in our case, there is not much evidence that supports such an approach. This is partly due to a lack of studies in the area of organisational interventions. The organisational intervention studies that have been done, however, do not yield a significant effect size.⁵ On the other hand, there seems to be enough evidence to conclude that cognitive behavioural interventions are effective in counterbalancing the effects of stress at work. So, even when only reliable evidence is used, there is still much to support counselling in the sense of cognitive behavioural treatment. In addition, there is a systematic review in the Cochrane Library on counselling in primary care, which concludes that it is associated with a modest improvement in short term outcome compared to "usual care" and not associated with more costs.⁷ Based on this evidence I would not simply reject counselling as ineffective.

This case illustrates that, in occupational health in general, there is a lack of awareness of the existence of evidence on effective interventions. That is the main reason why we are in the process of developing an Occupational Health Field within the Cochrane Collaboration. The Cochrane Collaboration is an international organisation, dedicated to making up-to-date, accurate information about the effects of healthcare readily available worldwide. Have a look at www.cochrane.org for more details.

We hope that, in the near future, the Occupational Health Field will fulfil its promises and will simplify the finding of evidence on occupational health interventions like counselling.

J Verbeek

Coronel Institute for Occupational and Environmental Health, University of Amsterdam, Occupational Health Field, Finnish Institute of Occupational Health, Kuopio, Finland; j.h.verbeek@amc.uva.nl

doi: 10.1136/oem.2003.012138

References

- 1 Henderson M, Hotopf M, Wessely S. Workplace counselling. *Occup Environ Med* 2003;60:899-900.

- 2 McLeod J. *Counselling in the workplace: the facts* Rugby. British Association for Counselling and Psychotherapy, 2001.
- 3 Verbeek JH, van Dijk FJ, Malmivaara A, et al. Evidence-based medicine for occupational health. *Scand J Work Environ Health* 2002;28:197-204.
- 4 Murphy LR. Stress management in work settings: a critical review of the health effects. *Am J Health Promot* 1996;11:112-35.
- 5 van der Klink JJ, Blonk RW, Schene AH, et al. The benefits of interventions for work-related stress. *Am J Public Health* 2001;91:270-6.
- 6 Roelofs CR, Barbeau EM, Ellenbecker MJ, et al. Prevention strategies in industrial hygiene: a critical literature review. *AIHA J* 2003;64:62-7.
- 7 Bower P, Rowland N, Mellor C, et al. Effectiveness and cost effectiveness of counselling in primary care. *Cochrane Database Syst Rev* 2002;CD001025.

Comments on article by Koh and Aw

Quoting both dictionary definitions and statutory requirements, Koh and Aw's education article¹ limits the definition of occupational "health surveillance" to the detection of adverse health effects resulting from occupational exposures. In doing so, they exclude international and national requirements for occupational health and medical surveillance to assess fitness for work.

Looking at the hazard of ionising radiation, international recommendations,² European Directives,³ and UK National Legislation⁴ all identify a requirement for surveillance where the primary purpose is an assessment of the individual's fitness for post. Similarly, in considering surveillance of divers, a key element of requirements is an assessment of fitness for work. On a more general level, both in the public and in the occupational setting, systems of health surveillance exist for drivers where it is clearly nonsense to suggest that this is aimed at the detection of adverse effects resulting from time behind the wheel. It is therefore suggested that the authors' conclusion needs to be expanded to identify a requirement for periodic examination of individuals, not only to detect reversible ill health, but also to assess fitness for work.

C Kalman

Salus Occupational Health & Safety, Centrum Park,
Hagmill Road, Coatbridge ML5 4TD, UK;
Chris.Kalman@lht.scot.nhs.uk

doi: 10.1136/oem.2003.011551

References

- 1 Koh D, Aw T-C. Surveillance in occupational health. *Occup Environ Med* 2003;60:705-10.
- 2 ICRP. *Recommendations of the International Commission on Radiation Protection*. Publication 60, 1992.
- 3 European Union. Basic safety standards for the health protection of the general public and workers against the dangers of ionising radiation. European Union Council Directive 1996/29/EURATOM, 13 May 1996.
- 4 The Ionising Radiations Regulations 1999 (UK statutory instrument).

BOOK REVIEW

Tolley's managing stress in the workplace

Carole Spiers (£60.00). Croydon, UK: LexisNexis UK. ISBN 0-7545-1269-X

"Not another book about workplace stress"—emanating in this case, from the "stress industry" would be an understandable reaction. Carole Spiers, the author, unequivocally describes herself as an "occupational stress consultant" and head of the Carole Spiers Group: "International Corporate Well-being Consultants".

She faces up to the implications immediately by asking "Why indeed another book about stress? What makes it different from the others?" Well, this one is intended to be practical and user friendly—a handbook that can sit on your shelf and act as a reference manual to be dipped into whenever required. It is aimed primarily at employers, employees, and their representatives rather than occupational health practitioners or academics; this is not a criticism—many occupational health practitioners will appreciate the way in which the subject of work related stress is assiduously presented in all its complexity.

Far from being all about the practicalities of managing stress in the workplace, there are chapters which go into some detail about the nature of stress, current legislation, and the health and safety framework in the UK and, to some extent, Europe. Naturally there has to be constant reference to health and safety and employment law but also to civil litigation, and here comes one of the problems: very few cases of work induced stress have in fact been litigated and those that have, have not, in many people's view, been very typical. Moreover, this is a fast changing field and the useful synopsis of appeal cases heard in 2002 may soon be out of date on account of impending House of Lords judgements. In another domain, namely identifying current workplace stressors, the template used: Culture, Demands, Control, Relationships, Change, Role, and Support has already been refined by the Health and Safety Executive as more is learned about measuring psychosocial factors at work.

However, the approach taken by the author to understanding the problem is straightforward, accepted by most, practical, and will be useful for people who want to systematise their approach to identifying workplace stressors or measure the effects of stress on an organisation.

There are chapters on bullying, post-traumatic stress, stress and health, and the effects of stress on the individual, whose conclusions will not be accepted by all. There is, unsurprisingly, a chapter on personal stress management strategies which sweeps up much of the advice, good and bad, offered by the self help industry and, to balance it, a chapter on healthy organisations. There is a short chapter on the future of stress.

In 400 pages the author covers most of what there is to know about the wider world of stress and has usefully interwoven a number of relevant themes. I was surprised how little mention was made of the medicalisation of stress—after all most employers receive their first intimation of an employee's stressed state by means by means of a sickness absence certificate signed by a general practitioner. This issue is only cursorily examined in chapter 10. I also failed to recognise many of the examples of stressed individuals which populate the book. They are all real cases, but where are the people with relatively undemanding jobs, beset by social problems, domestic difficulties, and unhealthy habits referred by harassed middle managers? It is often a toss up to know who will "go off with stress" first. I am not sure that this book is very enlightening about how to manage those people and how to prevent the seemingly inevitable slide of such individuals into resentment, long term sickness absence, and, eventually, Incapacity Benefit.

There does, also, seem to be an emphasis on larger organisations and not much about the dynamics within small and medium sized enterprises (where most people work these days), which are different.

The book does, however, deserve to be "dipped into" because there is a wealth of descriptive material on which to build.

D Snashall

NOTICE

28th ICOH International Congress on Occupational Health

The 28th ICOH International Congress on Occupational Health will be held in Milan, Italy, 11-16 June 2006.

Further information:

www.ich2006.it

Tel: +39 0250320110; fax: +39 025032011

Email: sabrina.braiati@unimi.it

CORRECTION

doi: 10.1136/oem.2002.006239corr1

With reference to the paper "Risk of selected birth defects by maternal residence close to power lines during pregnancy" (Blaasaas KG, Tynes T, Lie RT. *Occup Environ Med* 2004;61:174-6), the authors state:

"The total number of births inside the specified corridor given as 128 680 in the Results was wrong. We verified, however, that only 42 223 pregnancies were completed on specific addresses inside the corridor. These 42 223 births represented the cohort from which we identified the 465 cases and selected 930 controls. This should have been specified in the paper. The error gave a wrong impression of the prevalence of defects but had no implications for the results of the paper."