



# Impact of organisational change on mental health: a systematic review

Simon Grandjean Bamberger,<sup>1</sup> Anker Lund Vinding,<sup>2</sup> Anelia Larsen,<sup>3</sup> Peter Nielsen,<sup>4</sup> Kirsten Fonager,<sup>5</sup> René Nesgaard Nielsen,<sup>6</sup> Pia Ryom,<sup>1</sup> Øyvind Omland<sup>1</sup>

<sup>1</sup>Department of Occupational Medicine, Aalborg Hospital, Aarhus University Hospital, Danish Ramazzini Center, Aalborg, Denmark

<sup>2</sup>Department of Quality, North Denmark Region, Aalborg, Denmark

<sup>3</sup>Outpatient Department, Aalborg Psychiatric Hospital, Aalborg, Denmark

<sup>4</sup>Department of Political Science, Aalborg University, Aalborg, Denmark

<sup>5</sup>Department of Social Medicine, Aalborg Hospital, Aalborg, Denmark

<sup>6</sup>Department of Business and Management, Aalborg University, Aalborg, Denmark

## Correspondence to

Simon Grandjean Bamberger, Department of Occupational Medicine, Aalborg Hospital, Aarhus University Hospital, Havrevangen 1, Aalborg DK-9000, Denmark; sigb@rn.dk

Accepted 4 March 2012

Published Online First

27 April 2012

## ABSTRACT

Although limited evidence is available, organisational change is often cited as the cause of mental health problems. This paper provides an overview of the current literature regarding the impact of organisational change on mental health. A systematic search in PUBMED, PsychInfo and Web of Knowledge combining MeSH search terms for exposure and outcome. The criterion for inclusion was original data on exposure to organisational change with mental health problems as outcome. Both cross-sectional and longitudinal studies were included. We found in 11 out of 17 studies, an association between organisational change and elevated risk of mental health problems was observed, with a less provident association in the longitudinal studies. Based on the current research, this review cannot provide sufficient evidence of an association between organisational change and elevated risk of mental health problems. More studies of long-term effects are required including relevant analyses of confounders.

## INTRODUCTION

The last decade has brought increased attention to the possible detrimental effects of work on mental health. Existing research has focused on work strain and occupational health and found consistent evidence for associations.<sup>1</sup> Organisational change is often cited as a harmful exposure but few studies have been published to support this assumption.<sup>2–6</sup> One of the first studies on health risks was a study from Finland<sup>7</sup> where they found that the risk of health problems was at least two times greater after major downsizing than after no downsizing. A longitudinal study published few years later found a significantly faster decline in self-rated health even 4 years after downsizing among employees who had experienced major downsizing.<sup>8</sup> The increase in health problems was partially explained by concomitant increases in physical demands and job insecurity and a reduction in job control.<sup>8–9</sup> Besides, downsizing and repeated exposure to rapid personnel expansion may predict long-term sickness absence and hospital admissions.<sup>10</sup> Although these studies did not specifically address mental health problems, they should be interpreted within the framework of work stress.

Employees are increasingly confronted with frequent minor daily stressors related to changes in technology and workplace practices as well as the major upheavals of mergers, downsizing and

restructuring.<sup>11–12</sup> The imminence, duration and temporal uncertainty surrounding events of change can have a negative impact on employees.<sup>13</sup> Indeed, the increased uncertainty regarding job future or the direction of organisational change has been suggested to be a principal cause of stress.<sup>14–15</sup> Others propose that organisational change acts as a stressor through the individual's negative appraisal of the changes.<sup>16</sup>

The well-documented risks that might follow organisational changes are: intensification of job strain, time pressure, reduction of social support, lack of control and role ambiguity, which all have been associated with mental health problems.<sup>17–19</sup> Two systematic reviews of psychosocial factors at work and depression found evidence of a relation between perceived psychosocial job strains and an elevated risk of depressive symptoms or major depressive episode.<sup>20–21</sup> Job insecurity has also been consistently linked with detrimental mental health effects in both meta-analysis and reviews.<sup>22–23</sup> Another potential factor affected by organisational change: job dissatisfaction has shown strong associations with depression and anxiety according to meta-analysis.<sup>24</sup> Examining organisational change as a potential work stressor yields a certain benefit because organisational change is more tangible in nature than for instance change in meaningfulness of work is to the individual. Empirically it is possible to determine whether or not a change has happened and whether or not the employees are feeling worse (or better) hereafter.

The health effects of workplace reorganisation in intervention studies have previously been reviewed by Egan *et al*<sup>25</sup> and Bambra *et al*<sup>26</sup> who found small evidence for beneficial effects of increased employee participation and control. However, the reorganisation that occurs in intervention studies is typically aimed at addressing unhealthy environments, which is different from the economic motives of, for example, company downsizing.

Therefore, the objective of this paper is to provide a systematic review of observational studies on associations between organisational change and employee mental health problems.

## METHODS

The review was conducted in accordance with the PRISMA statement.<sup>27</sup>

### Eligibility criteria

#### Participants

The study population of interest was individuals employed in companies undergoing some sort of organisational change. Relevant comparators were

either employees not exposed to change or employees exposed to different levels or types of changes.

### Exposure

Our definition of organisational change was inspired by Rafferty's definition of *transformational* change and refers to modifications made to the core systems of an organisation including traditional ways of working, values, structure and strategy. Transformational changes can either be changes to the normal operational procedures (tasks, working hours) or changes in the system itself (eg, downsizing, changing the organisation's hierarchy or incorporating a new organisational system).<sup>28</sup> Only studies with organisational change explicitly listed as the primary exposure were considered.

### Outcome

This review was limited to studies examining mental health problems, specifically with a focus on depression, anxiety and stress. All health variables had to be measured by validated scale or based on diagnosis made by a mental health professional. Conversely studies where the outcome measured was readiness for change, commitment to change, coping with change or job satisfaction after change were not included.

### Study design

We focused on observational studies and in the case of duplicate publications of data from the same study, the paper with the most relevant analysis and risk estimate was chosen.

### Information sources

A series of computerised librarian-assisted systematic searches were undertaken in PubMed, PsycInfo, Excerpta Medica Database (EMBASE), EBSCO Business Source Complete and Web of Knowledge using a broad search strategy to find peer-reviewed articles of relevance. The searches were carried out using MeSH terms (medical subject headings) or free text when no MeSH term was available, and limited to English journal articles with abstracts. To find the most recent articles not yet indexed by MeSH terms, free text words were included for the period 2010–search date. The searches were performed in October 2011. Organisational change was operationalised in the search strategy as 'organizational change', 'organizational innovation', 'organizational restructuring', 'organizational transition', 'work change', 'downsizing' and 'reorganization'. The search terms for mental health problems were 'anxiety', 'depression', 'mental disorders', 'mental health', 'stress', 'occupational stress' and 'psychological stress'. Search terms were derived from keywords and titles of known core papers.

### Study selection

Initial eligibility assessment of titles was performed by the first author (SGB) under careful supervision of senior researcher ØO. Both ØO and SGB screened articles by abstract. Difficult judgements on relevance among reviewers were resolved by consensus.

### Data collection process

Data were extracted systematically from each study following a standardised format. Data were extracted by SGB under supervision of ØO. Information was extracted on: (1) Participants: (including nationality, sample size, type of organisations, work type, controls); (2) Exposure (type of organisational change); (3) Type of outcome measure (mental health problems using validated scale or other type of diagnostic tool); and (4)

Study aim and design, follow-up time, statistical analysis used, adjustments, bias and conclusion/results.


## RESULTS

The search strategy yielded 5443 records, which were screened for relevance based on title by one of the authors (SGB) and 5146 records were excluded. A duplicate search was performed and further 88 articles were removed. At step two the remaining 209 records were screened by abstract by SGB and ØO, and 156 records were removed. The reasons for exclusion was missing or ill-defined mental health outcome (57 studies), theoretical articles (22 studies), lack of organisational change as exposure (20 studies), intervention studies (17 studies), coping studies (17 studies), duplicate publications of data from same study (five studies), managers as study population (four studies) and otherwise irrelevant study design (14 studies).

At step three 53 full text articles were assessed for eligibility and 17 articles were included in the final synthesis. The reasons for exclusions were missing or ill-defined mental health outcome (22 studies), missing or ill-defined organisational change as exposure (12 studies), and duplicate publications of data from same study (two studies). The selection process is depicted in figure 1.

Descriptive statistics of the 17 eligible studies included in the review are listed in table 1. Six studies used a cross-sectional design<sup>29–34</sup> and 11 were longitudinal.<sup>2 3 35–43</sup> All but three studies<sup>2 32 43</sup> were published within the last 10 years. The included studies involved 40 993 participants, excluding the population and registry studies<sup>30 35 37</sup> which accounted for a total of 142 110 individuals. Besides the three large studies of the general working population, hospital staff and civil servants were the most frequent study subjects (table 1).

Data were primarily collected by questionnaire, though two studies were strictly registry based,<sup>35 37</sup> three studies used structured interviews<sup>29 30 41</sup> and finally one study<sup>36</sup> collected blood samples. The studies had a mean baseline response rate of 64%, with three studies below 50%<sup>3 39 43</sup> and one longitudinal study<sup>3</sup> was even below 20%; however, this was partly redeemed by a high retention rate. The median follow-up time of the longitudinal studies was approximately 2 years (ranging from 1 to 7). Four studies assessed participants twice<sup>36 39–41</sup> and four studies<sup>2 3 38 43</sup> assessed them thrice, while one study assessed



**Figure 1** Flowchart of the number of articles in different stages of the selection strategy.

**Table 1** Studies of organisational change and mental health problems

Cross-sectional studies						
Author, year	Sampling frame	Participation	Statistical analysis	Adjusted/controlled for	Exposure	Outcome (p<0.05%)
Bourbonnais, 2005 <sup>29</sup>	Nurses (n=2002)	77.1% (76% and 84%)	Binomial regression	Coping strategies, social support, domestic load, type A behaviour, personal and socio-occupational characteristics (age, familial status, seniority in the institution, job status, work shift)	Restructuring	Exposed PR 1.84 (1.56–2.16) of psychological distress vs controls
Dragano, 2005 <sup>30</sup>	Population (n=22 559)	61%	Bivariate analyses	Age, east/west residency, education, occupational status, physical demands and occupational hazards, weekly working hours and job insecurity	Downsizing	OR 1.53 (1.30–1.79) for exposed men of increased symptom load. OR 1.71 (1.43–2.06) for exposed women
Greubel, 2011 <sup>31</sup>	Police employees (n=1523)	76%	ANOVA	Age, gender and shift work	Relocation, extensive changes: downsizing and job changes	Anxiety ↑ depression ↑
Karasek, 1990 <sup>32</sup>	White collar (n=8504)	87%	Mantel-Haenszel	Age, sex	Company initiated job changes	Depression ↑
Pepper, 2003 <sup>33</sup>	State employees (n=5889)	55%	Hierarchical linear modelling	Nothing	Downsizing rate	Mental health component ↔ Perceived Stress ↔
Verhaeghe, 2006 <sup>34</sup>	Hospitals (n=2094)	51%	Logistic regression	Age, sex	Situational changes in working environment	Distress ↑
Longitudinal studies						
Author, year	Sampling frame	Participation (follow-up time)	Statistical analysis	Adjusted/controlled for	Exposure	Outcome (p<0.05%)
Dahl, 2011 <sup>35</sup>	Population (n=92 869)	Registry based study (6 years)	Multivariate analysis with logit models	Age, gender, children (3 age groups), marital status, stress of parents and spouse, firm tenure, wage, occupation level, firm size, firm age and industry classification	Organisational change	Stress ↑
Ferrie, 1998 <sup>2</sup>	White collar (n=7419)	73% (~3 years)	ANOVA and logistic regression	Age and employment grade	Change in job description, increase in workload and pace of work	OR 1.56 (1.30–1.86) of minor psychiatric morbidity for exposed men. For women ↔
Hansson, 2008 <sup>36</sup>	Hospital (n=226)	74% (1 year)	One-way and two-way ANOVA	Nothing	Reorganisation of work systems	Biological stress markers ↔
Kivimaki, 2007 <sup>37</sup>	Municipal employees (n=26 682)	Registry based study (7 years)	Negative binomial regression	Sex, age, education, occupational status and local government	Downsizing	RR 1.49 (1.10–2.02) of psychotropic drug use for exposed men. RR 1.12 (1.00–1.27) for exposed women
Loretto, 2010 <sup>3</sup>	Hospital (n=5385)	Baseline 18.4%, 84.3% retention rate of 1st cohort, 76.7 of 2nd cohort, (~1 year)	Logistic regression	Past GHQ caseness, personal and biographic factors, objective workplace and job characteristics	Perceived amount of overall change	OR 1.21 (1.06–1.38) of GHQ caseness when exposed to change
Moore, 2006 <sup>38</sup>	Manufacturing company (n=460)	62%–74% (~2.5 years)	ANCOVA	Age, gender, education and marital status	Downsizing	Depression ↔
Netterstrom, 2010 <sup>39</sup>	Civil servants (n=685)	44% (2 years)	Logistic regression	Age, leadership, department and occupation	Merger	Depression ↔
Probst 2003 <sup>40</sup>	State agency employees (n=313)	63% (6 months)	Multivariate ANOVA	Nothing	Restructuring	Mental health index ↔
Rohall, 2001 <sup>41</sup>	Military officers (n=1536)	85% (~2 years)	ANOVA	Nothing	Downsizing	Anxiety ↑ depression ↑
Vaananen 2011 <sup>42</sup>	Forest industry corporation (n=6511/4096)	82.3% (~4 years)	Cox proportional hazard models	Sex, age, marital status, occupational status, sense of coherence and job characteristics	Merger	Increased risk of postmerger psychiatric event HR 1.60 (1.19–2.14)
Woodward 1999 <sup>43</sup>	Hospital employees (n=346)	47% (~2 year)	ANOVA	Nothing	Re-engineering	Anxiety ↑ depression ↑

Where applicable ratios (HR, OR, RR and PR) and 95% CIs were available in the studies, these are listed.

Otherwise relevant significant results are reported using errors: ↓ = decrease in symptoms; ↑ = increase in symptoms; ↔ = no significant change in symptoms. ANOVA, analysis of covariance; GHQ, General Health Questionnaire; PR, prevalence ratio.

participants a total of four times.<sup>42</sup> The two remaining longitudinal studies were registry based and contained annual information from a 6-year<sup>35</sup> and 7-year period.<sup>37</sup>

Various types of organisational change were assessed in the 17 studies ranging from major changes like downsizing, mergers and reorganisations to lesser (though perhaps not less intrusive) job content changes. Downsizing as the best described exposure was analysed in two cross-sectional studies<sup>30 35</sup> and three longitudinal studies.<sup>37 38 41</sup> The three longitudinal studies<sup>37 38 41</sup> examined different degrees of contact with downsizing over time, that is, exposure–response relationships. One of the cross-sectional studies<sup>35</sup> used downsizing ratio as the exposure that is also a type of exposure–response study. The last downsizing study<sup>30</sup> simply examined the exposure–effect relationship of downsizing on mental health. Two longitudinal studies examine the effect of company mergers<sup>39 42</sup> while two other longitudinal studies<sup>3 35</sup> examined the effect of multiple types of organisational changes both quantitatively and qualitatively. Restructuring was assessed in one cross-sectional study<sup>29</sup> and three longitudinal studies.<sup>36 40 43</sup> Finally, job changes were examined by one longitudinal study<sup>2</sup> and three cross-sectional.<sup>31 32 34</sup>

Only one study<sup>42</sup> used clinical psychiatric criteria (International Classification of Disease) to define the mental health outcome. The remaining studies primarily relied on self-rated questionnaires or registry information to determine the mental health effects. Depression<sup>31 32 38 39 41 43</sup> and anxiety<sup>31 41 43</sup> were the most specific mental health problems reported; however, all the studies utilised different measurement scales. One study<sup>37</sup> used the ambiguous ‘psychotropic drug use’ as mental health indicator. Finally, a large part of the studies relied on the vague term ‘distress’<sup>29 33–36</sup> or general measures of self-rated mental health<sup>2 3 30 33 40</sup> as indicators of mental unbalance. Due to the disparity of the outcome data extracted from the studies and the heterogeneity of the outcome we chose not to perform a meta-analysis or any other type of data synthesis.

Most of the studies<sup>2 3 29–32 34 35 37–39 42</sup> adjusted results for socio-demographic factors such as age, gender, marital status and education. Work-related factors such as employment status and different job characteristics were less commonly adjusted for.<sup>2 3 29–31 35 37 39 42</sup> Only three studies adjusted for personality traits like type A behaviour,<sup>29</sup> neuroticism<sup>3</sup> or sense of coherence.<sup>42</sup> Though there is evidence of exacerbated risk of reoccurrence of mental health problems like depression,<sup>44</sup> only three studies<sup>3 40 42</sup> analysed for effects of previous mental health status. Five studies<sup>33 36 40 41 43</sup> did not adjust results for any possible confounders. Almost all of the studies lacked a non-respondents analysis and for the longitudinal studies analysis of attrition.

Associations between organisational change and mental health problems were found in five<sup>29–32 34</sup> of the six cross-sectional studies. The remaining study<sup>35</sup> found no association between downsizing rate and mental health effects, but did report lower mental health scores than national norm for all participating study sites. The longitudinal studies presented mixed results as only six out the 11 studies<sup>3 35 37 41–43</sup> found associations between exposure to organisational change and subsequent mental health problems, and one study<sup>2</sup> only found significant effects for men.

Three<sup>30 37 41</sup> out of five of the studies on downsizing found an association between mental health problems and exposure, and of these two<sup>37 41</sup> studies were longitudinal. In two<sup>29 43</sup> out of four studies on restructuring an effect on mental health was observed, and of these studies one was longitudinal.<sup>43</sup> The two negative studies<sup>36 40</sup> were both longitudinal. The effects of

mergers on mental health have been analysed in two studies.<sup>39 42</sup> Netterström *et al*<sup>39</sup> did not find an association between mergers and depression while Väänänen *et al*<sup>42</sup> found an association between the exposure and postmerger psychiatric events. Concerning the studies on exposure to multiple types of organisational changes both studies<sup>3 35</sup> found an association with negative health effects. All four studies on job changes<sup>2 31 32 34</sup> found an association between the exposure and mental health problems; however, in one study this association was only evident for men.

## DISCUSSION

Despite the broad search profile we found few studies which analysed the association between organisational change and mental health. A total of 17 studies were found eligible for this review, surprisingly few considering the high number of records initially evaluated by title, a finding experienced by others.<sup>2 3</sup> An association between exposure to organisational change and decreased mental health was found in 11 out of the 17 studies with a less provident association in the longitudinal studies. This might be explained by an observed acute short-term effect on mental health during the actual change process that later disappears or normalises over time such as the results by Dahl suggest.<sup>35</sup> Another reason for the discrepancy could be that the mental health impaired has already left work, or have not the energy to participate in such a study. None of the longitudinal studies provide drop-out analyses to counter this argument; however, three studies<sup>3 40 42</sup> at least take previous mental health status into account. The prospective cohort study by Kivimäki *et al*<sup>37</sup> included all employees who were at work before downsizing. The use of psychotropic drugs was examined in employees who kept their jobs after major downsizing and among those who lost their jobs during the downsizing, compared with the employees who did not experience downsizing. This strategy specifically allowed examination of the role of drop-out in the results.

An important aspect all but three of the studies<sup>33 34 42</sup> omit is the employees’ individual perception of change. The individual categorisation of a specific organisational change as threatening or not is connected to the psychological reaction. This process can be influenced by several factors like coping strategies,<sup>45</sup> negative affectivity,<sup>46</sup> stress prior to change,<sup>47</sup> perceived social support<sup>48</sup> or length of employment.<sup>49</sup> Personal characteristics such as personality type, temperament, intelligence and genetic constitution may all affect the way in which an individual understands and reacts to life experiences. Mental health problems can take years to develop but the studies explore a snapshot of a particular time period.<sup>50</sup> The effect of these confounding factors may influence the estimate. Kivimäki *et al*<sup>37</sup> also notes that survival bias may prevent detection of the adverse effects of work, as major psychiatric disorders are commonly related to work disability and, furthermore, a potential selective factor for unemployment.

Recall bias may influence the results as employees who experienced organisational change as traumatic might be more inclined to remember the experience than employees who were less affected. In relation to this, utilising self-reported data like Verhaeghe *et al*<sup>34</sup> introduces the risk of circular reasoning;<sup>51</sup> those who rate the organisational change as stressful are the stressed employees. In a broader perspective this approach addresses the issue whether organisational change really is harmful in itself, or if merely the employees’ individual perception of the change makes it harmful. This type of bias is avoided in population studies where the exposure (typically downsizing)

is determined by registry data. These types of studies, however, provide little information on the actual process of implementing the organisational change. Research into change processes suggests that the readiness for organisational change and the subsequent healthiness of the change process may reduce the experience of stress and increase the employees' abilities to cope with the changes,<sup>45 52 53</sup> that is, mediate or even moderate the change effect. Due to the number of different workplaces included in the population studies no specific information on the change process is reported. Furthermore, there is a risk of misclassification when organisational change is assessed at workplace level, as the particular change may have affected only parts of the organisation. Therefore, it would be preferable to use an independent measure of exposure in these studies.

Additional risk of misclassification is related to the possible effect of preceding organisational changes. This could skew the results in either way making the employees more vulnerable or resilient to mental health problems.<sup>38</sup> Pepper *et al*<sup>33</sup> noted that downsizing (as well as merger) is usually a change existing in a complex network of events. This would add to the likelihood of concomitant organisational changes such as restructuring or contract changes making it harder to measure the effect of downsizing as an independent exposure and increasing the risk of misclassification. Another important aspect emphasised by Dahl<sup>35</sup> is the rationale behind companies' change in the first place. He hypothesises that employees might be experiencing increased stress because their firms are poor performers trying to regain momentum by implementing organisational change. Thus, the mental health problems are related to the companies' poor performance and not the organisational change itself. If job insecurity is a central component in the pathway between organisational change and mental health problems as suggested,<sup>14 15 23</sup> then organisational changes in branches or industries with high competition might be at further risk. Indeed, the results of Dahl<sup>35</sup> suggest that performance variables influence the effect of change on stress; however, this needs to be further examined.

### Limitations

The epidemiologic evidence this review provides is limited mainly due to the low number of published relevant studies. Second, the definition of organisational change is at best vague including both major and minor changes to the organisation and work environment. To some extent, we compare the effect of major downsizing<sup>30 33 37 38 41</sup> and work schedule rearrangement<sup>2 32 34</sup> as being equal.

Subsequently, the definition of mental health as outcome is also marred by confinements. Limiting mental health to minor psychiatric disorders and not just mental discomfort narrows the available material considerably. Numerous studies otherwise relevant use job security, job satisfaction and morale as psychological outcome which can be relevant in other contexts but hardly describe mental health on a psychiatric level.<sup>54–61</sup> The most optimal mental health outcome would be diagnosis by psychiatrist; however, we had to settle for registry data, self-reported data or proxies like psychotropic drug use and stress hormone levels.

Several of the studies were cross-sectional leaving out any causality from the conclusions. However, it could be argued that it is probable that poor psychosocial work environment could be a reason to implement job changes and doubtful that the employee's mental health status is the cause of the major changes like downsizing and mergers.<sup>31</sup>

In three<sup>3 36 38</sup> of the included studies there is a high risk of confounding. The study by Hansson *et al*<sup>36</sup> identifies hormone

markers of mental distress, but fails to define valid baseline hormone levels. The analysis does not control for individual differences in cortisol awakening response, seasonal and diurnal variation, and lifestyle factors such as intake of coffee and alcohol, smoking and physical exercise.<sup>62</sup> Moore *et al*<sup>38</sup> comment in their discussion that due to small cell sizes, the results should only be regarded as exploratory, which is a conclusion we support. The study by Loretto *et al*,<sup>3</sup> otherwise solid, has a low response rate (18.4%) and no analysis of non-responders. The results might, therefore, be heavily biased. However, for the majority of the studies<sup>2 29 31 32 35 37 39 41–43</sup> we have no reason to believe that the results were confounded in such a degree that the associations would be altered significantly. Taken this into account and restricting the analyses to 14 studies with no major confounding errors, 10 studies<sup>29–32 34 35 37 41–43</sup> report a positive association between organisational changes and mental health, one study<sup>2</sup> reported an association for men but not women and finally three studies reported no association.<sup>33 39 40</sup> However, this restriction would alter the effect on the estimate increasing the positive association from ~65% including all 17 studies to ~71% when limited to the 14 studies. Both numbers are significantly lower than the ratio of 85% positive studies that Quinlan and Bohle<sup>23</sup> reported in their 2009 review on health and safety effects of downsizing and job insecurity. However, Quinlan and Bohle<sup>23</sup> included a wider array of (softer) outcomes and considered job insecurity an exposure in itself, and consequently their ratio is not directly comparable with the ratio found in this study.

### CONCLUSION

The exposure variables are multi-dimensional, and have been described, evaluated or measured differently. Multiple study designs have been used and the size of the population analysed has varied substantially. The outcomes in the analyses are inconsistent and similar outcomes have been defined differently among the studies. In 11 out of 17 studies an association between organisational change and elevated risk of mental health problems was observed, with a less provident association in the longitudinal studies. Given the heterogeneity of the studies, and the large number of cross-sectional studies this

#### What is already known on this subject

- ▶ The psychosocial work environment affects the mental health of employees.
- ▶ Perceived psychosocial job strains elevate the risk of depressive symptoms, anxiety and work stress.
- ▶ Organisational change is related to temporal increase of job insecurity.

#### What this paper adds

- ▶ The first systematic review of the association between organisational change and employee mental health problems.
- ▶ No convincing evidence of an association between these was found; however, studies are scarce and results should be considered with care.

review cannot provide convincing evidence of an association between organisational change and elevated risk of mental health problems. The studies are scarce and the findings should be considered with great care. Further analysing is needed to determine if the observed effect is of temporary or of more persistent nature. More studies with relevant analyses of confounders are recommended.

**Acknowledgements** We wish to thank Hjørdis Rasmussen who helped with the literature search.

**Contributors** SGB and ØO were responsible for the conduction of the study, interpretation of the data, study analysis including figures and tables and writing the report. Literature search was performed by SGB, ØO and Hjørdis Rasmussen. SGB is the guarantor. All members of the GOPA group (Globalisation, organisational change and psychosocial environment) participated in revising of the draft and all approved the final version for publication. All the authors are members of the GOPA group and ALV is the project leader.

**Funding** This research was financed by grant: 20080053113/12-2008-09 from the Foundation for Research of Work Environment (Denmark). The funders played no part in the conduct or reporting of the research.

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

1. Stansfeld S, Candy B. Psychosocial work environment and mental health—a meta-analytic review. *Scand J Work Environ Health* 2006;**32**:443–62.
2. Ferrie JE, Shipley MJ, Marmot MG, et al. The health effects of major organisational change and job insecurity. *Soc Sci Med* 1998;**46**:243–54.
3. Loreto W, Platt S, Popham F. Workplace change and employee mental health: results from a longitudinal study. *Br J Manage* 2010;**21**:526–40.
4. Corrigan J, Plagerson S, Lund C, et al. Global trade and mental health. *Glob Soc Policy* 2008;**8**:335–58.
5. Wilpert B. Impact of globalization on human work. *Saf Sci* 2009;**47**:727–32.
6. National Institute for Occupational Safety and Health. *The changing organization of work and the safety and health of working people*, Columbia Parkway Cincinnati: NIOSH Publications Dissemination 4676, 2002:2002–116. Publication No 2002-116 OH-45226-1998.
7. Vahtera J, Kivimäki M, Pentti J. Effect of organisational downsizing on health of employees. *Lancet* 1997;**350**:1124–8.
8. Kivimäki M, Vahtera J, Pentti J, et al. Downsizing, changes in work, and self-rated health of employees: a 7-year 3-wave panel study. *Anxiety, stress & coping. Int J* 2001;**14**:59–73.
9. Kivimäki M, Vahtera J, Pentti J, et al. Factors underlying the effect of organisational downsizing on health of employees: longitudinal cohort study. *BMJ* 2000;**320**:971–5.
10. Westerstam H, Ferrie J, Hagberg J, et al. Workplace expansion, long-term sickness absence, and hospital admission. *Lancet* 2004;**363**:1193–7.
11. Sikora PB, Beaty ED, Forward J. Updating theory on organizational stress: the asynchronous multiple overlapping change (AMOC) model of workplace stress. *Hum Resource Dev Rev* 2004;**3**:3–35.
12. Di Nunzio D, Hohnen P, Hasle P, et al. Impact of restructuring on health and safety and quality of work life. *Psychosocial Risks* 2009;**1**:90. CIT3-CT-2005–006193 Leuven, Belgium, ASBN 978-90-8836-017-6.
13. Rafferty AE, Restubog SLD. The impact of change process and context on change reactions and turnover during a merger. *J Manag* 2010;**36**:1309–38.
14. Bordia P, Hunt E, Paulsen N, et al. Uncertainty during organizational change: is it all about control? *Eur J Work Organizational Psychol* 2004;**13**:345.
15. Pollard T. Changes in mental well-being, blood pressure and total cholesterol levels during workplace reorganization: the impact of uncertainty. *Work Stress* 2001;**15**:14–28.
16. Pahkin K, Väinänen A, Koskinen A, et al. Organizational change and employees' mental health: the protective role of sense of coherence. *J Occup Environ Med* 2011;**53**:118–23.
17. LavoieTremblay M, Bonin J, Lesage AD, et al. Contribution of the psychosocial work environment to psychological distress among health care professionals before and during a major organizational change. *Health Care Manag* 2010;**29**:293–304.
18. Lindorff M. Predictors of symptom increase among managers—the role of environmental, personal and situational characteristics. *Stress Med* 1995;**11**:41–50.
19. Nelson A, Cooper CL. Uncertainty amidst change: the impact of privatization on employee job satisfaction and well-being. *J Occup Organizational Psychol* 1995;**68**:57–71.
20. Bonde JP. Psychosocial factors at work and risk of depression: a systematic review of the epidemiological evidence. *Occup Environ Med* 2008;**65**:438–65.
21. Netterstrom B, Conrad N, Bech P, et al. The relation between work-related psychosocial factors and the development of depression. *Epidemiol Rev* 2008;**30**:118–32.
22. Sverke M, Hellgren J, Näswall K. No security: a meta-analysis and review of job insecurity and its consequences. *J Occup Health Psychol* 2002;**7**:242–64.
23. Quinlan M, Bohle P. Overstretched and unreciprocated commitment: reviewing research on the occupational health and safety effects of downsizing and job insecurity. *Int J Health Serv* 2009;**39**:1–44.
24. Faragher EB, Cass M, Cooper CL. The relationship between job satisfaction and health: a meta-analysis. *Occup Environ Med* 2005;**62**:105–12.
25. Egan M, Bamba C, Thomas S, et al. The psychosocial and health effects of workplace reorganisation. 1. A systematic review of organisational-level interventions that aim to increase employee control. *J Epidemiol Community Health* 2007;**61**:945–54.
26. Bamba C, Egan M, Thomas S, et al. The psychosocial and health effects of workplace reorganisation. 2. A systematic review of task restructuring interventions. *J Epidemiol Community Health* 2007;**61**:1028–37.
27. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting item for systematic reviews and meta-analyses; the PRISMA-statement. *BMJ* 2009;**339**:332.
28. Bartunek JM, Moch MK. First-order, second-order, and third-order change and organization development interventions: a cognitive approach. *J Appl Behav Sci* 1987;**23**:483.
29. Bourbonnais R, Brisson C, Malenfant R, et al. Health care restructuring, work environment, and health of nurses. *Am J Ind Med* 2005;**47**:54–64.
30. Dragano N, Verde PE, Siegrist J. Organisational downsizing and work stress: testing synergistic health effects in employed men and women. *J Epidemiol Community Health* 2005;**59**:694–9.
31. Greubel J, Kecklund G. The impact of organizational changes on work stress, sleep, recovery and health. *Ind Health* 2011;**49**:353–64.
32. Karasek R. Lower health risk with increased job control among white collar workers. *J Organ Behav* 1990;**11**:171–85.
33. Pepper L, Messinger M, Weinberg J, et al. Downsizing and health at the United States department of energy. *Am J Ind Med* 2003;**44**:481–91.
34. Verhaeghe R, Vlerick P, Gemmel P, et al. Impact of recurrent changes in the work environment on nurses' psychological well-being and sickness absence. *J Adv Nurs* 2006;**56**:646–56.
35. Dahl MS. Organizational change and employee stress. *Manag Sci* 2011;**57**:240–56.
36. Hansson A, Vingard E, Arnetz BB, et al. Organizational change, health, and sick leave among health care employees: a longitudinal study measuring stress markers, individual, and work site factors. *Work Stress* 2008;**22**:69–80.
37. Kivimäki M, Honkonen T, Wahlbeck K, et al. Organisational downsizing and increased use of psychotropic drugs among employees who remain in employment. *J Epidemiol Community Health* 2007;**61**:154–8.
38. Moore S, Grunberg L, Greenberg E. Surviving repeated waves of organizational downsizing: the recency, duration, and order effects associated with different forms of layoff contact. *Anxiety Stress And Coping* 2006;**19**:309–29.
39. Netterstrom B, Blond M, Nielsen M, et al. Development of depressive symptoms and depression during organizational change—a two-year follow-up study of civil servants. *Scand J Work Environ Health* 2010;**36**:445–8.
40. Probst TM. Exploring employee outcomes of organizational restructuring: a Solomon four-group study. *Group Organ Manag* 2003;**28**:416–39.
41. Rohall D, Hamilton V, Segal D, et al. Downsizing the Russian army: quality of life and mental health consequences for former organizational members, survivors, and spouses. *Journal of Political & Military Sociology* 2001;**29**:73–91.
42. Vaananen A, Ahola K, Koskinen A, et al. Organisational merger and psychiatric morbidity: a prospective study in a changing work organisation. *J Epidemiol Community Health* 2011;**65**:682–7.
43. Woodward CA, Shannon HS, Cunningham C, et al. The impact of re-engineering and other cost reduction strategies on the staff of a large teaching hospital: a longitudinal study. *Med Care* 1999;**37**:556–69.
44. Brown GW, Moran P. Clinical and psychosocial origins of chronic depressive episodes. I: a community survey. *Br J Psychiatry* 1994;**165**:447–56.
45. Callan VJ, Terry DJ, Schweitzer R. Coping resources, coping strategies and adjustment to organizational change: direct or buffering effects? *Work Stress* 1994;**8**:372–83.
46. Mak AS, Mueller J. Negative affectivity, perceived occupational stress, and health during organisational restructuring: a follow-up study. *Psychol Health* 2001;**16**:125–37.
47. Vakola M, Nikolaou I. Attitudes towards organizational change: what is the role of employees' stress and commitment? *Employee Relations* 2005;**27**:160–74.
48. Swanson V, Power K. Employees' perceptions of organizational restructuring: the role of social support. *Work Stress* 2001;**15**:161–78.
49. Decker D, Wheeler GE, Johnson J, et al. Effect of organizational change on the individual employee. *Health Care Manag (Frederick)* 2001;**19**:1–12.
50. Zeitlin LR. Organizational downsizing and stress-related illness. *Int J Stress Manag* 1995;**2**:207–19.
51. Kasl S. Measuring job stressors and studying the health impact of the work environment: an epidemiologic commentary. *J Occup Health Psychol* 1998;**3**:390–401.
52. Tveit SD, Saksvik PO, Nytro K. Does change process healthiness reduce the negative effects of organizational change on the psychosocial work environment? *Work Stress* 2009;**23**:80–98.
53. Cunningham CE, Woodward CA, Shannon HS, et al. Readiness for organizational change: a longitudinal study of workplace, psychological and behavioural correlates. *J Occup Organ Psychol* 2002;**75**:377–92.
54. McDonough P. Job insecurity and health. *Int J Health Serv* 2000;**30**:453–76.

55. **Baillien E**, De Witte H. Why is organizational change related to workplace bullying? Role conflict and job insecurity as mediators. *Econ Ind Democr* 2009;**30**:348–71.
56. **Michel A**, Stegmaier R, Meiser D, *et al*. Burn out and unsatisfied? How change characteristics and change-specific job insecurity are related to emotional exhaustion, job satisfaction, and turnover intentions. *Z Arbeits- Organisations Psychologie* 2009;**53**:11–21.
57. **De Cuyper N**, Handaja Y, Witte H. Job insecurity during a restructuring: associations with engagement and emotional exhaustion. *Gedrag en Organisatie* 2008;**21**:97–112.
58. **Mansell A**, Brough P, Cole K. Stable predictors of job satisfaction, psychological strain, and employee retention: an evaluation of organizational change within the New Zealand customs service. *Int J Stress Manag* 2006;**13**:84–107.
59. **Lundgren SM**, Nordholm L, Segesten K. Job satisfaction in relation to change to all-RN staffing. *J Nurs Manag* 2005;**13**:322–8.
60. **Korunka C**, Scharitzer D, Carayon P, *et al*. Employee strain and job satisfaction related to an implementation of quality in a public service organization: a longitudinal study. *Work Stress* 2003;**17**:52–72.
61. **Woodward CA**, Shannon HS, Lendrum B, *et al*. Predictors of job stress and satisfaction among hospital workers during re-engineering: differences by extent of supervisory responsibilities. *Health Manage Forum* 2000;**13**:29–35.
62. **Hansen AM**, Larsen AD, Rugulies R, *et al*. A review of the effect of the psychosocial working environment on physiological changes in blood and urine. *Basic Clin Pharmacol Toxicol* 2009;**105**:73–83.

## Have confidence in your decision making.



The best clinical decision support tool is now available as an app for your iPhone. Visit [bestpractice.bmj.com/app](http://bestpractice.bmj.com/app)

**BestPractice**  
FROM THE BMJ EVIDENCE CENTRE

clinicians • medical students • nurses • healthcare practitioners