

consideration of time since median exposure and dose rate) and arsenic no increase in risk below 10 mg/m³-years was observed. Fixing the parameter estimate of the ERR in this range at 0 provided the best model fit with an ERR of 0.065 (95% CI: 0.043, 0.087) above 10 mg/m³-years. The silica-induced lung cancer risk decreased with increasing attained age. Data of a nested case-control study did not indicate a major correlation between silica and smoking status.

Conclusions The study confirms a positive exposure-response relationship between silica and lung cancer, particularly for high exposures. No clear evidence was observed in the low-dose range. Confounding by smoking is unlikely but residual confounding cannot be fully excluded.

363 DOES INHALATION OF SILICA DUST CAUSE CHRONIC RENAL DISEASE? - A SYSTEMATIC REVIEW

M M Möhner, Gellissen, Pohrt. *Federal Institute for Occupational Safety and Health, Berlin, Germany*

10.1136/oemed-2013-101717.363

Objective To prove the evidence for a causal relationship between exposure to silica dust and chronic renal disease, in particular Glomerulonephritis (GN)

Methods The review is based on a search of the Medline database for relevant epidemiological studies published between 1987 and 2012. The quality of selected papers was evaluated based on the Newcastle-Ottawa Scale and the CASP checklist. Meta-analyses were performed separately for cohorts of workers compensated for silicosis and for industry-based cohorts. In addition, synopses were undertaken for studies based on quantitative exposure estimates and for studies using end-stage renal disease (ESRD) as the outcome.

Results The review includes 10 cohorts of silicotics, 12 industry-based cohort studies, 5 case-control-studies and one pooled analysis. The pooling of the cohorts of silicotics did not show any excess risk for renal diseases (ICD-9: 580–589; SMR = 1,00 [0.84–1.18]_{95%}). By contrast, the meta-analysis of the industry-based cohort studies showed an elevated risk (SMR = 1,42 [1.15–1.75]_{95%}) when applying a random effects model to account for the strong heterogeneity of these studies. Five of those studies investigated the incidence of ESRD as the outcome. The corresponding relative risk estimators were in the range of 0.77 to 1.97, two of them were significantly elevated ($p < 0.05$). Three studies went further and investigated GN as the outcome. They resulted in markedly higher risk estimators (SIR in the range of 3.85 to 4.27) then for ESRD. Results could have been affected by a possible diagnostic bias and confounding by other occupational risk factors.

Conclusion The review does not provide sufficient evidence for establishing a causal link between exposure to silica dust and chronic renal disease. To clarify this relationship, further longitudinal studies are needed, which should make use of clinical or even pre-clinical data, including data of health surveillance programs for workers with high exposure to silica dust.

364 SMOKING CESSATION AND LUNG CANCER MORTALITY IN A HISTORICAL COHORT OF WORKERS WITH SILICOSIS IN HONG KONG, 1981–2006

¹Tse, ²Leung, ¹Qiu, ¹Yu. ¹The Chinese University of Hong Kong, NT, Hongkong; ²Pneumoconiosis Clinic, Department of Health, Hong Kong SAR, China

10.1136/oemed-2013-101717.364

Objectives To evaluate the impact of smoking cessation on the risk of lung cancer mortality among workers with silicosis.

Methods We recruited all 3202 incident cases of silicosis in Hong Kong from 1981 to 2005 and followed up them till 2006 to ascertain the causes of death. The follow-up rate was 97.5%. We collected each worker's socio-demographics, lifetime smoking habits, lifetime occupational history, and medical history at the initial assessment of diagnosis. We obtained the most recent information on smoking status from medical record. Multiple Cox's regression analysis was performed to examine the impact of smoking cessation.

Results A total of 1562 deaths (48.8%) occurred and 157 of them were from lung cancer. Lung cancer mortality was strongly associated with smoking [former smoking: hazard ratio (HR) = 3.41, 95% CI: 1.23 - 9.46; current smoking: HR = 5.80, 95% CI: 2.13 - 15.74], and a significantly positive gradient was indicated with smoking pack-year and the years of smoking. Lung cancer mortality tended to decreasing at the 10th year of cessation (HR = 0.64, 95% CI: 0.40 - 1.04) and a substantial decrease was observed after the abstinence of smoking for 20 or more years (HR = 0.22, 95% CI: 0.11 - 0.44). Relative to never smokers, the hazard ratio for lung cancer mortality for silicotics who had never quit smoking was 7.09 (95% CI: 2.57 - 19.59) and it became 4.75 (95% CI: 1.71 - 13.16) for the new quitters who changed their behaviour during the follow-up period, whilst a relatively lower risk (HR = 3.47, 95% CI: 1.25 - 9.61) was found among the persistent quitters.

Conclusions Lung cancer mortality decreased substantially with smoking cessation in workers with silicosis, particularly for those who persistently quit smoking for a longer period.

Acknowledgement Pneumoconiosis Compensation Fund Board, Hong Kong

365 PREDICTORS OF SILICOSIS IN AN INDUSTRY WIDE STUDY IN THE SOUTH AFRICAN GOLD MINING INDUSTRY

¹R I Ehrlich, ¹Knight, ²Fielding, ²Jeffery, ²Grant, ³Churchyard. ¹Centre for Occupational and Environmental Health Research, Cape Town, South Africa; ²London School of Hygiene and Tropical Medicine, London, United Kingdom; ³Aurum Institute, Johannesburg, South Africa

10.1136/oemed-2013-101717.365

Objectives Despite the scale of the silicosis problem in the South African goldmining industry, studies in active miners are scarce. A large randomised cluster trial of isoniazid tuberculosis prophylaxis provided an opportunity to investigate the predictors of silicosis and variation in silicosis prevalence across goldmining shafts.

Methods This analysis was based on a random sample of routine annual chest radiographs of active miners entering the trial, taken between 2004 and 2009. All were read for silicosis using the ILO classification by an experienced lay reader. All films classified as abnormal and a random sample of normals were re-read by a 'B'-reader and these readings substituted for those of the lay reader in case of disagreement. The association between silicosis defined as ILO > 1/0 and age at radiograph, years since first employment, shaft and other occupational markers, was analysed.

Results A total of 14,434 radiographs from 15 goldmining shafts were read. Silicosis prevalence varied from 0.68% to 6.28% across shafts; 7.8 fold across the range in adjusted analysis. Silicosis showed a strong dose response relationship with years since

first employment [OR 8.1 (95% CI 3.2, 21) for > 30 vs < 15 years]. Surprisingly, after adjusting for years since first employment, age was still predictive of silicosis risk ($p < 0.001$). Miners from Lesotho, who make up a large minority of the workforce, had an increased odds (OR 1.55; 95% CI 1.2, 1.9) of silicosis relative to South African miners.

Conclusions Goldmining shafts differ in silicosis risk independently of workforce age and length of mining service. Speculative explanations include higher silica fraction of the rock and greater intensity of exposure to silica. Basotho miners are at increased relative risk of silicosis, suggesting greater involvement in dustier jobs. The strong independent effect of age on silicosis risk, reported previously, remains to be explained.

Session: 28. Health impact analysis physical factors

366 THE ASSOCIATION BETWEEN HEAT EXPOSURE AND WORK-RELATED INJURIES IN SOUTH AUSTRALIA, 2001–2010

¹J X Xiang, ¹Bi, ²Pisaniello, ¹Hansen. ¹The University of Adelaide, Adelaide, Australia; ²The University of Adelaide/Discipline of Public Health, Adelaide, Australia

10.1136/oemed-2013-101717.366

Objective To investigate the association between work-related injuries and temperature, to identify the groups of workers at high risk of heat-related injuries, and to explore the possible lagged effects of extreme heat on work-related injuries.

Method Workers' compensation claims data obtained from SafeWork South Australia for the period of 2001–2010 were transformed into time series format and merged with daily meteorological data. The impacts of temperature on daily work-related injury rates were estimated by using generalised estimating equation model with negative binomial distribution, a log link function and a first order autocorrelation structure. A piecewise linear spline function was utilised to quantify the effect of temperature on work-related injury rates below and above thresholds. The day of the week and long-term trends were adjusted.

Result Overall, there was an association between work-related injuries and temperature in South Australia. One degree Celsius increase in temperature below 38 was associated with 0.2% increase of injury rate. However, the injury risk declined significantly above this temperature. Specifically, the following groups of workers were at high risk of heat-related injury: male workers (IRR 1.004, 95% CI 1.002–1.005); and young workers aged < 24 (IRR 1.003, 95% CI 1.000–1.006). Occupations at risk were labourers and related workers (IRR 1.004, 95% CI 1.001–1.006), intermediate production and transport workers (IRR 1.002, 95% CI 1.000–1.004), and tradespersons and related workers (IRR 1.003, 95% CI 1.001–1.006). Industries showing an association between injuries and temperature were agriculture (IRR 1.007, 95% CI 1.002–1.013), construction (IRR 1.005, 95% CI 1.002–1.008), finance, property and business services (IRR 1.006, 95% CI 1.001–1.012), and overall outdoor industries (IRR 1.004, 95% CI 1.002–1.006). A lagged effect of extreme heat on work-related injury rates has not been found.

Conclusion The risk of work-related injuries is significantly associated with heat exposure, especially for vulnerable groups in the workplace.

367 INVESTIGATING OCCUPATIONAL FACTORS AND BIOMARKERS OF KIDNEY FUNCTION AMONG NICARAGUAN WORKERS

¹M M McClean, ²Laws, ³Amador, ⁴Weiner, ⁵Kaufman, ⁶Ramirez Rubio, ⁷Sánchez Rodríguez, ⁸Brooks. ¹Boston University, Boston, United States of America; ²Rebecca Laws, Boston, United States of America; ³Juan Jose Amador, Managua, Nicaragua; ⁴Daniel Weiner, Boston, United States of America; ⁵James Kaufman, Boston, United States of America; ⁶Oriana Ramirez Rubio, Madrid, Spain; ⁷Jose Marcell Sánchez Rodríguez, Leon, Nicaragua; ⁸Daniel Brooks, Boston, United States of America

10.1136/oemed-2013-101717.367

Objective In Nicaragua, an epidemic of chronic kidney disease (CKD) with unknown aetiology has been described primarily among young, male sugarcane workers. Our goals were to characterise the type of kidney damage and evaluate the potential role of occupational factors.

Methods Our study population included 284 sugarcane workers, 51 miners, 60 construction workers, and 53 port workers in Western Nicaragua. For sugarcane workers in seven jobs (e.g. cane cutters, pesticide applicators, etc.), blood and urine samples were collected at the beginning and end of the 6-month sugarcane harvest. One round of samples were also collected from workers who were employed in the other three industries (but had never worked in the sugarcane industry). Biomarkers of kidney injury included serum creatinine, urinary albumin, and urinary neutrophil gelatinase-associated lipocalin (NGAL). Linear regression models were used to determine whether biomarkers of kidney injury increased during the harvest and/or varied by job.

Results Biomarkers of kidney injury were significantly different by sugarcane job and estimated glomerular filtration rate (eGFR) declined significantly during the harvest among those engaged in the most strenuous work tasks, while urine albumin remained low in all groups. Compared to factory workers, eGFR was lowest among cane cutters ($p = 0.006$) and urinary NGAL was highest among cane cutters ($p = 0.04$). Workers in other industries also had higher than expected prevalence of eGFR <60 mL/min/1.73 m².

Conclusions Our results provide evidence that the type of kidney damage occurring in sugarcane workers, as well as among workers in other industries, is primarily tubulointerstitial (and not glomerular) in nature. Biomarkers of kidney injury varied by job and were highest among workers engaged in more strenuous physical labour. If acute kidney damage is on the causal pathway to CKD, heat or other work-related exposures may be contributing to this epidemic.

368 OCCUPATIONAL NOISE EXPOSURE AND SERUM LIPIDS: THE IMPACT OF NOISE EXPOSURE LEVEL AND HEARING PROTECTION

A S S Schmedes, Arlien-Søborg, Stokholm, Hansen, Bonde, K L Christensen, Frederiksen, Kristiansen, Lund, Vestergaard, Wetke, Kolstad. Danish Ramazzini Center, Aarhus, Denmark

10.1136/oemed-2013-101717.368

Objectives Environmental and occupational noise exposure have been related to increased risk of cardiovascular disease. Activation of the physiological stress response and altered serum lipid levels have been proposed as alternative causal pathways. The objective of this study was to investigate the relation between occupational noise exposure and serum levels of total cholesterol, LDL-cholesterol, HDL-cholesterol, and triglycerides when