aimed to identify determinants of PM₁₀ and PM_{2.5} levels in poultry barns and evaluate the effectiveness of electrostatic precipitators (ESP) to reduce environmental air pollution.

Methods PM₁₀ and PM_{2.5} Harvard impactor samples were collected in “side-by-side” barns (one with an ESP, one without) at five poultry farms in British Columbia, Canada from 2008 to 2012. Measured particulate levels were analysed using multimodel inference and linear mixed-effects models after log-transformation. Random effects were added to account for clustering within farms, barns, and rearing cycles.

Results A total of 234 PM_{2.5} and 230 PM₁₀ valid samples were modelled. Geometric means of PM_{2.5} and PM₁₀ were significantly lower in barns with ESP control in place (151 and 427 g/m³, respectively) compared to barns with no control (334 and 969 g/m³), resulting in unadjusted % reductions of 47% and 50% respectively.

In statistical models, the fixed-effects explained 57% and 72% of the total variance in the PM_{2.5} and PM₁₀ measurements, respectively. The residual (i.e. within rearing cycle) and between rearing cycle variance were the most affected by adding the fixed effects structure. Strongest determinants in the models for both dust types were ESP use (i.e. approximately halving particulate levels for both PM_{2.5} and PM₁₀), bird age (i.e. 10–30 fold increase in particulate levels depending on bird and particulate type), and type of bird (i.e. approximately a 2.5 fold increase for PM_{2.5}, and four fold increase in PM₁₀ for turkey compared to chicken). Interactions were suggested in PM₁₀ models between type of bird, bird age, and ESP use.

Conclusions The use of ESP resulted in significant reductions in in-barn particulate even after controlling for other determinants such as bird age, type of bird, ventilation, and date.

Session: 3. Electro magnetic fields and health

239 MEASUREMENTS OF OCCUPATIONAL EXPOSURE TO EMF EMITTED BY HIGH-SPEED MAGLEV TRANSPORTATION SYSTEM AND ITS HEALTH EFFECTS

Z L Zhang, Z Z Zhou, W Y Wang, Y J Yang, C F Chu, Y Z Yu. *Third Military Medical University, Chongqing, China*

10.1136/oemed-2013-101717.239

Objectives High-speed Maglev transportation system is an advanced technology using magnetic forces to propel, levitate, and guide the train. Our study was to assess the EMF emitted by High-Speed Maglev Transportation System and the possible health effects of occupational exposure to the Maglev EMF.

Methods The static magnetic field of Maglev were measured by Narda ETM-1 magnetic field measurement system. The time-varying EMFs of Maglev were evaluated by PMM8053 EMF measurement system. 48 employees exposed to maglev EMF were selected as occupationally exposed group while 54 employees without any Maglev EMF exposure were selected as control. Questionnaires were sent to two groups. Complete blood count was done by haematology analyzer. Blood lipid level was detected by enzymatic method. Thyroid function related hormones was evaluated by chemical immune assay. Serum melatonin level was measured by an ELISA kit.

Results EMF strength of all Maglev sites was lower than the safety exposure limits of current international and Chinese national standard.

The symptoms of drowsiness, memory impairment, irritability, alopecia were related to occupational exposure to maglev EMF. Abnormal rate of cholesterol and high density lipid protein in exposed group were significantly higher than control. The blood cell counts, thyroid function, and serum melatonin level had no statistical difference between two groups.

Conclusions EMF emitted by the High-speed Maglev transportation system in different frequencies were lower than the exposure limits of current international and Chinese national standard.

Due to the population limits, we can not draw a conclusion that occupational exposure to Maglev EMF may have adverse health effects from the previous data. However, it may give us a clue that occupational exposure to maglev EMF may result in the alteration of neuronal function and lipid metabolism.

240 NO INFLUENCE OF MAGNETIC RESONANCE IMAGING SCAN ON MALE REPRODUCTIVE HORMONES

¹O J M Møllerlækken, ¹Moen, ²Magerøy. ¹University of Bergen, Bergen, Norway; ²Haukeland University Hospital, Bergen, Norway

10.1136/oemed-2013-101717.240

Objectives To investigate if magnetic resonance imaging (MRI) influences male reproductive hormones. The use of MRI is increasing around the world and the possible adverse effects on reproductive health of electromagnetic fields (EMF) in MRI are not previously studied. Previous articles have suggested that radio frequency (RF) electromagnetic fields may influence on male reproductive hormones follicle stimulating hormone (FSH) and luteinizing hormone (LH), and that this may cause reduced sperm quality.

Methods A randomised balanced cross-over study using real and sham MRI scan among 24 healthy male volunteers were conducted. Serum-blood samples of Inhibin B, testosterone, LH, FSH, sex-hormone binding globulin, oestrogen, prolactin and thyreotropine were taken immediately before and after the different scans and at a control session without any scan. Questionnaire data was gathered regarding possible confounding factors among the participants at each session. The RF EMF exposure caused by the MRI scanner was described as RF estimates for each sequence driven during the scan.

Results When investigating the hormone levels immediately before and immediately after the scan we found no differences, neither in the MRI scan setting nor in the sham setting. We also investigated if the hormones changed differently in the MRI setting versus the sham setting, but found no differences. We could neither find an effect after 11 days, which was the average number of days between the different sessions.

Conclusion EMF exposure during a MRI did not cause changes in male reproductive hormones. Adverse effects on other endpoints than male reproduction or possible chronic effect of multiple MRI scans have not been investigated in this study. To evaluate this, further studies should be carried out.

241 OCCUPATIONAL EXPOSURE TO EXTREMELY LOW FREQUENCY MAGNETIC FIELDS AND BRAIN TUMOURS IN THE INTEROCC STUDY

¹C Turner, ²Benke, ³Bowman, ¹Figueroa-Alquezar, ⁴Fleming, ⁵Hours, ⁶Kindl, ⁷Krewski, ⁸Lavoue, ⁹McClean, ¹⁰Parent, ⁸Richardson, ¹¹Sadetzki, ¹²Schlaefler, ¹²Schlehofer, ⁸Siemietycki, ¹³Van Tongeren, ¹Cardis. ¹Centre for Research in Environmental Epidemiology, Barcelona, Spain; ²Monash University, Melbourne, Australia; ³National Institute for