uranium co-exposure. A potential excess risk of lung cancer related to chromium (VI) exposure observed among nuclear workers should be confirmed in further studies.

247 MORTALITY STUDY IN TWO ITALIAN COHORTS OF WORKERS EXPOSED TO PCBs: EXTENSION OF THE FOLLOW-UP UP TO 2006

¹A C Pesatori, ²Grillo, ²Consonni, ³Caironi, ³Sampietro, ¹Bertazzi. ¹University of Milan, Milan, Italy; ²IRCCS Foundation, Ospedale Maggiore Policlinico, Milan, Italy; ³Local Health Unit (ASL), Bergamo, Italy

10.1136/oemed-2013-101717.247

Objectives PCBs have been classified as carcinogenic in animals and probably carcinogenic to humans (IARC1987). Epidemiological studies showed increased risks for Non-Hodgkin's Lymphoma, liver and biliary tract, digestive, skin, prostate and brain cancer but findings are still controversial. We present the results of the extension of the follow-up as of 2006 of two cohorts of workers potentially exposed to PCBs in capacitor manufacturing plants in Lombardy, Northern Italy. The mortality of the two cohorts was previously examined up to 1991 (Plant 1) and 2002 respectively (Plant 2).

Methods The study population included 544 males and 1551 females employed between 1946 and 1978 in Plant 1 and 373 males and 97 females employed between 1950 and 1988 in plant 2. Standardised mortality ratios (SMR) and 95% Confidence Intervals (95% CI) were calculated (in the whole cohort, separately in plant 1 and 2 and among males and females) using as reference the regional mortality rates specific by age and calendar period.

Results In the whole cohort, increased mortality from biliary tract and "other digestive" cancer, prostate, brain cancer, Hodgkin and non-Hodgkin lymphoma (8 deaths, SMR 1.65; 95% CI 0.8–3.3) were detected. The excess mortality from biliary tract cancer was confined to males (4 deaths, SMR 3.9; 95% CI 1.5–10.4), where as the increased mortality from Hodgkin lymphoma (4 deaths, SMR 3.6, 95% CI 1.3–9.5) and brain cancer (7 deaths, SMR 2.1; 95% CI 1.02–4.5) was observed in plant 1 only. The excess mortality from other digestive cancer and NHL was equally distributed in the two cohorts and among males and females. Analyses by duration of employment and time since first employment did not show consistent trends.

Conclusions Our findings suggest increased mortality from cancers of "a priori" interest in PCB exposed workers, however these excesses were not clearly related to duration of employment and latency.

248 OCCUPATIONAL EXPOSURE TO TRICHLOROETHYLENE AND PERCHLOROETHYLENE AND THE RISK OF LYMPHOMA, LIVER AND KIDNEY CANCER IN FOUR NORDIC COUNTRIES

¹J J Vlaanderen, ¹Straif, ²Pukkala, ³Kauppinen, ⁴Kyyronen, ⁵Martinsen, ⁵Kjaerheim, ⁶Tryggvadottir, ⁷Hansen, ⁸Sparen, ⁹Weiderpass. ¹IARC, Lyon, France; ²School of Public Health, University of Tampere, Tampere, Finland; ³Finnish institute of occupational health, Helsinki, Finland; ⁴Finnish cancer registry, Helsinki, Finland; ⁵Cancer registry of Norway, Oslo, Norway; ⁶Faculty of Medicine, University of Iceland, Reykjavik, Iceland; ⁷Institute of Cancer Epidemiology, Danish cancer society, Copenhagen, Denmark; ⁸Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden; ⁹Folkhalsan Research Center, Samfundet Folkhalsan, Helsinki, Finland

10.1136/oemed-2013-101717.248

Objectives Trichloroethylene (TCE) and Perchloroethylene (PER) are two chlorinated solvents that have been applied widely as degreaser of metal parts and in dry cleaning and other applications. In 2012 the International Agency for Research on Cancer classified TCE as *carcinogenic to humans* and PER as *probably carcinogenic to humans*. We explored exposure-response relations for TCE and PER and non-Hodgkin lymphoma (NHL), multiple myeloma (MM), and cancers of the kidney and liver in the Nordic Occupational Cancer (NOCCA) cohort which holds information on occupation and long-term follow-up for cancer for the populations of Norway, Sweden, Finland and Iceland.

Methods The cohort was set-up by linking occupational information from censuses to national cancer registry data utilising personal identity codes in use in all the Nordic countries. Country, time-period, and job-specific exposure estimates were generated for TCE, PER, and potentially confounding occupational exposures with a comprehensive job-exposure matrix. Two exposure metrics were created: cumulative exposure and 'average intensity times prevalence' of exposure. Conditional logistic regression was conducted for exposure groups as well as for continuous cumulative exposure.

Results Hazard ratios for liver cancer, NHL, MM, but not kidney cancer were slightly elevated in groups with high exposure to PER (compared to occupationally unexposed subjects). Hazard ratios for liver cancer and NHL also increased with increasing continuous exposure to PER. We did not observe evidence for an association between exposure to TCE and NHL, MM, and liver and kidney cancer.

Conclusions Although this study was subject to limitations related to the low prevalence of exposure to PER and TCE in the Nordic population and a necessarily rudimentary exposure assessment strategy, we observed some evidence indicative of an excess risk of cancer of the liver and NHL in subjects exposed to PER. Exposure misclassification may have diluted the observed associations in this study.

249 A COMPREHENSIVE DANISH OCCUPATIONAL CANCER LINKAGE

J H Hansen, Meersohn. Danish Cancer Society, Copenhagen, Denmark

10.1136/oemed-2013-101717.249

Objectives Linkages of cancer and occupation provide an important resource to aid in our understanding of the role of occupational exposures in the aetiology of cancer. The majority of existing surveys have, however, been based on census data including occupational information from only one certain date, thus lacking the general situation with several changed between different jobs during a work career. We report on a series of 60 Danish nested case-control studies based on nationwide data linkages, including information on all employments since 1964.

Methods All cancer cases (N = 1,031,504; 50.4% women and 49.6% men) diagnosed between 1970 and 2010, aged 18 and 84 years old, was retrieved from the Danish Cancer Registry. Each subject was individually linked with employment information from the Nationwide Pension Scheme with compulsory membership, including information on start and end of each employment, and a 5-digit hierarchically trade code. Information on job title, civil and vital status was obtained from the Central Population Register (CPR). Controls free of cancer and matched on birth year and sex have been selected from the