important in compensation cases. The cause is probably a considerable indirect exposure.

273 THE LATENCY PERIOD OF MESOTHELIOMA AMONG A COHORT OF BRITISH ASBESTOS WORKERS (1978–2005)

G F Frost, Harding. *Health & Safety Laboratory, Buxton, United Kingdom*

10.1136/oemed-2013-101717.273

Objectives To investigate the hypothesis of an inverse relationship between intensity of asbestos exposure and mesothelioma latency among a cohort of asbestos workers in Britain.

Methods The Great Britain (GB) Asbestos Survey recruited asbestos workers attending medical examinations as required under the asbestos regulations in GB. Job details and smoking information were updated at any subsequent examinations. The primary outcome for this analysis was the time from first occupational exposure to asbestos to death with mesothelioma. Univariable and multivariable generalised gamma accelerated failure-time (AFT) models were used to estimate time ratios (TRs). Those who died with mesothelioma between 1978 and 2005 were included in the analysis.

Results After exclusions for missing data, there were 614 workers in the GB Asbestos Survey who died with mesothelioma between 1978 and 2005. Total follow-up time was 9,280 person-years, with a median latency of 22.8 years [95% confidence interval (CI) 16.0–27.2 years]. In the final multivariable AFT model, mesothelioma latency was around 26% longer for females compared to males (TR = 1.26, 95%CI = 1.15–1.37), and 5% shorter for those who died with asbestosis compared to those who did not (TR = 0.95, 95%CI = 0.91–0.99) after adjustment for year of first exposure, age at first exposure, and duration of exposure. There was no evidence of associations between mesothelioma latency and occupation, smoking status and mesothelioma type in either univariable or multivariable analyses.

Conclusions The associations between mesothelioma latency and both sex and presence of asbestosis were in the directions expected if the intensity hypothesis was true. However, there was no evidence that asbestos insulation workers, who tended to experience greater asbestos exposures than other occupations, had shorter latencies than other asbestos workers. Therefore this study did not find sufficient evidence to either support or refute the intensity hypothesis. © Crown Copyright (2012).

274 BIOMARKERS FOR PLEURAL MESOTHELIOMA IN THE MEXICAN INSTITUTE SOCIAL SECURITY

¹G A M Aguilar-Madrid, ²Juarez-Perez, ³Calderón Aranda, ²Kelly García, ¹Alvarado Cabrero, ²Ochoa Vázquez. ¹Instituto Mexicano Del Seguro Social, Mexico, Distrito Federal, Mexico; ²Instituto Mexicano Del Seguro Social, Mexico, Distrito Federal, Mexico; ³Centro de Investigación y Estudios Avanzados. CINVESTAV. Depto Toxicología, México, DF, Mexico

10.1136/oemed-2013-101717.274

Objective Developing a new diagnostic strategy for Malignant Pleural Mesothelioma measuring soluble proteins in plasma and pleural fluid using ELISA and qRT-PCR.

Methods Case-control study, in which the cases were incident patients with malignant pleural mesothelioma in two hospitals from Mexican Social Security (Neumology and Oncology) and there were 4 controls for each case. A questionnaire was applied for asbestos exposure. ELISA with double antibody was used to

determine the presence of thrombomodulin and mesothelin, using known concentration curves for each one. An statistical descriptive analysis was made as well as logistic regression models and ROC curves.

Results In this study were enrolled 35 cases and 136 controls. The mean concentrations, (SD) and [min-max] in plasma for thrombomodulin were: 10.2 (7) ng/mL [1.6–32.2] for the cases, and 5.4 (5.1) ng/mL [0.78–32.5] for the controls. For mesothelin values were: 104.3 (78.2) ng/mL, [12.2–351.2] for the cases, and 25.2 (3.2) ng/mL, [6.5–330.3] for the controls. Both biomarkers were statistically significant. In the ROC curve for both biomarkers sensibility of 82.3% and specificity of 95.5% were achieved. In the nonconditional logistic regression model, mesothelin and thrombomodulin had an OR = 1.03 (CI: 1.02–1.05) and OR = 1.08 (CI: 1.01–1.16), respectively, adjusted for age and sex.

Conclusions A unique parameter to evaluate a diagnostic test in every situation doesn't exist, but following Sackett and Álvarez-Cáceres guidelines, our two biomarkers have a high specificity (95.5%) and sensibility (82.35%) with a cut-off point, in this sample, ≥83 ng/mL for mesothelin and ≥10.5 ng/mL. Nevertheless we consider that our sample is small and that we also need to control the factors, such as diabetes mellitus, hypercholesterolemia and renal failure, that have influence in the level of both biomarkers.

275 OCCUPATION AND MESOTHELIOMA IN PLEURA AND PERITONEUM IN SWEDEN - UPDATED INCIDENCE FOR MALES AND FEMALES 1961–2009

¹N P Plato, ¹Weiderpass, ¹Sparren, ²Martinsen. ¹Karolinska Institutet, Stockholm, Sweden; ²Cancer Registry of Norway, Oslo, Norway

10.1136/oemed-2013-101717.275

Objectives Mesothelioma incidence increased in Sweden between years 1970 and 2000, and seems to have reached a plateau in the last decade. We used data from the Swedish component of the Nordic Occupational Cancer Study - NOCCA - to study potential occupational exposures other than asbestos that may be associated with mesothelioma occurrence in men and women.

Methods Follow up was done by linkages of the Swedish NOCCA study database, including 53 occupational categories, with the population based Swedish registries (cancer, mortality, total population) from years 1961 to 2009, and matched with the Swedish version of NOCCA-JEM, which include 25 carcinogenic substances with exposure levels for 283 occupations in different employment periods from year 1945 to 1994. Multivariate analysis with 20 years delay for cumulative exposure were done.

Results A total of 4180 incident mesothelioma of the peritoneum and pleura were reported to the Swedish Cancer Registry (1961 - 2009): 26.6% were women. 83.9% were located in the pleura (18% women). We found a significant over risk in mesothelioma in 16 of 53 occupational categories, and a clear gender difference. We observed increased risk of mesothelioma among female textile workers, building cleaners, seamstresses, printers, packers and postal workers. Men had over risks in construction workers, electricians, masons, and insulation workers; the greatest excess risk of pleural mesothelioma was observed among plumbers (SIR 5.1, 95% CI 4.30 to 6.01). In multivariate analysis, controlling for other occupational exposures, significant associations were observed for asbestos exposure and mesothelioma risk.

Conclusion Exposure to asbestos is associated with incidence of mesothelioma in Sweden. There were some occupations