(aOR=2.4, 95% CI=1.8–3.1 for SRH; aOR=3.0, 95% CI=2.0–4.5 for PHU), and client-related burnout (aOR=1.6, 95% CI=1.2–2.3 for SRH; aOR= 2.1, 95% CI=1.2–3.5 for PHU) while adjusted for significant variables.

Conclusions We concluded that nurses worked in public health units and secondary referral hospitals had higher job strain and work-related burnout as compared to primary clinics. Further study should examine the stressors from these workplaces and follow up the health effects of high strain and burnout status.

0164 METAGENOMIC DETECTION OF BACTERIA IN AEROSOL SAMPLES IN ANIMAL SLAUGHTERHOUSES TO DEVELOP EXPOSURE PROFILES FOR AN EPIDEMIOLOGICAL ANALYSIS

¹David McLean, ²Patrick Biggs, ³Mily Leblanc-Maridor, ⁴Richard Hall, ²Nigel French, ⁵Neil Pearce, ¹Jeroen Douwes. ¹Massey University, Wellington, New Zealand; ²Massey University, Palmerston North, New Zealand; ³Université de Nantes, Nantes, France; ⁴Institute of Environmental Science and Research, Upper Hutt, New Zealand; ⁵London School of Hygiene and Tropical Medicine, London, UK

10.1136/oemed-2014-102362.65

Objectives Significant excess risks of lung cancer and haematologic neoplasms have been observed in slaughterhouse workers in eight New Zealand studies, and numerous studies conducted elsewhere. No specific causal agents have been identified, although a biological aetiology is suggested as the risk is highest in those areas where workers are exposed to live animals or to biological material containing animal urine, faeces or blood. This study aimed to assess the airborne bacterial microflora in the slaughterhouse environment in order to develop exposure categories for reanalysis of a meat workers' cohort.

Method Bulk air samples (n = 31) were collected for between 5 and 8 h in five areas in both sheep and beef slaughterhouses using a SASS3100 sampler (fitted with a proprietary SASS filter) located between 0.5 and 2 metres from the worker. Nucleic acid was extracted from each filter and amplified using commercially available kits, then sequenced on an Illumina MiSeq instrument. Bioinformatics analyses conducted included comparative taxonomic analyses, gene function (including virulence factor) analyses, and principal component analyses to compare profiles in samples taken in different areas.

Results Of the bacteria identified over 95% were in the classes Actinobacteria, Firmicutes and Proteobacteria. Clear differences in all parameters were apparent in the different areas, however, and the full results of the comparative analyses and the development of exposure profiles will be presented.

Conclusions Metagenomic analysis of bioaerosol samples represents a promising method for the development of exposure categories for the epidemiological analysis of the effect of biological exposures in an occupational environment.

0168 SOMATISING TENDENCY, OCCUPATIONAL STRAIN AND MUSCULOSKELETAL SYMPTOMS: RESULTS FROM A LONGITUDINAL STUDY AMONG ITALIAN NURSES

^{1,2}Matteo Bonzini, ¹Lorenza Bertù, ²Marco Conti, ¹Alessia D'Amato, ¹Giovanni Veronesi, ³David N Coggon, ^{1,2}Marco M Ferrario. ¹Epidemiology and Preventive Medicine Research Centre, Insubria University, Varese, Italy; ²MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK; ³Ospedale Di Circolo Fondazione Macchi, Varese, Italy

10.1136/oemed-2014-102362.66

Objectives Musculoskeletal symptoms are a common cause of disability, with major impact on workforce wellbeing, absenteeism and productivity. Several, mainly cross-sectional, studies have linked such symptoms to physical workload, and also to psychological and socio-cultural factors.

We investigated whether prolonged or increasing job strain, tendency to somatise and other individual characteristics, are associated with worsening musculoskeletal pain.

Method As part of the CUPID study, we investigated a cohort of nurses employed on medical wards at the Varese University Hospitals (Italy). Participants were asked, at baseline and after one year of follow-up, about individual and occupational risk factors, psychological characteristics (including tendency to somatise), occupational strain (by Siegrist's Effort/Reward Imbalance Questionnaire-ERI), and musculoskeletal symptoms. Associations of worsening musculoskeletal pain with perceived job strain were assessed by multivariate log-binomial regression.

Results Occupational stress was associated with pain in the lower back (LBP) and neck/shoulder (NSP) in both cross-sectional questionnaires.

Comparing baseline and follow-up answers, workers who reported an increase in perceived stress showed more frequent worsening of both LBP (prevalence of worsening symptoms=41%, OR when compared with not stressed=1.7, 95% CI=1.1–2.7) and NSP (prevalence of worsening=51%, OR=1.2, 95% CI=0.8–1.8).

This relationship persisted after adjustment for gender, age and BMI, and exposure to physical workload, and was more evident among subjects with a tendency to somatise (OR=2.8, 95% CI=1.0–7.4 for LBP; OR=1.6, 95% CI=0.8–3.2 for NSP).

Conclusions Our observation suggests that tendency to somatise modifies individual responses to "triggering exposures", such as psychological workload, with important implications for the health, and productivity of workers.

0169 SINONASAL CANCERS: IS INTESTINAL TYPE ADENOCARCINOMA THE ONLY RELATED TO OCCUPATIONAL EXPOSURES? RESULTS FROM AN ITALIAN CASE-CONTROL STUDY

^{1,2}Matteo Bonzini, ³Laura Zanetta, ¹Lorenza Bertù, ^{3,4}Davide Parassoni, ³Mario Turri Zanoni, ³Davide Lepera, ^{3,2}Paolo Castelnuovo, ^{1,2}Marco M Ferrario. ¹Epidemiology and Preventive Medicine Research Centre, Insubria University, Varese, Italy; ²Insubria University, Varese, Italy; ³Ospedale Di Circolo, Fondazione Macchi, Varese, Italy; ⁴School of Specialisation in Occupational Health, University of Brescia, Brescia, Italy

10.1136/oemed-2014-102362.67

Objectives Epithelial sinonasal cancers (SNC) are rare, severe diseases associated to the exposure to several well-established carcinogens (IARC). The etiologic role of these carcinogens in different histological subtypes is still disputed, with several studies focusing on intestinal-type adenocarcinoma (ITAC) as the most (and maybe the only) occupational-related subtype. To assess the role of occupational exposures in SNC aetiology we designed a case control study, in which occupational exposures prevalence in two group of ITAC cases and non-ITAC were compared to controls.

Method In a large Italian hospital we enrolled 50 consecutive surgical non-ITAC cases (mainly squamous-cell carcinoma), 50 consecutive ITAC cases and 50 non-neoplastic patients (controls). Previous occupational exposures to wood and leather dust, solvents, metals, formaldehyde were investigated through a

Oral presentation

standardised interview. Gender, age, residence area and smoking were collected as potential confounders. Multivariate logistic regression was applied.

Results Considering all tumours together, we observed large increased risks for wood exposure (OR=6.9, 95% CI=3.0–16.3) and leather (prevalence 24% in tumours, 0% among controls) only. Compared to controls, we observed an increased risk for wood exposure [OR=7.7 (95% CI=2.6–22.5)] in ITAC cases, but not in non-ITAC cases [OR=0.8 (95% CI=0.2–3.1)]. Prevalence of leather exposure was 42% among ITAC and 6% in non-ITAC.

Conclusions Our case control study confirmed that ITAC cases but not other histotypes were strongly related to occupational exposures, and in particular to leather and wood dusts. Grouping together all SNC types reduce the causal role of occupation exposures. Larger samples size are needed to investigate other work-related carcinogens.

0171 DO FAMILY AND INDIVIDUAL CHARACTERISTICS AFFECT THE EXPERIENCE OF PHYSICAL AND PSYCHOSOCIAL WORK ENVIRONMENT IN DANISH 20/21 YEAR OLDS?

¹Trine Nohr Winding, ^{1,2}Merete Labriola, ^{3,4}Ellen Aagaard Nohr, ¹Johan Hviid Andersen. ¹Danish Ramazzini Centre, Department of Occupational Medicine, Regional Hospital Herning, Herning, Denmark; ²Department of Clinical Social Medicine, Public Health and Quality Management, Central Denmark Region and Section of Clinical Social Med, and Rehabilitation, School of Public Health, Aarhus University, Aarhus, Denmark; ³University of Southern Denmark, Institute of Clinical Research, Research Unit of Gynaecology and Obstetrics, Odense, Denmark; ⁴Department of Public Health, Section for Epidemiology, Aarhus University, Aarhus, Denmark

10.1136/oemed-2014-102362.68

Objectives To describe the work environment of Danish 20/21 year olds and to investigate the influence of family socioeconomic background or individual characteristics at age 14/15 on later experience of physical and psychosocial work environment.

Method The study population consisted of 695 young people with primary work affiliation at age 20/21 who were derived from a prospective youth cohort. Outcome information from the questionnaire in 2010 consisted of six questions about psychosocial work environment and two questions about physical work environment. Exposure information about school performance, vulnerability, health and parental socioeconomic status was derived from the questionnaire in 2004 and from registers.

Results Overall, the psychosocial work environment of the young people was good but they experienced more repetitive movements and hard physical work than older workers. Individual as well as family factors in late childhood all together only had limited impact on how young people report later work environment. Low self-esteem at age 14/15 was associated with experiencing high demands, low trust and low fairness at work. In girls low self-esteem and low sense of meaningfulness were associated with experiencing low influence at work. Low parental socioeconomic status was associated with poor physical work environment.

Conclusions This study showed a social gradient in experiencing poor physical work environment at age 20/21. The psychosocial work environment in young people was on average good, but it seems that vulnerable young people need special intention in order to prevent them from being selected into psychosocial demanding job functions later in life.

0173 GROUPING STRATEGIES FOR EXPOSURE ASSESSMENT OF THE PSYCHOSOCIAL WORK ENVIRONMENT

¹Morten Vejs Willert, ^{1,2}Vivi Schlünssen, ²Ioannis Basinas, ¹Zara Ann Stokholm, ³Matias Brødsgaard Grynderup, ⁴Johan Hviid Andersen, ⁵Reiner Rugulies, ³Åse Marie Hansen, ⁶Linda Kærlev, ⁷Jane Frølund Thomsen, ⁷Marianne Agergaard Vammen, ¹Henrik Kolstad. ¹Danish Ramazzini Centre, Department of Occupational Medicine, Aarhus University Hospital, Region Midt, Denmark; ²Danish Ramazzini Centre, Department of Occupational Medicine, Herning University Hospital, Region Midt, Denmark; ³Research Unit of Clinical Epidemiology, Institute of Clinical Research, University of Southern Denmark, Region Southern Denmark, Denmark; ⁴Department of Public Health, Aarhus University, Aarhus, Denmark; ⁵Department of Public Health, University of Copenhagen, Copenhagen, Denmark; ⁶Research Centre for the Working Environment, Copenhagen, Denmark; ⁷Department of Occupational and Environmental Medicine, Bispebjerg University Hospital, Capital Region, Denmark

10.1136/oemed-2014-102362.69

Objectives Individual response style, mood, expectations, and health status may affect reporting of the psychosocial work environment, and bias associations with outcomes. Reporting bias may be avoided by aggregating individual responses, ideally preserving exposure contrast. In this study, we examined the degree of exposure contrast yielded by different grouping strategies.

Method In 2007, we enrolled 4489 public employees from Aarhus, Denmark in the PRISME-cohort, with follow-up in 2009. From pay-roll registers we grouped workers at 2 organisational levels: department (n = 22) and work unit (n = 751), and 3 occupational levels: sector (n = 7), profession (n = 46), and job title (n = 77). Exposures, calculated as means of items scored on 5-point Likert scales, included psychological demands, decision latitude, social support, effort, reward, and procedural and relational justice. To assess variance components, we fitted linear mixed effect models with exposures as dependent variables, and id and grouping variables as random effects. Results are reported as the contrast in mean exposure levels e.g. between-group variance/ (between-group variance + within-group variance).

Results Within each hierarchy contrasts rose with increasing group-level detail. Grouping by either work unit (wu) or by job title (jt) contrasts were: psychological demands: 0.28(wu); 0.26(jt), decision latitude: 0.24(wu); 0.32(jt), social support: 0.24(wu); 0.06(jt), effort: 0.23(wu); 0.16(jt), reward: 0.19(wu); 0.12(jt), procedural justice: 0.24(wu); 0.14(jt), and relational justice: 0.29(wu); 0.04(jt).

Conclusions Grouping by work unit gave the most consistent contrasts (0.19–0.29), while grouping by job title varied considerably (0.04–0.32). These preliminary findings suggest that grouping by work unit provided better exposure contrasts than grouping by job title for all exposures, but decision latitude.

0182 PREDICTORS OF WORKERS' COMPENSATION CLAIM DURATION AMONG WORKERS DISABLED DUE TO LOW BACK PAIN

¹Jason Busse, ²Ivan Steenstra, ¹Shanil Ebrahim, ¹Diane Heels-Ansell, ¹Stephen Walter, ¹Gordon Guyatt. ¹McMaster University, Hamilton, Ontario, Canada, ²Institute for Work and Health, Toronto, Ontario, Canada

10.1136/oemed-2014-102362.70

Objectives Low back pain (LBP) is a common complaint among workers receiving Workers' Compensation wage replacement benefits. We used the administrative data from the Ontario Workplace Safety and Insurance Board (WSIB) to explore the association between baseline characteristics and commonly reimbursed therapies and time to claim closure among workers disabled due to LBP.