from pleura cancer and melanoma. This cohort study is the most informative ever conducted in France among nuclear workers.

0278 THE PESTIMAT PROGRAM: DEVELOPMENT OF A CROP EXPOSURE MATRIX FOR PESTICIDE EXPOSURE ASSESSMENT IN AGRICULTURE

^{1,2}Isabelle Baldi, ^{1,2}Camille Carles, ¹Anne Cantagrel, ^{3,4}Yannick Lecluse, ^{3,4}Elodie Niez, ⁵Pascal Fabbro-Peray, ^{3,4}Pierre Lebaillly. ¹Univ. Bordeaux, ISPED, Laboratoire Santé Travail Environnement, Centre INSERM U 897 Epidemiologie-Biostatistique, F-33000 Bordeaux, France; ²CHU de Bordeaux, Service de Médecine Du Travail, F-33000 Bordeaux, France; ³INSERM, UMR 1086- Cancers Et Préventions, F-14000 Caen, France; ⁴Centre François Baclesse, F-14000 Caen, France; ⁵DIM, Hôpital G Doumergue, F-34000 Montpellier, France

10.1136/oemed-2014-102362.108

Objectives Pesticides include ~1000 chemicals with various toxicological properties. Pesticide exposure assessment represents a crucial methodological issue for epidemiological studies. A reconstitution of pesticides used in entire professional careers based only on the memory is questionable. PESTIMAT is a crop exposure matrix, which intends to reconstitute parameters of pesticide exposure in France.

Method PESTIMAT is made of tables crossing crops and active ingredients (a.i.) by year from 1950 to 2010 and providing the following exposure parameters: 1) probability corresponding to the proportion of farmers having used the a.i. (in%), 2) frequency expressed as the number of treatment days, 3) intensity documented by the application rate of the a.i. (in kg/ha). Parameters were obtained by the combination of six complementary sources: i) registrations and withdrawals from the Agriculture Ministry, ii) information from ACTA* on products marketed iii) agricultural recommendations by the Plant Health Protection body, iv) treatment calendars provided by farmers, v) information from professional associations of farmers, vi) data from the industry

Results To date, 529 a.i. usable between 1950 and 2010 on 4 crops are included in PESTIMAT, corresponding to 160 fungicides (30%), 160 herbicides (30%) and 209 insecticides (40%). The total numbers were comparable on wheat/barley (N = 293), vineyards (N = 280), and apple-growing (N = 267) but lower on corn (N = 196). We will present an illustration with data obtained for dithiocarbamates fungicides, and examples of use in epidemiological studies, e.g. PHYTONER and AGRICAN cohorts.

Conclusions The multisource information available in the PESTIMAT matrix should enable to assess associations between pesticides and health at the a.i. level.

0282 A CROSS-SECTIONAL STUDY OF MARKERS OF EARLY IMMUNOLOGICAL AND CARDIOVASCULAR HEALTH EFFECTS AMONG A POPULATION EXPOSED TO CARBON NANOTUBES: THE CANTES STUDY

¹Roel Vermeulen, ²Anjoeka Pronk, ¹Jelle Vlaanderen, ³Dean Hosgood, ⁴Nat Rothman, ⁴Allan Hildesheim, ⁴Debra Silverman, ¹Annemarie Melis, ²Suzanne Spaan, ²Eef Voogd, ⁵Peter Hoet, ⁵Lode Godderis, ⁴Qing Lan. ¹IRAS Utrecht University, Utrecht, The Netherlands; ²TNO, Zeist, The Netherlands; ³Albert Einstein College of Medicine, New York, NY, USA; ⁴National Cancer Institute, Bethesda, MD, USA; ⁵University of Leuven, Leuven, Belgium

10.1136/oemed-2014-102362.109

Objectives To assess the association between occupational exposure to carbon nanotubes (CNTs) and early immunological and cardiovascular health effects.

Method We collected biological samples (blood, urine, buccal and nasal epithelium cells) from a group of workers in a CNT production factory and approximately age and sex matched unexposed individuals working in a different workplace. Based on an initial assessment of exposure data collected in the weeks before biological sampling we further classified CNT production workers as operators (high probability of exposure) and R&D workers (lower probability of exposure). We analysed blood samples for a panel of inflammatory markers.

Results We included 8 operators (median age 34.8), 16 R&D workers (median age 32.2), and 43 controls (median age 30.6). Preliminary analyses provide some evidence for an association between CNT exposure and selected cytokines. We observed a similar pattern when we restricted our analyses to non-smoking men. Concentrations of cytokines were exposure dependently upregulated with higher levels among operators than R&D workers; both higher than unexposed workers (p for trend <0.05).

Conclusions We observed some indications of early immunological health effects in a pilot study conducted among workers exposed to CNTs. Further analyses are planned, including assessment of lung function, heart rate variability, oxidative stress, and blood coagulation markers. Extensive exposure measurements were conducted in the CNT production factory as well and additional analyses will use these individual exposure measurements to more thoroughly explore exposure dependent effects.

0285 DEVELOPMENT OF A SOURCE-BASED APPROACH TO ASSESSING OCCUPATIONAL EXPOSURE TO ELECTROMAGNETIC FIELDS IN THE INTEROCC STUDY

¹Javier Vila, ²Joseph D Bowman, ³Laurel Kincl, ²Dave L Conover, ⁴Martie van Tongeren, ¹Jordi Figuerola, ³Lesley Richardson, ¹Elisabeth Cardis, ⁶on behalf of the INTEROCC Study Group. ¹Centre for Research in Environmental Health (CREAL), Barcelona, Spain; ²The National Institute for Occupational Safety and Health (NIOSH), Cincinnati, OH, USA; ³Oregon State University (OSU), Corvallis, OR, USA; ⁴Institute of Occupational Medicine (IOM), Edinburgh, UK; ⁵University of Montreal Hospital Research Centre (CRCHUM), Montreal, Canada; ⁶Institute of Occupational Medicine (IOM), Edinburgh, Spain

10.1136/oemed-2014-102362.110

Objectives Exposure to electromagnetic fields (EMF) has become ubiquitous in modern life and concern has increased regarding possible associated health effects. To date, assessment of occupational exposure has relied on job-exposure matrices, with exposure estimates for very broad occupational categories. To move EMF research forward, a new approach was necessary. A source-based strategy, incorporating detailed information on tasks, equipment used and work organisation could allow a more individualised exposure assessment.

Method Information on occupational histories and sources of EMF was collected as part of the INTERPHONE-INTEROCC study, providing an opportunity to assess occupational EMF exposure by assigning exposure to each source used. A source-exposure matrix (SEM) was developed based on measurements identified in the literature and estimates obtained through experts' elicitation, for sources without available measurements. This paper focuses on the SEM development methodology to ensure the quality and representativeness of the estimates.

Results Estimates of exposure for 138 EMF sources were obtained from measurements (1424 aggregated records) extracted from 71 papers and hygiene reports (1974–2013). For each source, exposure was calculated by frequency band and dosimetry type, as the arithmetic and geometric means of all measurements identified. Standard deviations were included in order to characterise the variability of the estimates.

Conclusions A source-exposure matrix has been constructed for the most common sources of EMF in the workplace, based on the responses to the INTERPHONE-INTEROCC study questionnaire. This database currently represents the most comprehensive source of information on occupational EMF exposure and is available on request to researchers.

0286 OCCUPATIONAL USE OF INSECTICIDES, FUNGICIDES AND FUMIGANTS AND RISK OF NON-HODGKIN LYMPHOMA AND MULTIPLE MYELOMA IN THE AGRICULTURAL HEALTH STUDY

¹Michael Alavanja, ¹Jonathan Hofmann, ²Charles Lynch, ³Cynthia Hines, ¹Kathryn Barry, ⁴Joseph Barker, ⁴Dennis Buckman, ⁵Kent Thomas, ⁶Dale Sandler, ⁷Jane Hoppin, ¹Stella Koutros, ¹Gabriella Andreotti, ¹Jay Lubin, ¹Aaron Blair, ¹Laura Beane Freeman. ¹Occupational and Environmental Epidemiology Branch, National Cancer Institute, Bethesda, MD, USA; ²University of Iowa, Department of Epidemiology, Iowa City, IA, USA; ³National Institute for Occupational Safety and Health, Cincinnati, OH, USA; ⁴IMS, Calverton, MD, USA; ⁵National Exposure Research Laboratory, U. S. Environmental Protection Agency, Research Triangle, NC, USA; ⁶National Institute of Environmental Health Sciences, Research Triangle, NC, USA; ⁷North Carolina State University, Raleigh, NC, USA

10.1136/oemed-2014-102362.111

Objectives Farming and exposure to pesticides have been linked to non-Hodgkin lymphoma (NHL), and multiple myeloma (MM) in previous studies. We evaluated use of insecticides, fungicides and fumigants and risk of NHL, including MM and other NHL sub-types in the Agricultural Health Study, a US-based prospective cohort study.

Method A total of 527 cases occurred among 55 875 pesticide applicators from enrollment (1993–1997) through 2011 in Iowa and 2010 in North Carolina. Information on pesticide use, other agricultural exposures and other factors was obtained from questionnaires at enrollment and follow-up approximately five years later (1999–2005). Information from these questionnaires was used to create lifetime-days and intensity-weighted lifetime-days of pesticide use. Poisson regression and polytomous logit models were used to calculate relative risks (RR) and 95% confidence intervals (CI) to evaluate associations between 26 pesticides and NHL and five NHL-subtypes including multiple myeloma, while adjusting for potential confounding factors.

Results Statistically significant positive exposure-response trends occurred between overall NHL risk and lindane (p-trend = 0.004) and DDT (p-trend = 0.02). In addition, ever use of terbufos was associated with NHL overall (RR=1.2; CI=1.0–1.5), but with no exposure-response trend. In sub-type analyses, terbufos and DDT were associated with small cell lymphoma/chronic lymphocytic leukaemia/marginal cell lymphoma. In addition, lindane and diazinon were associated with follicular lymphoma and permethrin with MM although tests of homogeneity did not show significant differences in exposure-response among NHL-subtypes for any chemical.

Conclusions These findings are among the first to suggest links between DDT, lindane, permethrin, diazinon and terbufos and specific NHL subtypes.

0288 GENE-SPECIFIC DNA METHYLATION AS A VALUABLE TOOL FOR RISK ASSESSMENT: THE CASE OF OCCUPATIONAL EXPOSURE TO DIFFERENT VOC'S IN MEXICAN WORKERS

¹Octavio Jiménez-Garza, ²Andrea Baccarelli, ²Hyang-Min Byun, ³Giovanni Battista Bartolucci, ³Mariella Carrieri. ¹University of Guanajuato Campus León, Health Sciences Division, León, Guanajuato, Mexico; ²Harvard School of Public Health, Laboratory of Human Environmental Epigenomics, Boston, Massachusetts, USA; ³University of Padova, Department of Medicine, Padova, Italy

10.1136/oemed-2014-102362.112

Objectives To determine gene-specific methylation levels (promoter region) on genes from critical cellular pathways in persons occupationally exposed to a single volatile organic compound (VOC) or to a mixture of them

Method Workers from two tanneries, two shoe factories, and two gas stations were included (exposed groups to VOC's). We also included administrative workers (reference group), all of them from the city of León Guanajuato, México. In tannery workers we measured individual exposure levels to three different VOC's; for the rest of the groups we measured up to 7 VOC's. After exposure characterisation, we took blood samples and extracted DNA in order to determine, by PCR-pyrosequencing, methylation levels in genes involved in inflammation, DNA repair, oxidative stress and xenobiotic metabolism pathways.

Results Only toluene environmental levels were higher in tannery workers. Workers from the leather shoe factory showed the highest exposure levels for up to five different VOC's. There was no statistical significant difference in gene-specific methylation for tannery workers when compared to the control group. For the leather shoe factory workers, we found hypermethylation in the TNF α , SOD1 and TOP2A promoter regions compared to the control group or to other exposed groups. Gas station attendants showed hypermethylation for the IL6 gene compared to controls

Conclusions Occupational exposure to a mixture of VOC's has important effects on the methylation status of genes involved in inflammation, DNA repair and oxidative stress. These epigenetic changes, detectable at a pre-clinical stage, represent a valuable tool for performing an early risk assessment in these populations

0289 PREVENTING NEEDLESTICKS AND OTHER SHARPS INJURIES TO HOME CARE AIDES: RESULTS OF A SURVEY TO IDENTIFY HAZARDS DURING HOME VISITS

¹David Kriebel, ¹Natalie Brouillette, ¹Pia Markkanen, ¹Catherine Galligan, ¹Susan Sama, ¹Rebecca Gore, ²Angela Laramie, ¹Daniel Okyere, ¹Chuan Sun, ²Letitia Davis, ¹Margaret Quinn. ¹University of Massachusetts Lowell, Lowell, MA, USA; ²Massachusetts Department of Public Health, Boston MA, USA

10.1136/oemed-2014-102362.113

Objectives To follow up on our previous finding that home care (HC) aides have a substantial risk of injury with used sharp medical devices, this study's goal was to identify modifiable aspects of aide-client encounters during HC visits that increase risk of sharps injuries (SI).

Method A survey of 1249 HC aides was conducted in eastern Massachusetts. Approximately half the participants (634) were employed by HC agencies, and half (615) directly by HC clients and their families. A questionnaire gathered data on aides' most recent home visits, enabling quantification of hazardous working conditions. The specific investigation described here focused on understanding determinants of hazardous conditions likely to increase the risk of SI including: the aide finding used sharps