Editor's choice
Scan to access more
free content

ORIGINAL ARTICLE

Trends in mortality from occupational hazards among men in England and Wales during 1979–2010

E Clare Harris,¹ Keith T Palmer,¹ Vanessa Cox,¹ Andrew Darnton,² John Osman,³ David Coggon¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/oemed-2015-103336>).

¹MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK

²Statistics and Epidemiology Unit, Science Directorate, Health and Safety Executive, UK

³Centre for Occupational and Environmental Health, University of Manchester, Manchester, UK

Correspondence to Professor David Coggon, MRC Lifecourse Epidemiology Unit, Southampton General Hospital, Southampton SO16 6YD UK; dnc@mrc.soton.ac.uk

Received 22 September 2015

Revised 5 February 2016

Accepted 2 March 2016

Published Online First

14 March 2016

ABSTRACT

Objectives To monitor the impact of health and safety provisions and inform future preventive strategies, we investigated trends in mortality from established occupational hazards in England and Wales.

Methods We analysed data from death certificates on underlying cause of death and last full-time occupation for 3 688 916 deaths among men aged 20–74 years in England and Wales during 1979–2010 (excluding 1981 when records were incomplete). Proportional mortality ratios (PMRs), standardised for age and social class, were calculated for occupations at risk of specified hazards. Observed and expected numbers of deaths for each hazard were summed across occupations, and the differences summarised as average annual excesses.

Results Excess mortality declined substantially for most hazards. For example, the annual excess of deaths from chronic bronchitis and emphysema fell from 170.7 during 1979–1990 to 36.0 in 2001–2010, and that for deaths from injury and poisoning from 237.0 to 87.5. In many cases, the improvements were associated with falling PMRs (suggesting safer working practices), but they also reflected reductions in the numbers of men employed in more hazardous jobs, and declining mortality from some diseases across the whole population. Notable exceptions to the general improvement were diseases caused by asbestos, especially in some construction trades and sinonasal cancer in woodworkers.

Conclusions The highest priority for future prevention of work-related fatalities is the minority of occupational disorders for which excess mortality remains static or is increasing, in particular asbestos-related disease among certain occupations in the construction industry and sinonasal cancer in woodworkers.

INTRODUCTION

Monitoring trends in the burden of illness and injury attributable to work is an important element in the control of occupational health hazards.¹ It enables evaluation of the effectiveness of measures to reduce risk, and informs the prioritisation of future preventive strategies. The best methods for such monitoring vary according to the nature of the health outcome (eg, whether it is frequently fatal) and the confidence with which it can be ascribed to work in the individual case.

For some outcomes, the link to occupation can readily be established in the individual because they

What this paper adds

- It has been unclear to what extent improved working conditions in Britain over recent decades have reduced the burden of serious occupational disease, and which should now be the highest priorities for prevention.
- Analysis of data from death certificates indicates that during 1979–2010 excess mortality from most established occupational hazards declined substantially, most likely reflecting a combined effect of reduced employment in more hazardous jobs, improved health across the population as a whole, and safer working conditions.
- Notable exceptions to the general improvement are diseases caused by asbestos, especially in some construction trades and sinonasal cancer in woodworkers.
- Asbestos-related disease in the construction industry and sinonasal cancer in woodworkers remain priorities for future prevention of work-related mortality.

are specific to exposures in the workplace (eg, byssinosis, coal workers' pneumoconiosis), occur with very high relative risk in people who have undertaken certain jobs (eg, adenocarcinoma of the nose and nasal sinuses in furniture makers using hardwoods), or have distinctive clinical features when caused by work (eg, the temporal relationship of an injury to an occupational activity, demonstrable dermal sensitisation to an allergen encountered in the workplace). In these circumstances, the resultant burden can be assessed simply by counting attributable cases.

For other health outcomes, the relation to work is less specific, and cases that are caused by occupation are clinically indistinguishable from those that are not. For example, there are no distinctive clinical features of lung cancer that is caused by asbestos. In this situation, the attributable population burden can only be established epidemiologically—by comparing risks in people with and without the relevant exposure. Estimates may be derived by direct assessment of attributable fractions in representative populations, or indirectly by applying estimates of relative risk from one source to data

CrossMark

To cite: Harris EC, Palmer KT, Cox V, et al. *Occup Environ Med* 2016;**73**:385–393.

collected separately on the distribution of exposures in the population of interest.

National analyses of occupational mortality contribute importantly to this endeavour because they incorporate data on exposures and health outcomes for an entire population. They are particularly useful in the assessment of diseases and injuries with high fatality, but even where case fatality is lower, they may still provide helpful insights into trends over time, especially when there is little change in the proportion of cases that lead to death.

We have previously reported a national analysis of occupational mortality in England and Wales over two consecutive periods (1979–1990 and 1991–2000),² and have now extended that investigation with addition of data from 2001 to 2010. Our aim was to evaluate trends across the three time periods in mortality from established occupational hazards, and the extent to which changes reflected altered levels of risk in relevant occupations as opposed to changes in the numbers at risk in those jobs, or an overall decline in mortality across all occupations (eg, because of improved medical care). Of particular interest was whether previously persistent excess mortality attributable to asbestos and wood dust² had now started to fall.

METHODS

The analysis used data supplied by the Office for National Statistics (ONS) (previously the Office of Population Censuses and Surveys (OPCS)) on all deaths among men aged 20–74 years in England and Wales during 1979–2010 (but excluding 1981 when records were incomplete). Information from death certificates on age at death, underlying cause of death and last full-time occupation was used to derive proportional mortality ratios (PMRs) with associated 95% CIs for combinations of occupation and cause of death, the expected numbers being calculated with stratification by 5-year age band and social class (determined from the last full-time occupation and classified to seven levels), taking all occupations combined as the standard.

For this purpose, occupations were assigned to ‘job groups’ in a classification similar to that which had been used in our earlier analysis.² However, the coding schemes that were employed by ONS and OPCS had changed over time, and a few occupational categories that were distinguished from each other in one period were merged in another. To achieve compatibility across all periods, analyses for these occupations were based on the merged grouping, and for the period(s) in which the finer classification had been used, observed and expected numbers of deaths were derived separately for the subcategories and then summed to give combined values for the merged job group. Online supplementary table S1 sets out the definitions of the job groups analysed, according to the coding schemes that were applied in each period.

Similarly, cause of death categories were mostly as had been analysed previously,² but with a few adjustments because of changes between the 9th and 10th revisions of the International Classification of Diseases (ICD-9 and ICD-10; see online supplementary table S2). Again, where for consistency it was necessary to merge cause of death categories that had not been distinguished in all time periods, observed and expected numbers of deaths were derived separately for each subcategory and then aggregated to give values for the merged grouping. One important category in ICD-10, mesothelioma unspecified, had been included in ICD-9 with all unspecified neoplasms (which were much more numerous), and therefore could not be considered in our analysis of trends.

For each cause of death that was considered an established occupational hazard, we listed those job groups in which excess mortality might plausibly be expected—for example, cancer of the pleura in carpenters and joiners (as a consequence of exposure to asbestos), railway accidents in rail construction and maintenance workers. We started with the hazards and the job groups that had been classed as at risk in our previous analysis,² and reviewed the observed and expected numbers of deaths for those combinations of cause of death and job group during each of the three periods, 1979–1990, 1991–2000 and 2001–2010, and overall. For injury and poisoning (fatality from which is likely to be acute) and pneumonia in occupations exposed to metal fume (for which the excess risk resolves following cessation of exposure), we considered only deaths at normal working ages (20–64 years). For other hazards, we included deaths at 20–74 years. A few of the combinations of hazard and job group were excluded from further analysis because changes in classification meant that they could not be examined meaningfully, or there was no excess mortality in any of the three periods of study (see online supplementary table S3).

For each hazard and time period, we summed the observed and expected numbers of deaths across the job groups designated as being at risk, and divided the difference between the total observed and expected numbers by the length of the observation period to derive an estimate of annual excess mortality in the relevant period. To avoid bias, we applied this approach even where, in a particular job group and time period, the number of deaths observed was fewer than the number expected (ie, the excess was negative). For diseases that could occur only as a consequence of occupational exposures (eg, pneumoconiosis, farmers’ lung disease), we counted also the deaths in occupations that were not designated as being at risk, and included them in our estimate of excess mortality. Where a major excess of deaths was observed for a hazard, we noted the main job groups that contributed, and where there was a substantial change in excess mortality over the duration of the study, we checked to which job groups it applied, and whether it was driven more by changes in their PMRs, in the numbers of men at risk (indicated by the number of deaths from all causes in the job groups concerned), or in mortality from that cause across all occupations.

As in our previous analysis,² we did not attempt to evaluate trends in excess mortality from lung cancer that might be attributable to work. This was because of the major potential for confounding by smoking, the prevalence of which was expected to vary importantly between job groups and over time. Nor did we examine mortality from coronary heart disease since although it has been linked with both shift work³ and occupational stress,⁴ there were no job groups in which an excess of deaths could confidently be ascribed to such exposures.

RESULTS

The analysis was based on a total of 3 688 916 deaths, of which 1 564 981 occurred during 1979–1990, 1 199 234 in 1991–2000 and 924 701 in 2001–2010. Table 1 sets out the numbers of deaths in each time period, across all job groups combined, from the causes of death that were examined as occupational hazards. There were marked declines in mortality from some causes including tuberculosis, cancer of the scrotum, asthma, farmers’ lung disease, most types of pneumoconiosis, byssinosis and many categories of injury. On the other hand, increases were observed in deaths from viral hepatitis, cancer of the liver, other alcohol-related diseases, other and unspecified allergic pneumonitis and asbestosis.

Table 1 Causes of death analysed and numbers of deaths by time period in all job groups combined, men aged 20–74 years, England and Wales, 1979–2010

Cause of death	Number of deaths in all job groups combined at ages 20–74 years*			Per cent change in annual number of deaths 1979–1990 to 2000–2010
	1979–1980 1982–1990	1991–2000	2000–2010	
Infectious diseases				
Tuberculosis	3019	1433	945	–66
Viral hepatitis	544	592	998	102
Cancers				
Cancer of the oral cavity	3687	3283	3533	5
Cancer of the pharynx (specified)	2830	2566	3064	19
Cancer of the liver	5339	6472	8678	79
Cancer of the peritoneum	362	365	363	10
Cancer of the nose and nasal sinuses and middle ear	839	518	400	–48
Cancer of the larynx	4832	3962	3213	–27
Cancer of the pleura	2848	2894	2314	–11
Mesothelioma at other sites and unspecified combined			6114	
Other cancer of skin (excluding melanoma)	1219	910	722	–35
Cancer of the scrotum	85	29	16	–79
Urothelial cancer	18 403	14 184	10 691	–36
Other alcohol-related diseases				
Other alcohol-related diseases (excluding alcohol poisoning)	8216	14 530	26 701	257
Respiratory diseases				
Combined pneumonia (excluding bronchopneumonia)	3672	4410	4191	26
Chronic bronchitis and emphysema	85 096	54 941	38 955	–50
Asthma	5586	3441	1630	–68
Farmer's lung disease	66	34	17	–72
Other and unspecified allergic pneumonitis	24	36	68	212
Coal worker's pneumoconiosis	942	498	210	–75
Asbestosis	281	331	407	59
Silicosis	163	50	39	–74
Other pneumoconiosis combined	244	98	49	–78
Byssinosis	39	6	1	–97
Digestive diseases				
Cirrhosis (not specified as biliary)	5327	5533	6469	34
Pancreatitis	2785	2718	2758	9
Injury and poisoning				
Railway accidents combined	562	239	142	–72
Motor vehicle traffic accidents combined	20 428	11 952	10 154	–45
Off-road motor vehicle traffic accidents combined	476	261	132	–69
Animal transport accidents rider/passenger	53	26	20	–58
Water transport accidents	499	173	126	–72
Air transport accident	427	216	92	–76
Pesticide poisoning	11	5	5	–50
Poisoning by other gases and vapours combined	617	263	177	–68
Fall from ladder or scaffolding	606	294	208	–62
Fall from building	900	416	358	–56
Fall into hole	96	35	115	32
Other fall	428	291	171	–56
Slipping and tripping	121	91	74	–33
Fall unspecified	1040	877	1230	30
Injured by fire	1202	871	577	–47
Heat injury	14	10	5	–61
Injury by animals and plants combined	41	42	46	23
Injury by lightning	20	14	5	–73
Injury by falling, thrown or projected object	647	263	160	–73
Injury by being caught between objects	54	31	24	–51
Injury by machinery	907	341	89	–89
Injury by cutting and piercing instruments or objects	88	72	29	–64
Injury by explosion of pressure vessel	30	16	5	–82

Continued

Table 1 Continued

Cause of death	Number of deaths in all job groups combined at ages 20–74 years*			Per cent change in annual number of deaths 1979–1990 to 2000–2010
	1979–1980 1982–1990	1991–2000	2000–2010	
Injury by firearms	95	44	17	–80
Injury by explosive material	148	52	12	–91
Injury by hot substances	54	28	13	–74
Injury by electric current	452	255	144	–65
Homicide	1234	1038	1098	–2
War	28	3	1	–96

*For pneumonia, injury and poisoning, the numbers of deaths are at ages 20–64 years.

Table 2 lists the job groups that were considered a priori to be at risk of one or more of the identified occupational hazards, and gives the total number of deaths in each job group by time period. In most job groups, the number of deaths declined over the period of study, the fall being most marked in traditional manufacturing occupations, and in rail travel assistants and dockers. There was, however, a notable increase in deaths in professional athletes and sports officials. A reduction in deaths among electrical, energy, boiler operatives and attendants was largely offset by an increase for other transport, plant and machine operatives.

Table 3 summarises the trends in excess mortality from diseases caused by occupational exposure to dusts and fumes (here and in tables 4 and 5, some of the results for 1979–1990 and 1991–2000 differ slightly from those in our earlier paper² because of the aggregation of job groups that was necessary to achieve compatibility across the three time periods, and because in this study, deficits of observed vs expected deaths were counted as negative numbers rather than 0). The overall excess of deaths declined from an annual average of 414.8 during 1979–1990 to 272.6 in 1991–2000. The new category of deaths from mesothelioma at unspecified sites, which was distinguished only during 2001–2010, accounted for 135.9 excess deaths per year. In the same period, total excess mortality from the other causes of death showed a further decrease to 312.8–135.9=176.9 deaths per year.

The main contributors to the progressive decline in excess mortality were chronic bronchitis and emphysema, and coal workers' pneumoconiosis, for which the annual excesses of deaths fell from 170.7 to 36.0 and 85.6 to 21.0, respectively. These reductions were driven largely by the fall in the population of coal miners, as evidenced by a decrease in all-cause mortality from 47 249 deaths during 1979–1990 to 14 208 in 2001–2010 (table 2). However, as illustrated in online supplementary table S4, which shows a detailed breakdown of excess mortality for individual job groups, there was in addition, a small decrease in the PMR for chronic bronchitis and emphysema in coal miners from 144 to 128. For silicosis, farmers' lung disease and pneumonia, the decrease in excess mortality was also beyond that which could be explained by a fall in the numbers at risk. The lowering in excess occupational mortality from tuberculosis and asthma coincided with falls in the annual numbers of deaths from those diseases across all occupations by 66% and 68%, respectively (table 1), but there was also a reduction in the PMRs of the job group at highest risk of tuberculosis—mine (excluding coal) and quarry workers—from 431 to 172.

In contrast to the improving picture for most diseases caused by dusts and fumes, excess mortality from sinonasal cancer and

from diseases caused by asbestos remained fairly constant, or in the case of asbestosis, increased. The main job groups contributing to the rise in mortality from asbestosis were carpenters and joiners (18.9 excess deaths in 2001–2010 vs 3.8 in 1979–1991), plumbers (23.1 vs 9.8) and platers (13.4 vs 1.9; see online supplementary table S4). Mesothelioma at unspecified sites was only distinguished as a separate diagnostic category in ICD-10, but PMRs for the asbestos-exposed occupations during 2001–2010 correlated closely with those for cancer of the pleura (correlation coefficient 0.81, see online supplementary figure S1).

Over the entire study period, the calculated excess mortality among occupations designated as being at risk accounted for 88% of the 1650 deaths from coal workers' pneumoconiosis, 58% of 1019 deaths from asbestosis, 65% of 252 deaths from silicosis, 85% of 117 deaths from farmers' lung disease, and 52% of 9146 deaths from cancers of the peritoneum and pleura combined (see online supplementary table S4).

Trends in excess mortality from other diseases caused by work are summarised in table 4. Reductions in excess deaths were observed for non-melanoma skin cancer in occupations particularly exposed to sunlight, and for cancer of the scrotum in those with exposure to mineral oils, while for viral hepatitis in doctors and urothelial cancer in chemical workers, excess mortality was small with no clear trends across the study period (see online supplementary table S5). Publicans and bar staff (in the UK, publicans own or manage 'public houses' that are licensed to sell alcoholic drinks for consumption on the premises) experienced consistently elevated mortality from diseases caused by alcohol, but their annual excess of deaths from cirrhosis and pancreatitis declined over time, whereas that from cancers of the pharynx and liver increased (see table 4 and online supplementary table S5). There was a parallel rise in deaths from cancer of the liver in all occupations combined, but total mortality from cirrhosis and pancreatitis showed no overall decline across the study period (table 1).

Table 5 and online supplementary table S6 summarise patterns of mortality from injury and poisoning. The overall annual excess of deaths in occupations at risk fell from 237.0 during 1979–1990 to 137.4 in 1991–2000, and 87.5 in 2001–2010. This substantial decline was driven by reductions in most of the major categories of injury, including a notable fall in the excess of deaths from railway accidents among railway workers from 11.7 per year in 1979–1990 to 0.4 deaths per year during 2001–2010. One exception to the general downward trend was excess mortality from injury by animals and plants in farmers, which remained fairly constant at 1.0–1.4 deaths per year (see online supplementary table S6).

Table 2 Job groups analysed and numbers of deaths from all causes by time period, men aged 20–74 years, England and Wales, 1979–2010

Job group	Number of deaths from all causes combined at ages 20–74 years			Per cent change in annual number of deaths 1979–90 to 2000–10
	1979–1980 1982–90	1991–2000	2000–2010	
Marketing and sales managers combined	14 454	14 132	19 105	45
Vocational trainers, social scientists, etc	3828	3855	3415	–2
Doctors	4086	3263	2807	–24
Other health professions combined	2360	2431	1962	–9
Professional athletes, sports officials	482	624	1068	144
Chemical engineers and scientists	2554	2539	2018	–13
Other professional engineers	16 766	17 230	14 787	–3
Draughtspersons	6057	5076	3365	–39
Laboratory technicians	5795	3325	2331	–56
Aircraft flight deck officers	619	814	574	2
Seafarers	8600	6963	4780	–39
Other technicians	4593	6308	6276	50
Production and maintenance managers	27 591	21 689	14 224	–43
Managers in construction	7036	5578	5465	–15
Managers in transport, mining and energy industries	16 247	13 517	9562	–35
Butchers	9739	6364	4087	–54
Publicans and bar staff	17 415	12 446	10 346	–35
Farming and fishing combined	58 113	40 782	30 071	–43
Armed forces	10 037	8204	8266	–9
Police	7273	7426	6812	3
Fire service personnel	2805	2643	1920	–25
Sales representatives	24 234	15 869	10 839	–51
Other service personnel combined	40 181	34 143	23 204	–36
Rail travel assistants combined	9749	6331	1032	–88
Forestry workers	1459	1090	675	–49
Leather and related trades combined	6108	3189	1078	–81
Weavers and knitters combined	3386	1782	860	–72
Other textile processing operatives combined	10 185	2339	1419	–85
Chemical workers combined	18 146	9821	5117	–69
Bakers	5371	4020	1950	–60
Other food drink and tobacco process operatives nec combined	10 391	7330	5886	–38
Paper and wood machine operatives combined	8882	4946	3104	–62
Glass and ceramic workers combined	7846	4422	2628	–63
Coal miners combined	47 249	27 896	14 208	–67
Upholsterers	2620	1574	1265	–47
Carpenters and joiners	25 914	22 049	20 817	–12
Cabinet makers combined	3972	2822	2276	–37
Smiths and forge workers	2633	1570	836	–65
Moulders, core makers, die casters	4607	2198	909	–78
Electroplaters combined	1273	849	509	–56
Other metal manufacturers combined	12 224	7613	3825	–66
Metal machining setters and setter-operators combined	11 349	10 482	4582	–56
Metal working machine operatives combined	54 442	34 154	24 181	–51
Production fitters	49 781	38 535	23 220	–49
Motor mechanics, auto engineers combined	14 788	13 162	11 536	–14
Electricians, electrical maintenance fitters combined	20 719	18 381	21 253	13
Cable jointers, lines repairers	1728	1362	824	–48
Radio, TV and video engineers	2025	1698	1220	–34
Electrical engineers (not professional) combined	7803	6753	4836	–32
Plumbers, heating and ventilating engineers and related trades	16 019	14 797	14 046	–4
Sheet metal workers	7529	4806	2919	–57
Metal plate workers, shipwrights, riveters	5560	4120	2151	–57
Steel erectors	3517	3675	1837	–43
Scaffolders, riggers combined	3321	3484	2898	–4
Welding trades	11 783	10 414	8140	–24
Coach and vehicle body builders and repairers combined	2122	2324	2006	4

Continued

Table 2 Continued

Job group	Number of deaths from all causes combined at ages 20–74 years			Per cent change in annual number of deaths 1979–90 to 2000–10
	1979–1980 1982–90	1991–2000	2000–2010	
Painters and decorators	28 103	21 241	18 661	–27
Assemblers/lineworkers (electrical/electronic goods)	1031	1903	1363	45
Packers, sorters and testers combined	7947	15 376	8563	19
Bricklayers, masons combined	17 359	11 288	8412	–47
Roofers and glaziers	3267	3908	5610	89
Rail construction and maintenance workers	3020	1590	761	–72
Road construction workers and paviors	4947	3329	2979	–34
Other construction workers combined	48 602	45 154	59 230	34
Mine (excluding coal) and quarry workers	2580	1658	1279	–45
Rail transport operatives combined	2994	1707	3415	25
Railway engine drivers	4179	3836	1682	–56
Lorry drivers	58 475	54 545	48 106	–10
Mechanical plant drivers and operatives (earth moving and civil engineering)	2805	3736	3673	44
Crane drivers	8153	4887	2713	–63
Fork lift and mechanical truck drivers	6330	6967	6771	18
Storekeepers and warehousemen/women	46 739	28 467	19 928	–53
Dockers goods porters and slingers combined	14 154	9443	2829	–78
Electrical, energy, boiler operatives and attendants combined	9428	4393	1818	–79
Other transport, plant and machine operatives combined	3146	12 523	9418	229

Table 3 Excess mortality from diseases caused by occupational exposure to dusts and fumes, men aged 20–74 years,* England and Wales, 1979–2010

Cause of death	Exposure	1979–1980, 1982–1990			1991–2000			2001–2010		
		Deaths observed†	Deaths expected†	Annual excess†	Deaths observed†	Deaths expected†	Annual excess†	Deaths observed†	Deaths expected†	Annual excess†
Coal workers' pneumoconiosis	Coal mine dust	942		85.6	498		49.8	210		21.0
Asbestosis	Asbestos	281		25.5	331		33.1	407		40.7
Cancer of the peritoneum	Asbestos	178	94.7	7.6	168	93.6	7.4	173	100.3	7.3
Cancer of the pleura	Asbestos	1491	792.5	63.5	1495	836.5	65.8	1239	681.4	55.8
Mesothelioma at unspecified and other sites combined	Asbestos							3149	1790.3	135.9
Silicosis	Silica dust	163		14.8	50		5.0	39		3.9
Tuberculosis	Silica dust	88	60.8	2.5	42	24.5	1.7	23	15.0	0.8
Other pneumoconiosis combined	Various	244		22.2	98		9.8	49		4.9
Byssinosis	Textile dust	39		3.5	6		0.6	1		0.1
Farmers' lung disease	Spores in mouldy hay	66		6.0	34		3.4	17		1.7
Other and unspecified allergic pneumonitis	Various	7	0.9	0.6	3	1.1	0.2	8	1.8	0.6
Cancer of the nose and nasal sinuses and middle ear	Leather dust, wood dust	50	25.4	2.2	35	15.4	2.0	30	11.7	1.8
Combined pneumonia (excluding bronchopneumonia)	Metal fume	164	87.0	7.0	122	80.2	4.2	89	75.6	1.3
Chronic bronchitis and emphysema	Coal mine dust, silica dust, metal fume	7391	5513.8	170.7	3889	3041.7	84.7	2063	1702.6	36.0
Asthma	Various	156	122.1	3.1	142	93.7	4.8	44	34.3	1.0
Total		11 260	6697.2	414.8	6913	4186.7	272.6	7541	4413.0	312.8

*Deaths from pneumonia in relation to metal fume are at ages 20–64 years.

†For coal workers' pneumoconiosis, asbestosis, silicosis, other pneumoconiosis combined, byssinosis and farmers' lung disease, the number of deaths observed is for all occupations combined, and was used to calculate the annual excess. For all other hazards, the numbers of deaths observed and expected are for job groups designated as being at risk (see text on methods and online Supplementary table S4), and the excess rate was based on the difference between observed and expected.

Table 4 Excess mortality from diseases with other occupational causes: men aged 20–74 years, England and Wales 1979–2010

Cause of death	Exposure	1979–1980, 1982–1990			1991–2000			2001–2010		
		Deaths observed*	Deaths expected*	Annual excess*	Deaths observed*	Deaths expected*	Annual excess*	Deaths observed*	Deaths expected*	Annual excess*
Viral hepatitis	Hepatitis infection	14	3.2	1.0	8	2.7	0.5	10	3.0	0.7
Cancer of the oral cavity	Alcohol	117	42.6	6.8	110	35.3	7.5	109	43.1	6.6
Cancer of the pharynx	Alcohol	71	30.8	3.7	67	28.2	3.9	83	38.7	4.4
Cancer of the liver	Alcohol	112	69.1	3.9	125	81.8	4.3	165	107.6	5.7
Cancer of the larynx	Alcohol	119	45.4	6.7	100	36.1	6.4	75	34.6	4.0
Cirrhosis (not specified as biliary)	Alcohol	243	80.9	14.7	170	72.0	9.8	145	79.9	6.5
Pancreatitis	Alcohol	45	33.6	1.0	32	31.4	0.1	31	30.5	0.0
Other alcohol-related diseases (excluding alcohol poisoning)	Alcohol	458	125.4	30.2	449	185.9	26.3	662	319.0	34.3
Non-melanoma skin cancer	Sunlight	99	83.8	1.4	70	60.6	0.9	71	63.8	0.7
Urothelial cancer	Aromatic amines	224	208.8	1.4	145	116.8	2.8	85	68.7	1.6
Cancer of the scrotum	Mineral oils	24	7.7	1.5	7	2.1	0.5	2	0.9	0.1
Total		1526	731.1	72.3	1283	653.0	63.0	1438	789.8	64.8

*The numbers of deaths observed and expected are for job groups designated as being at risk (listed in online supplementary table S5), and the excess rate was based on the difference between observed and expected.

Table 5 Excess mortality from occupational injuries and poisoning: men aged 20–64 years, England and Wales 1979–2010

Cause of death	1979–1980, 1982–1990			1991–2000			2001–2010		
	Deaths observed*	Deaths expected*	Annual excess*	Deaths observed*	Deaths expected*	Annual excess*	Deaths observed*	Deaths expected*	Annual excess*
Railway accidents combined	134	4.8	11.7	25	1.5	2.4	5	0.7	0.4
Motor vehicle traffic accidents combined	2358	1561.8	72.4	1655	1034.1	62.1	1547	991.1	55.6
Off-road motor vehicle traffic accidents combined	190	76.6	10.3	118	43.6	7.4	75	21.4	5.4
Animal transport accidents	14	2.4	1.1	14	1.1	1.3	6	0.5	0.5
Water transport accidents	168	31.0	12.5	52	9.3	4.3	27	4.5	2.3
Air transport accidents	116	6.9	9.9	55	2.3	5.3	17	1.1	1.6
Pesticide poisoning	4	0.2	0.3	4	0.3	0.4	1	0.1	0.1
Poisoning by other gases and vapours combined	53	43.8	0.8	24	18.4	0.6	21	14.2	0.7
Fall from ladder or scaffolding	321	96.7	20.4	138	55.0	8.3	90	46.5	4.3
Fall from building	416	143.1	24.8	164	71.0	9.3	140	90.8	4.9
Fall into hole	22	6.1	1.4	6	2.4	0.4	7	8.2	−0.1
Other fall	91	45.3	4.2	55	26.9	2.8	22	16.6	0.5
Slipping and tripping	12	4.7	0.7	8	3.8	0.4	7	2.9	0.4
Fall unspecified	94	78.6	1.4	105	75.3	3.0	162	138.7	2.3
Injured by fire	8	1.8	0.6	5	1.8	0.3	1	1.3	0.0
Heat injury	4	0.2	0.3	2	0.1	0.2	0	0.1	0.0
Injury by animals and plants combined	17	2.2	1.3	16	1.6	1.4	11	1.3	1.0
Injury by lightning	3	0.9	0.2	3	0.6	0.2	0	0.0	0.0
Injury by falling, thrown or projected object	286	121.5	15.0	117	49.9	6.7	63	35.1	2.8
Injury by being caught between objects	15	7.0	0.7	11	3.3	0.8	5	2.1	0.3
Injury by machinery	521	201.8	29.0	187	73.3	11.4	45	21.4	2.4
Injury by cutting and piercing instruments or objects	22	8.1	1.3	15	7.3	0.8	9	5.3	0.4
Injury by explosion of pressure vessel	6	1.7	0.4	6	1.2	0.5	1	0.0	0.1
Injury by firearms	32	5.4	2.4	14	2.3	1.2	3	0.8	0.2
Injury by explosive material	55	20.7	3.1	14	7.0	0.7	3	1.8	0.1
Injury by hot substances	18	3.7	1.3	8	1.6	0.6	1	0.8	0.0
Injury by electric current	145	61.6	7.6	80	34.6	4.5	41	28.2	1.3
Homicide	16	5.8	0.9	7	6.2	0.1	4	4.6	−0.1
War	10	0.3	0.9	2	0.1	0.2	1	0.0	0.1
Total	5151	2544.5	237.0	2910	1535.9	137.4	2315	1440.0	87.5

*The numbers of deaths observed and expected are for job groups designated as being at risk (listed in online supplementary table S6), and the excess rate was based on the difference between observed and expected.

DISCUSSION

Our findings indicate that over recent decades, there has been a large and continuing fall in mortality from most of the occupational hazards considered. In many cases, this appears to have come about through improved safety, but it also reflects reductions in the numbers of men employed in more hazardous jobs such as coal mining, and falling mortality from some diseases across the whole population. Notable exceptions to the general improvement are diseases caused by asbestos, especially in some construction trades, sinonasal cancer in woodworkers, and injuries by animals and plants in farmworkers, for which there was still no clear decline in excess mortality.

As in our previous report,² we limited our analysis of trends to men. This was because in the early part of the study period, most women who died at age 20–74 years had spent only limited time in paid work, and during 1979–1990, only 30% had occupations recorded on their death certificates.⁵ Furthermore, with a few notable exceptions (eg, production of asbestos textiles⁶), the jobs in which they have been employed were not those in which excess mortality could confidently be attributed to occupational hazards. We plan to report on occupational mortality among women during 2001–2010 in a separate paper.

Our analysis had the advantage of covering all deaths nationally among men of working age over a prolonged period, which allowed trends to be evaluated with maximal statistical confidence. Against this, however, must be set a number of methodological limitations, which relate principally to the accuracy with which causes of death and occupational exposures could be determined, and the potential for confounding by non-occupational factors.

Cause of death was ascertained from death certificates, a source that is known not always to be accurate.⁷ Fewer errors might be expected for established occupational diseases such as pneumoconiosis, and for injury and poisoning, all of which are classed as unnatural causes of death and therefore are subject to more detailed investigation. However, diagnostic errors may have had a greater impact on, for example, cancer of the liver, which can be difficult to distinguish from hepatic metastasis of malignancies arising in other tissues. In general, misclassification of causes of death is likely to have been non-differential with respect to job group, biasing PMRs towards the null.

A further problem in assessing causes of death was the lack of specificity of some ICD categories. This applied particularly to mesothelioma at unspecified sites, which in ICD-9 was classed along with other cancers of unknown origin that were much more numerous. Analysis of data from 2001 to 2010, when ICD-10 was used, indicated that PMRs from unspecified mesothelioma correlated closely with those for pleural cancer (see online supplementary figure S1), and that excess numbers of deaths were some 2.5 times higher (see online supplementary table S4).

Incomplete diagnostic specificity was also a problem for deaths caused by injury and poisoning, the detailed circumstances of which were unknown. As a consequence, attribution to work could only be made with reasonable confidence when the category of injury was likely to be a particular hazard in a job group. Other injuries resulting from work will have been missed, and this explains why the annual excess deaths are substantially lower than the incidence of fatal occupational injuries assessed from closer investigation of individual cases. During 2001–2010, the UK Health and Safety Executive (HSE) recorded in excess of 200 fatal injuries per year to workers.⁸ Our estimate of 87.8 was limited to men, but unlike the HSE

statistics, it included deaths from road traffic accidents and in the armed forces.

Data on occupation were also derived from death certificates, and again will not have been completely accurate.⁹ Analysis of proportional mortality avoided the biases that can occur when different sources of occupational information are used for the numerator and denominator of death rates, but at the expense of possible bias if mortality in a job group from all causes combined was unusually low or high. In addition, as with causes of death, some occupational categories lacked desirable specificity. For example, because of changes in occupational classification, stone masons (who can have high exposures to silica) were grouped with bricklayers. The resultant 'dilution' will have led to lower PMRs, although it should not have biased estimates of excess numbers of deaths.

More important was the potential to miss long-term effects of hazardous exposures where men had changed employment and the last full-time occupation as recorded on the death certificate was not that in which the relevant exposure occurred. For example, as they approach retirement, some manual workers may move to jobs that are less physically demanding, and others may have changed their employment as some of the more hazardous industries contracted. Some indication of magnitude of this underascertainment can be gleaned from the findings for disease that are specific to occupational exposures. For example, coal mining occupations were reported for only 88% of deaths from coal workers' pneumoconiosis, and only 65% of deaths from silicosis occurred in men whose last occupation was classed as potentially exposed to silica. For mesothelioma, which can develop as a consequence of relatively short exposures to asbestos many years in the past, the underascertainment because of changes in employment may have been even greater. Over the full study period, excess mortality in the occupations designated as being at risk accounted for only 52% of all deaths from cancers of the peritoneum and pleura, whereas the true attributable fraction is likely to have been closer to 85%.¹⁰

Confounding by non-occupational factors will have been reduced by the adjustment of expected numbers of deaths for social class. However, it remains a potential problem for diseases caused by smoking, the prevalence of which can vary importantly between occupations, even within the same social class. The relative risk from smoking, and therefore the scope for confounding, is particularly high for lung cancer, and trends in occupational lung cancer, which require more complex assessment, will be considered in a separate report. Confounding may also have caused us to overestimate the burden of alcohol-related diseases attributable to work in bars. Some of the diseases (notably cancers of the oral cavity, pharynx and larynx) are also caused by smoking, which may be more prevalent in bar workers. Furthermore, while ready access to alcohol poses an important hazard in such employment, there may also be a tendency for heavy drinkers to take on jobs in bars selectively.

Overall, the effect of the various limitations discussed will have been a net underascertainment of the burden of mortality attributable to occupation. However, we think it unlikely that many of the estimates for occupational diseases will have been out by a factor of >3. Bearing in mind this scope for error, the estimates for sinonasal cancer in workers exposed to wood and leather dust can be compared with those derived by Slack *et al*¹¹ using different methodology. Applying risk estimates derived from published studies to estimates of the proportion of the national population ever exposed, they calculated population attributable fractions for mortality from sinonasal cancer in men during 2005 of 0.1533 (95% CI 0.0594 to 0.2912) for wood

dust and 0.0668 (95% CI 0.0328 to 0.1231) for leather dust. If applied to the average annual number of deaths from sinonasal cancer among men aged 20–74 during 2001–2010, which was 40 (see [table 1](#)), these figures give annual excess numbers of deaths of 6.1 (95% CI 2.4 to 11.6) for wood dust and 2.7 (95% CI 1.3 to 4.8) for leather dust. In comparison, we estimated annual excesses of 1.9 deaths for all woodworkers combined, and 0 for leather workers. Moreover, across all of 1979–2010, the annual excess of deaths in leather workers was less than 0.1. If extrapolated to this period, the lower 95% confidence limit of Slack's estimate would correspond to a cumulative excess of $1.3 \times 31 = 40.3$ deaths, whereas we observed only 8 deaths in total with 5.1 expected (see online supplementary table S4). While the two estimates for wood dust might be compatible, those for leather dust are not, and call into question the accuracy of the figure calculated by Slack.

While the inevitable shortcomings of our method are a source of uncertainty when estimating the absolute burden of mortality caused by work, the resultant biases are unlikely to have varied much over the 32-year period of study. Thus, we can be fairly confident that overall, excess mortality from the occupational hazards examined declined substantially in England and Wales between 1979 and 2010. For deaths from injury and poisoning, this conclusion is supported by other data collected independently.⁸ For some hazards, the fall in mortality has been a consequence of changing patterns of employment or a general reduction in mortality across all occupations. However, there is strong evidence also of improvements in safety.

The highest priority for future prevention of work-related fatalities is the minority of occupational disorders for which excess mortality remains static or is increasing. The continuing burden of disease caused by asbestos has been noted before,² and patterns of risk by birth cohort, including for asbestosis, suggest that reductions in exposure to asbestos since the 1970s have brought benefits.¹² However, there is still more to be achieved, with a special focus needed on certain trades in the construction industry. A useful starting point in planning further action might be to analyse patterns of incidence or mortality by birth cohort separately for relevant occupations. An absence of decline in recent birth cohorts would be a particular cause for concern. Also a priority is the persistent excess of sinonasal cancer in woodworkers. This was highlighted in our previous report, and data for the most recent 10-year period again show

no improvement. It is a well-established hazard, and should be manageable through controls on exposure to hardwood dust, particularly in the furniture industry.

Acknowledgements We thank the staff of the Office for National Statistics, who provided us with the data files for our analysis, and Stefania d'Angelo, who helped with part of the statistical analysis.

Contributors ECH contributed to the design of the study, jointly carried out the statistical analysis and jointly wrote the first draft of the paper. KTP contributed to the design of the study and to revision of the draft paper. VC jointly carried out the statistical analysis and contributed to revision of the draft paper. AD and JO contributed to the design of the study and to revision of the draft paper. DC led the design of the study, oversaw the statistical analysis, and jointly wrote the first draft of the paper.

Funding Medical Research Council. 10.13039/5011000002 65. MRC_MC_UP_A620_1018. MRC_MC_UU_12011/5.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data used in the analysis can be made available to other researchers subject to agreement from the Office for National Statistics.

REFERENCES

- 1 't Mannetje A, Pearce N. Occupational mortality studies: still relevant in the 21st century. *Occup Environ Med* 2010;67:802–3.
- 2 Coggon D, Harris EC, Brown T, et al. Work-related mortality in England and Wales, 1979–2000. *Occup Environ Med* 2010;67:816–22.
- 3 Vyas MV, Garg AX, Iansavichus AV, et al. Shift work and vascular events: systematic review and meta-analysis. *BMJ* 2012;345:e4800.
- 4 Glozier N, Tofler GH, Colquhoun DM, et al. Psychosocial risk factors for coronary heart disease. *Med J Aust* 2013;199:179–80.
- 5 Inskip H, Coggon D, Winter P, et al. Occupational mortality of women. In: Drever F, ed. *Occupational health: decennial supplement*. London: HMSO, 1995:44–61.
- 6 Newhouse ML, Berry G, Wagner JC. Mortality of factory workers in east London 1933–80. *Br J Indust Med* 1985;42:4–11.
- 7 Swift B, West K. Death certification: an audit of practice entering the 21st century. *J Clin Pathol* 2002;55:275–9.
- 8 Health and Safety Executive. Statistics on fatal injuries in the workplace in Great Britain, 2015. <http://www.hse.gov.uk/statistics/pdf/fatalinjuries.pdf>
- 9 Alderson MR. Some sources of error in British occupational mortality data. *Br J Indust Med* 1972;29:245–54.
- 10 Rake C, Gilham C, Hatch J, et al. Occupational, domestic and environmental mesothelioma risks in the British population: a case-control study. *Br J Cancer* 2009;100:1175–83.
- 11 Slack R, Young C, Rushton L, British Occupational Cancer Burden Study Group. Occupational cancer in Britain: nasopharynx and sinonasal cancers. *Br J Cancer* 2012;107(Suppl):S49–55.
- 12 Darnton A, Hodgson J, Benson P, et al. Mortality from asbestosis and mesothelioma in Britain by birth cohort. *Occup Med* 2012;62:549–52.