

Appendix Table 1: Survey wave in which a participant was diagnosed with a chronic condition and Province resided. Weighted N includes all participants with and without diagnosis in each model.

	Arthritis		Back pain		Diabetes		Hypertension		Heart disease	
	Weighted N	Diagnosed (%)	Weighted N	Diagnosed (%)	Weighted N	Diagnosed (%)	Weighted N	Diagnosed (%)	Weighted N	Diagnosed (%)
Survey wave diagnosed										
1996	1045	15.9	1420	17.4	121	18.2	570	14.8	143	17.1
1998	598	20.4	895	19.4	126	19.7	458	15.5	144	23.1
2000	598	17.3	684	20.3	120	19.8	450	15.4	131	10.1
2002	590	20.5	574	18.3	174	12.3	435	18.6	134	24.3
2004	464	26.4	469	17.7	164	14.3	448	21.1	115	19.5
2006	274	25.5	364	17.9	206	12.4	403	19.9	109	19.9
2008	305	24.0	249	17.9	174	21.1	356	25.0	107	15.4
Province*										
Newfoundland and Labrador	57	21.9	68	16.0	14	21.4	49	20.2	11	19.0
Prince Edward Island	21	20.1	23	15.7	6	22.9	14	28.9	4	16.5
Nova Scotia	109	30.0	128	17.6	29	17.1	82	22.3	24	25.7
New Brunswick	91	21.0	114	14.4	25	17.7	73	18.4	22	25.3
Quebec	925	14.4	1146	17.4	265	13.1	741	19.6	189	19.1
Ontario	1524	22.5	1772	17.5	426	19.0	1203	17.2	356	21.3
Manitoba	149	22.9	182	16.5	37	18.3	118	23.0	27	10.9
Saskatchewan	127	24.1	156	20.8	28	18.0	117	23.4	25	23.0
Alberta	389	22.9	479	21.7	118	16.3	320	16.2	92	15.9
British Columbia/Territories	481	16.4	587	22.0	137	12.6	403	16.1	133	10.1

Notes: * = Measured at Time 0, when respondents were employed and not diagnosed with any health condition; † = Work hours in wave previous to disease diagnosis; ‡ = Working status at current wave.