Oral Presentation

Risk Assessment

0362 RISK OF PLEURAL MM AND RESIDUAL ASBESTOS BURDEN IN THE LUNG: A RETROSPECTIVE CASE-CONTROL STUDY

^{1,2}Paolo Girardi*, ⁴Anna Somigliana, ⁵Pietro Gino Barbieri, ³Enzo Merler. ¹Venetian Mesothelioma Registry, Occupational Health Unit, Padua, Italy; ²SER Epidemiological Department, Veneto Region, Padua, Italy; ³Promotion and Development of Hygiene and Public Health, Venice, Italy; ⁴Lombardy Environmental Protection Agency (ARPA), Centre of Electronic Microscopy, Milan, Italy; ⁵Mesothelioma Register of Brescia, Occupational Health Unit, Local Health Authority, Brescia, Italy

10.1136/oemed-2017-104636.297

Introduction Results of Malignant Pleural Mesothelioma (MPM) occurrence (mortality and incidence) by cumulative exposure dose clearly showed a proportional relation of MPM risk with dose, confirmed among studies by fibre burden. We evaluated the association between residual fibre content and MPM risk by circumstance of asbestos exposure.

Methods and materials Lung samples obtained from pleuro-pneumonectomies or autopsies (349 MPMs, and 41 controls) among subjects investigated for probability and circumstance of asbestos exposure were examined through Scanning Electron Microscopy; 291 cases had an occupational asbestos exposure, 38 MPMs a non-occupational exposure (familiar or environmental), whereas among 20 MPM an asbestos exposure was not identified. The MPM risk was evaluated by means of Odds Ratio (OR).

Results The residual asbestos fibre burden was higher among MPMs occupationally exposed (Geometric Mean:2.10 Million fibres/gram of dried tissue; 95% CI:1.5–2.58) in comparison with non-occupational (GM:0.66 Mff/gdt; 95% CI:0.47–0.95) or with unknown exposures (GM:0.59 Mff/gdt; 95% CI:0.34–1.03) and controls (GM:0.26 Mff/gdt; 95% CI:0.20–0.34). Among occupationally exposed, the MPM risk increased according to the asbestos fibre burden reaching an OR of 36.8 (95%CI:11.9–113.5) for concentrations higher than 1 Mff/g dt, compared to the reference level (<0.25 Mff/gdt). Higher ORs were observed at any concentration of amphibole fibres in comparison those for chrysotile fibres.

Conclusions The MPM risk was strongly associated to the residual asbestos fibre lung burden. The MPM risk due to non-occupational exposure shows a magnitude comparable with that with unknown asbestos exposures. The residual lung burden of chrysotile is strongly influenced by clearance and time since exposures ceased.

Oral Presentation

Other

0363 OCCUPATIONAL EXPOSURE TO HIGH FREQUENCY ELECTROMAGNETIC FIELDS AND RISK OF BRAIN TUMOURS IN THE INTEROCC STUDY

^{1,3}Javier Vila*, ^{1,4}Michelle C Turner, ^{1,2}Esther Gracia-Lavedan, ^{1,2}Jordi Figuerola, ⁵Joseph D Bowman, ⁶Laurel Kincl, ⁷Lesley Richardson, ⁸Geza Benke, ⁹Sarah Fleming, ¹⁰Martine Hours, ⁴Daniel Krewski, ¹¹Dave McLean, ¹²Marie-Elise Parent, ^{13,14}Siegal Sadetzki, ¹⁵Klaus Schlaefer, ¹⁵Brigitte Schlehofer, ¹⁶Joachim Schüz, ⁷Jack Siemiatycki, ^{17,18}Martie van Tongeren, ^{1,3}Elisabeth Cardis. ¹Barcelona Institute for Global Health (ISGlobal), Barcelona, Spain; ²Universitat Pompeu Fabra (UPF), Barcelona, Spain; ³CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain; ⁴McLaughlin Centre for Population Health Risk Assessment, University of Ottawa, Ottawa, Canada; ⁵National Institute for Occupational Safety and Health (NIOSH), Ohio, USA; ⁶Oregon State University (OSU), Corvallis, Oregon, USA; ⁷University of Montreal Hospital Research Centre (CRCHUM), Montreal, Canada; ⁸School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia; ⁹Faculty of Medicine and Health, University of Leeds, Leeds, UK; ¹⁰Université de Lyon, Institut national de Veille Sanitaire, Unité Mixte de Recherche Épidémiologique et de Surveillance Transports Travail Environnement, Lyon, France; ¹¹Massey University, Wellington, New Zealand; ¹²INRS-Institut Armand-Frappier, Université du Québec, Laval, Canada; ¹³Cancer and Radiation Epidemiology Unit, Gertner Institute, Chaim Sheba Medical Centre, Tel-Hashomer, Israel; ¹⁴Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; ¹⁵German Cancer Research Centre, Heidelberg, Germany; ¹⁶International Agency for Research on Cancer (IARC), Section of Environment and Radiation, Lyon, France; ¹⁷Institute of Occupational Medicine (IOM), Edinburgh, UK; ¹⁸Centre for Occupational and Environmental Health, Centre for Epidemiology, University of Manchester, Manchester, UK; ¹⁹INTEROCC Study Group, Barcelona, Spain

10.1136/oemed-2017-104636.298

Introduction The International Agency for Research on Cancer classified radiofrequency (RF) electromagnetic fields (EMF) as *possibly carcinogenic* based on limited evidence in human studies of cell phone use and in animal experiments, while occupational evidence was found *inadequate* due in part to limitations in exposure assessment. This study assesses possible associations between occupational exposure to RF or intermediate frequency (IF) EMF in INTEROCC participants using novel exposure assessment methodologies.

Methods A plausible index of cumulative exposure to RF and IF EMF was calculated using a source-exposure matrix and detailed interviews on work with or nearby EMF sources, both overall and in specific exposure time windows. Conditional logistic regression was used to investigate associations with glioma and meningioma risk.

Results Only ~10% (n=769) of participants (n=7,330) were ever exposed to RF and ~1% (n=44) to IF EMF sources. Overall, there was no positive association between exposure to

RF or IF EMF and glioma or meningioma risk, and the majority of odds ratios (ORs) were <1.0. Some elevated ORs were observed in the highest exposure quantile (90th) for both RF electric fields and IF magnetic fields in the 1–4 year exposure time window for glioma, but were not statistically significant.

Conclusions Lifetime occupational RF and IF exposures based on our index were not associated with the risk of glioma or meningioma. Further work should include more exposed participants and examine alternative exposure or dose indices such as those incorporating thresholds on biological effects or combinations of static and RF magnetic fields.

Oral Presentation

Cancer

0364 RISK OF PROSTATE CANCER IN FIREFIGHTERS: A META-ANALYSIS

^{1,2}Jeavana Sritharan*, ³Marie-Elise Parent, ¹Manisha Pahwa, ^{1,4}Shelley Harris, ⁴Donald Cole, ^{1,4}Paul Demers. ¹Occupational Cancer Research Centre, Cancer Care Ontario, Toronto, Ontario, Canada; ²Institute of Medical Science, University of Toronto, Toronto, Ontario, Canada; ³INRS-Institut Armand-Frappier, University of Quebec, Laval, Quebec, Canada; ⁴Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada

10.1136/oemed-2017-104636.299

Objectives Although epidemiological studies have generally shown inconsistent findings for prostate cancer risk and occupation, some associations with firefighting have been observed. Firefighters are exposed to harmful substances that include known carcinogens. This meta-analysis aimed to synthesise the available published evidence on prostate cancer risk in firefighters.

Methods A comprehensive search of PubMed and Web of Science was conducted for all epidemiological studies on firefighting and prostate cancer published in English from 1980–2017. Studies that reported on risk estimates for incidence and/or mortality with corresponding ever/never employment were extracted from included studies. Meta risk estimates (mRE) were determined in a random effects model and between-study heterogeneity was evaluated based on I^2 values.

Results Of the 101 studies retrieved, 25 firefighter studies were included in this analysis. Ever employment as a firefighter, based on results from all studies, was associated with a 15% increase in prostate cancer risk (95% CI 1.04–1.27, $I^2=71\%$). By study design, the mRE was 1.14 (95% CI 1.03–1.26, $I^2=67\%$) for 18 cohort studies and 1.22 (95% CI 0.90–1.67, $I^2=82\%$) for 5 case-control studies. Incidence studies had an mRE of 1.17 (95% CI 1.07–1.28, $I^2=74\%$) and mortality studies had an mRE of 1.12 (95% CI 0.92–1.36, $I^2=50\%$). For linkage and registry-based studies, an mRE of 1.19 (95% CI 1.05–1.33, $I^2=61\%$) was observed. Estimates showed moderate to substantial heterogeneity.

Conclusions This analysis showed evidence of a small excess risk of prostate cancer in firefighters across study designs. Further evaluation of heterogeneity, potential biases, and quality assessment are currently underway.

Poster Presentation

Burden of Disease

0365 ESTIMATION OF THE BURDEN OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE DUE TO OCCUPATION IN GREAT BRITAIN

¹Sally Hutchings*, ¹Lesley Rushton, ²Steven Sadhra, ^{3,4}David Fishwick. ¹Imperial College London, Department of Epidemiology and Biostatistics, London, UK; ²Occupational and Environmental Medicine, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK; ³Health and Safety Laboratory, Buxton, UK; ⁴Infection, Immunity and Cardiovascular Disease, University of Sheffield, Sheffield, UK

10.1136/oemed-2017-104636.300

Introduction We have estimated the burden of COPD for deaths and prevalence due to occupational exposures in Britain using published literature and available national data plus a population based job exposure matrix (ACEJEM). The prevalence estimate avoids underestimation of the burden as estimates of COPD deaths, or hospitalisations, only account for the most serious cases.

Methods We used population attributable fractions (PAFs), requiring estimates of the population exposed to airborne pollutants at work, derived from 2001 Census data by job code matched to a JEM giving proportions exposed and average levels of exposure to pollutants including vapours, gases, dusts, fumes (VGDF) by UK SOC2000 job code. Exposures were matched to the best currently available population-based risk estimates. We estimated prevalent COPD PAFs using a novel age-time window approach to estimate prevalent numbers exposed, and deaths from COPD which requires numbers ever exposed were estimated using a risk exposure period methodology, taking into account age.

Results The PAF for prevalent COPD due to occupational exposure to VGDF was 18.7% for men, 8.0% for women and 11.5% overall giving 22 700 cases in men and 13 400 in women. Equivalent PAFs for COPD deaths, based on exposure 30+ years before 2010, were 18.3%/6.1%/12.3% (2,480 deaths in men, 790 in women). Prevalent PAF estimates which included a 30 year onset lag and mortality PAFs assuming 0–80 year latencies produced very similar results.

Conclusions The novel approach, making best use of available data, provides burden estimates which may inform strategies for reducing workplace COPD risks.

Oral Presentation

Burden of Disease

0366 COMPARING ALTERNATIVE METHODS OF ESTIMATING FUTURE DISEASE BURDEN DUE TO OCCUPATION

¹Sally Hutchings*, ¹Lesley Rushton, ²Martie van Tongeren, ³Damien McElvenny. ¹Imperial College London, Department of Epidemiology and Biostatistics, London, UK; ²University of Manchester, Manchester, UK; ³Institute of Occupational Medicine, Edinburgh, UK

10.1136/oemed-2017-104636.301