RF or IF EMF and glioma or meningioma risk, and the majority of odds ratios (ORs) were <1.0. Some elevated ORs were observed in the highest exposure quantile (90th) for both RF electric fields and IF magnetic fields in the 1–4 year exposure time window for glioma, but were not statistically significant.

Conclusions Lifetime occupational RF and IF exposures based on our index were not associated with the risk of glioma or meningioma. Further work should include more exposed participants and examine alternative exposure or dose indices such as those incorporating thresholds on biological effects or combinations of static and RF magnetic fields.

Oral Presentation

Cancer

0364 RISK OF PROSTATE CANCER IN FIREFIGHTERS: A META-ANALYSIS

^{1,2}Jeavana Sritharan*, ³Marie-Elise Parent, ¹Manisha Pahwa, ^{1,4}Shelley Harris, ⁴Donald Cole, ^{1,4}Paul Demers. ¹Occupational Cancer Research Centre, Cancer Care Ontario, Toronto, Ontario, Canada; ²Institute of Medical Science, University of Toronto, Toronto, Ontario, Canada; ³INRS-Institut Armand-Frappier, University of Quebec, Laval, Quebec, Canada; ⁴Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada

10.1136/oemed-2017-104636.299

Objectives Although epidemiological studies have generally shown inconsistent findings for prostate cancer risk and occupation, some associations with firefighting have been observed. Firefighters are exposed to harmful substances that include known carcinogens. This meta-analysis aimed to synthesise the available published evidence on prostate cancer risk in firefighters.

Methods A comprehensive search of PubMed and Web of Science was conducted for all epidemiological studies on firefighting and prostate cancer published in English from 1980–2017. Studies that reported on risk estimates for incidence and/or mortality with corresponding ever/never employment were extracted from included studies. Meta risk estimates (mRE) were determined in a random effects model and between-study heterogeneity was evaluated based on I² values.

Results Of the 101 studies retrieved, 25 firefighter studies were included in this analysis. Ever employment as a firefighter, based on results from all studies, was associated with a 15% increase in prostate cancer risk (95% CI 1.04–1.27, I²=71%). By study design, the mRE was 1.14 (95% CI 1.03–1.26, I²=67%) for 18 cohort studies and 1.22 (95% CI 0.90–1.67, I²=82%) for 5 case-control studies. Incidence studies had an mRE of 1.17 (95% CI 1.07–1.28, I²=74%) and mortality studies had an mRE of 1.12 (95% CI 0.92–1.36, I²=50%). For linkage and registry-based studies, an mRE of 1.19 (95% CI 1.05–1.33, I²=61%) was observed. Estimates showed moderate to substantial heterogeneity.

Conclusions This analysis showed evidence of a small excess risk of prostate cancer in firefighters across study designs. Further evaluation of heterogeneity, potential biases, and quality assessment are currently underway.

Poster Presentation

Burden of Disease

0365 ESTIMATION OF THE BURDEN OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE DUE TO OCCUPATION IN GREAT BRITAIN

¹Sally Hutchings*, ¹Lesley Rushton, ²Steven Sadhra, ^{3,4}David Fishwick. ¹Imperial College London, Department of Epidemiology and Biostatistics, London, UK; ²Occupational and Environmental Medicine, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK; ³Health and Safety Laboratory, Buxton, UK; ⁴Infection, Immunity and Cardiovascular Disease, University of Sheffield, Sheffield, UK

10.1136/oemed-2017-104636.300

Introduction We have estimated the burden of COPD for deaths and prevalence due to occupational exposures in Britain using published literature and available national data plus a population based job exposure matrix (ACEJEM). The prevalence estimate avoids underestimation of the burden as estimates of COPD deaths, or hospitalisations, only account for the most serious cases.

Methods We used population attributable fractions (PAFs), requiring estimates of the population exposed to airborne pollutants at work, derived from 2001 Census data by job code matched to a JEM giving proportions exposed and average levels of exposure to pollutants including vapours, gases, dusts, fumes (VGDF) by UK SOC2000 job code. Exposures were matched to the best currently available population-based risk estimates. We estimated prevalent COPD PAFs using a novel age-time window approach to estimate prevalent numbers exposed, and deaths from COPD which requires numbers ever exposed were estimated using a risk exposure period methodology, taking into account age.

Results The PAF for prevalent COPD due to occupational exposure to VGDF was 18.7% for men, 8.0% for women and 11.5% overall giving 22 700 cases in men and 13 400 in women. Equivalent PAFs for COPD deaths, based on exposure 30+ years before 2010, were 18.3%/6.1%/12.3% (2,480 deaths in men, 790 in women). Prevalent PAF estimates which included a 30 year onset lag and mortality PAFs assuming 0–80 year latencies produced very similar results.

Conclusions The novel approach, making best use of available data, provides burden estimates which may inform strategies for reducing workplace COPD risks.

Oral Presentation

Burden of Disease

0366 COMPARING ALTERNATIVE METHODS OF ESTIMATING FUTURE DISEASE BURDEN DUE TO OCCUPATION

¹Sally Hutchings*, ¹Lesley Rushton, ²Martie van Tongeren, ³Damien McElvenny. ¹Imperial College London, Department of Epidemiology and Biostatistics, London, UK; ²University of Manchester, Manchester, UK; ³Institute of Occupational Medicine, Edinburgh, UK

10.1136/oemed-2017-104636.301

Occupational burden of disease studies to identify priorities for regulatory interventions, and to estimate the impact of these interventions, are becoming more frequent. There are various ways to estimate burden of disease in a population, each method's appropriateness depends on available data and the research question to be answered. We review the available methods, and assess their usefulness for predicting future occupational disease burden and for testing population-level interventions.

There are several approaches to disease burden estimation: Population Attributable Fraction (PAF) applied to incident or prevalent disease cases, Lifetime Risk/Future Excess Fraction and Disease Projection (Age-Period-Cohort and structural model/'g' formula) approaches. The PAF takes into account past exposures, with intervention effects estimated from the point of intervention onwards. Lifetime risk estimates disease that workers, currently exposed to a disease-causing agent, will contract in their lifetimes; no latency assumptions are required, with interventions assumed to take effect immediately. Disease projection methods use current and past disease rates to predict rates into the future. The 'g-formula', a generalised regression model, is used to estimate the joint distribution of outcome and risk factors; Monte Carlo methods can then simulate disease risk following intervention, compared to no intervention. The methods are tested on examples of occupational disease/exposure pairings, including lung cancer from respirable crystalline silica exposure and COPD from vapour, gases, dust and fume exposure, and results are compared.

The results will provide critical information on the sensitivity of some examples of occupational burden of disease and health impact studies on the modelling approach.

Poster Presentation

Respiratory

0367 THE DIAGNOSIS OF PNEUMOCONIOSIS USING GAS CHROMATOGRAPHY/MASS SPECTROMETRY

^{1,2}Hsiao-Yu Yang*, ³Ruei-Hao Shie, ¹Che-Jui Chang, ^{1,2}Pau-Chung Chen. ¹*Institute of Occupational Medicine and Industrial Hygiene, National Taiwan University College of Public Health, Taipei, Taiwan;* ²*Department of Environmental and Occupational Medicine, National Taiwan University Hospital, Taipei, Taiwan;* ³*Green Energy and Environmental Research Laboratories, Industrial Technology Research Institute, Hsinchu, Taiwan*

10.1136/oemed-2017-104636.302

Breath metabolites could reflect pathogenic events at the respiratory level and might be used to screen for the presence of occupational environmental lung diseases. The objective of this study was to estimate the accuracy of breath analysis for the diagnosis of pneumoconiosis using gas chromatography/mass spectrometry (GC/MS). A prospective study was design. We screened 200 subjects from a cohort of stone workers and their family members. Diagnosis of pneumoconiosis was made based on occupational history, the presence of abnormal pulmonary function, and parenchymal abnormalities consistent with pneumoconiosis on chest X-rays. After excluding subjects with asthma and those using steroids or taking nonsteroidal anti-inflammatory drugs on the examination day, we conducted a case-control study that

enrolled 25 cases of pneumoconiosis and 154 controls. Breath and environmental air samples were collected and analysed by GC/MS. After subtraction of environmental volatile organic compounds (VOCs), the concentrations of VOCs in breath were used to discriminate between pneumoconiosis and controls. The discrimination accuracy was validated by both leave-one-out cross-validation and 5-fold cross-validation. The results showed that 94.4% of subjects were accurately classified and the cross-validated accuracy was 88.8%. The area under curve (AUC) for the diagnosis of pneumoconiosis was 0.9 (95% CI=0.8–1.0). Mean AUC of 5-fold cross-validation was 0.9. Breath test might have potential in the screening for pneumoconiosis; however, a multi-centre study is warranted to establish a reliable model and all procedures must be standardised before clinical application.

Oral Presentation

Methodology

0368 IMPROVING INCIDENCE RATES OF WORK-RELATED ILL-HEALTH CALCULATED FROM GENERAL PRACTICE

Louise Hussey*, Matthew Gittins, Roseanne McNamee, Raymond Agius. *The University of Manchester, Manchester, UK*

10.1136/oemed-2017-104636.303

Introduction The calculation of incidence rates of work-related ill-health (WRIH) is essential when assessing employment sectors at risk. The Health and Occupational Reporting network in General Practice (THOR-GP) collects information on WRIH from approximately 250 GPs. To calculate rates, GP numerator data has to be divided by a compatible denominator.

Aims To formulate the best methods of determining incidence rates from THOR-GP data

Methods The THOR-GP denominator is the registered population of all participating GPs' practices. Census data linked to practice postcodes were used to establish how the population was employed; practice list data were used to estimate the size of the population and adjusted for GP participation. Numerator data (incidence cases of WRIH) from 2006–2015 were adjusted for response rates and time sampling frequency. Numerator data were then divided by the denominator and rates calculated per 1 00 000 persons employed per annum

Results The incidence rate of WRIH (from 6491 numerator cases) was 1350 per 1 00 000 persons employed per annum. Musculoskeletal disorders had the highest rates (640) followed by mental ill-health (516). Rates for musculoskeletal disorders were highest in agriculture (2232) and construction (1048) and for mental ill-health, in financial services (1113) and public administration/defense (1124).

Conclusion All employed persons have access to a GP therefore calculating incidence rates from GP data allow identification of industries/occupations at higher risk of WRIH Using postcode-based census data to characterise the denominator (therefore specifically related to the numerator) enables rates to be calculated with greater accuracy compared with using national employment data.