Oral Presentation

Dusts and Fibres

0402 DOES LUNG CANCER INCIDENCE AND MORTALITY DIFFER WITH THE TYPE OF ASBESTOS FIBRE? : EVIDENCE FROM WESTERN AUSTRALIA

¹Nita Sodhi, ^{1,2}Nick deKlerk*, ¹Peter Franklin, ^{1,3}Fraser Brims, ⁴Susan Peters, ¹Nola Olsen, ^{1,3}Bill Musk. ¹The University of Western Australia, Perth, WA, Australia; ²Telethon Kids Institute, Perth, WA, Australia; ³Sir Charles Gairdner Hospital, Perth, WA, Australia; ⁴Utrecht University, Utrecht, The Netherlands

10.1136/oemed-2017-104636.330

Background Crocidolite is known to be the most carcinogenic form of asbestos for mesothelioma. This study aimed to assess if there was a difference between crocidolite and mixed asbestos fibre exposure for rates of lung cancer incidence and mortality after adjusting for smoking.

Methods Administrative datasets were used to identify lung cancer incidence and deaths in three Western Australian cohorts of people exposed to: crocidolite as miners and millers at Wittenoom; crocidolite as residents of Wittenoom; and mixed asbestos fibres occupationally. Multivariable Poisson regression models were used to compare the 5 year incidence and mortality rates between these cohorts after starting a health surveillance program. Explanatory variables included radiological evidence of asbestosis and cumulative asbestos exposures as determined through our Asbestos Job-Exposure Matrix (AsbJEM) after adjusting for smoking pack years, time since smoking cessation, age and sex.

Results The 5 year lung cancer incidence (n=176) and mortality (n=125) rates increased with radiological evidence of asbestosis (2.4-fold), increasing cumulative asbestos exposure and smoking pack-years. Compared with crocidolite miners and millers, Wittenoom ex-residents had the lowest rates (Incidence Rate Ratio (RR): 0.31, 95% CI: 0.15–0.67; and Mortality RR: 0.39 95% CI: 0.17–0.88). However, there were no significant differences identified between people occupationally exposed to mixed asbestos fibres (Incidence Rate Ratio (RR): 0.92, 95% CI: 0.61–1.39; and Mortality RR: 1.07 95% CI: 0.66–1.73) and Wittenoom workers.

Conclusion Mixed asbestos fibre exposure seems to be an equally potent cause of lung cancer as crocidolite after adjusting for smoking history.

Oral Presentation

Occupational Medicine (SCOM/Modernet)

0403 FINDINGS FROM THE FIRST YEAR OF MAREL: THE ITALIAN NETWORK ON WORK-RELATED DISEASES

¹Stefania Curti*, ¹Stefano Mattioli, ²Pierluigi Cocco, ³Alfonso Cristaudo, ⁴Marco dell'Omo, ⁵Giovanni Mosconi, ⁶Giuseppe Campo. ¹Department of Medical and Surgical Sciences, University of Bologna, Bologna, Italy; ²Department of Public Health, Clinical and Molecular Medicine, University of Cagliari, Cagliari, Italy; ³Unit of Occupational Medicine, University of Pisa, Pisa, Italy; ⁴Unit of Occupational and Environmental Medicine, Department of Medicine, University of Perugia, Perugia, Italy; ⁵Unit of Occupational Medicine, Azienda Ospedaliera Papa Giovanni XXIII, Bergamo, Italy; ⁶Department of Work and Environmental Medicine, Epidemiology and Hygiene, National Institute for Insurance against Accidents at Work (INAIL), Rome, Italy

10.1136/oemed-2017-104636.331

Background Detecting new disease-exposure associations is an important public health issue. We created a pilot network (MAREL - MALattie e Rischi Emergenti sul Lavoro) of occupational disease consultation centres of Italian university hospitals to which patients are referred for potentially work-related diseases.

Methods The MAREL network currently includes five occupational disease consultation centres of university hospitals located in central-northern Italy. Patients are referred to these consultation centres by their general practitioners, occupational physicians or other specialists for the investigation of the putative occupational origin of the disease. Each centre collects cases of putative occupational origin through a structured and standardised data collection form. We collect data on: diagnosis; personal habits; occupational history; exposure assessment for potentially associated risk factors; physician's opinion on the possible causal relationship between disease and occupation. Data are coded according to national and international classifications.

Results The data collection started in 2016. We collected 1516 cases of putative occupational diseases. Intervertebral lumbar disc degeneration was the most represented condition (n=170, 11.2%). Apart from musculoskeletal disorders, cases of asthma were the most frequently reported (n=59, 3.9%). About 11% of patients were construction workers. The most frequently reported exposures were: manual material handling (29.5%, out of 1811), hand-arm vibrations (13.9%), and repetitive movements of the upper limb (10.9%).

Discussion The MAREL network will be expanded to other occupational disease consultation centres in 2017 with the aim to contribute to already existing surveillance systems (i.e. MALPROF) by the detection of new and emerging occupational diseases and risks.