Poster Presentation

Disease Surveillance

0404 SURVEILLANCE OF OCCUPATIONAL EYE INJURY THROUGH EMERGENCY SERVICES: TWO-YEAR EXPERIENCES IN A MEDICAL CENTRE OF TAIWAN

¹Ting-Chia Weng*, ^{2,3}Lin Hsu, ²Chia-Chang Chuang, ^{1,3}Jung-Der Wang, ^{1,4}How-Ran Guo, ^{1,4}Yau-Chang Kuo. ¹Department of Occupational and Environmental Medicine, National Cheng Kung University Hospital, Tainan, Taiwan; ²Department of Emergency Medicine, National Cheng Kung University Hospital, Tainan, Taiwan; ³Department of Public Health, College of Medicine, National Cheng Kung University, Tainan, Taiwan; ⁴Department of Environmental and Occupational Health, National Cheng Kung University, Tainan, Taiwan

10.1136/oemed-2017-104636.332

Ocular trauma in the workplace is a worldwide cause of visual morbidity, but is largely preventable. We established a pilot surveillance system of occupational eye injuries at emergency service of National Cheng Kung University Hospital (NCKUH) from Feb 2015 to Feb 2017. During the study period, 114 events of occupational eye injuries were collected with 22 patients further hospitalised. Among them, 3 cases of severe ocular trauma were accompanied with other injuries resulting in an Injury Severity Score of more than 16. Foreign body injury (28%, n=32) and chemical burn (21%, n=24) were the most frequent eye injuries. Notably, 12 cases (10.5%) eyeball rupture were identified. The total medical costs charged through the NCKUH were about 66,000 USD for these ocular trauma patients, counting emergency and hospitalisation fees within 90 days after the first encounter. Eyeball ruptures accounted for 57.5% of the total medical charges. Only about 18% of occupational ocular trauma cases were reported to the National Labour Insurance and would usually be compensated. This study highlights the urgent needs of comprehensive compulsory regulations of recognition and report, compensation, and prevention and control of occupational eye injuries in the workplace of Taiwan.

Oral Presentation

Occupational Medicine (SCOM/Modernet)

0405 PREDICTING THE IMPACT OF THE EU VIBRATION DIRECTIVE ON THE PREVALENCE OF VIBRATION WHITE FINGER (VWF), CARPAL TUNNEL SYNDROME (CTS) AND SENSORINEURAL SYMPTOMS ACROSS EUROPE

¹S Jill Stocks*, ²Begoña Martínez-Jarreta, ³Riitta Sauni, ¹David A Berk. ¹University of Manchester, Manchester, UK; ²University of Zaragoza, Zaragoza, Spain; ³Ministry of Social Affairs and Health, Tampere, Finland

10.1136/oemed-2017-104636.333

Objectives EU directives can lend themselves to evaluation through natural experiments because of the freedom regarding their implementation within individual countries. Natural experiments may be criticised because of the lack of direct casual evidence for a relationship between the intervention and the exposure or disease outcome. Such experimental designs could be strengthened by an a priori estimate of the impact of an intervention. Here we aim to predict the impact

of EU Directive 2002/44/EC on VWF, CTS and sensorineural symptoms (HAVS).

Methods Previously we described the development of a Markov chain Monte Carlo accelerated failure time model to predict the incidence and prevalence of HAVS. The model was developed using longitudinal data from Italian workers and validated using published data for Swedish and UK workers, and compensation data from the Czech Republic. For the next step we have used the exposure data from the Eurofound European Working Conditions Survey from 2000 to 2015 and population level demographic data to predict the impact of the directive on HAVS in Europe.

Results The model predicted that a reduction in vibration exposure to 5 m/s² (i.e. complete success of the directive) would result in a reduction in lifetime prevalence of VWF of around 25% in the UK. Predictions of the variation in impact according to differing reductions in exposure and across different age groups will be presented for the UK and other countries.

Conclusion Future work will compare these estimates of the impact of the directive using routinely collected data in European countries.

Poster Presentation

Cancer

0406 ANALYSIS OF INFECTIOUS DISEASE PREVALENCE AMONG SEMICONDUCTOR MANUFACTURING WORKERS WITH NON-HODGKIN'S LYMPHOMA

Jung-min Sung*, Sang-gil Lee, Eun-A Kim. KOSHA, Ulsan, Republic of Korea

10.1136/oemed-2017-104636.334

Introduction The Occupational Safety and Health Research Institute (OSHRI) established a cohort of former and current workers in six semiconductor-manufacturing companies in South Korea, to determine the incidence of occupational cancer. This study compared the prevalence and incidence of general illnesses between patients with non-Hodgkin lymphoma (NHL) and a control group.

Research Methods After identifying 43 workers with NHL, OSHRI established a sex- and age-matched control group. The NHL cohort and control groups were compared for the prevalence of infectious diseases before and after diagnosis. The prevalence of infectious diseases was based on diagnostic data collected from National Health Insurance information. Disease names were determined by the ICD-10 disease code.

Outcomes Prior to the diagnosis of NHL, there was no significant difference in the prevalence of bacterial and viral diseases between the groups. However, following the diagnosis of NHL, the prevalence rates of viral and bacterial infections were significantly higher in the NHL group. Significant differences were found in the prevalence of herpes zoster, cytomegalovirus, and tuberculosis.

Conclusion Prior to the diagnosis of NHL, there were no significant underlying diseases in the NHL cohort group when compared to the control group. Therefore, we concluded there is no relationship between underlying infectious disease and NHL development. However, the probability of contracting opportunistic infections was higher in the NHL group

after diagnosis. Presumably, cancer progression and treatment may cause opportunistic infections. Further evaluation is necessary to support this result.

Poster Presentation

Exposure Assessment

0407

RESPIRATORY IRRITANT EXPOSURES DURING CLEANING AND DISINFECTING IN HOME CARE: PRELIMINARY RESULTS FROM THE SAFE HOME CARE STUDY

Margaret Quinn*, Catherine Galligan, John Lindberg, Pia Markkanen, Susan Sama, Rebecca Gore, Noor Sheikh, Nicole Karlsson, David Kriebel, Nancy Goodyear. *University of Massachusetts, Lowell, MA, USA*

10.1136/oemed-2017-104636.335

Objectives While there is evidence of asthma and other respiratory illness among workers performing cleaning and disinfecting (C and D) of environmental surfaces in healthcare and in residential and institutional settings, quantitative measurements of C and D exposures are limited. Previously, we found that 80% of home care (HC) aide visits to clients' homes involve C and D; commonly-used C and D products contain bleach, a respiratory irritant. Objectives of this study were to measure quantitatively airborne exposures generated during C and D tasks performed by HC aides using a bleach-containing product. The work reported here is part of a larger evaluation of exposures and respiratory effects of a range of C and D products and practices and their efficacy in reducing pathogens in HC.

Methods A bathroom was constructed in a laboratory according to home building construction specifications. Twenty HC aides were recruited from employer agencies to perform C and D tasks for 20 min sessions in the bathroom following typical HC practices using a C and D spray product containing bleach (1%–5% by weight sodium hypochlorite). Aides wore a vest holding a direct-reading instrument to measure chlorine breathing-zone concentrations as they performed C and D on tub/shower, toilet, and sink.

Results Maximum chlorine concentrations generated during the 20 min sessions ranged from 0.35ppm to 3.40ppm. The tub/shower C and D task produced the highest exposures. The US Occupational Safety and Health Administration ceiling limit for chlorine is 1 ppm; nearly 70% of the aides conducted a C and D session exceeding this value.

Conclusions C and D in HC using a product containing bleach can produce over-exposure to chlorine, a respiratory irritant.

Poster Presentation

Chemicals

0408

WORKERS' EXPOSURE TO BROMINATED FLAME RETARDANTS: A GLANCE AT AMERICAN AND CANADIAN POPULATION DATABASES

^{1,2}Sabrina Gravel*, ^{1,3}Jérôme Lavoué, ^{1,2}France Labrèche. ¹*School of Public Health, Université de Montréal, Montreal, QC, Canada;* ²*Institut de recherche Robert-Sauvé en santé et en sécurité du travail, Montreal, QC, Canada;* ³*Centre de recherche du Centre hospitalier de l'Université de Montréal, Montreal, QC, Canada*

10.1136/oemed-2017-104636.336

We aim to determine biological concentrations of four congeners of brominated flame retardants, which have no occupational exposure limits, in American and Canadian workers.

Serum concentrations of four polybrominated diphenyl ethers congeners (BDE47, 99, 100 and 153) were obtained from the American National Health And Nutrition Examination Survey (NHANES) 2003–2004 and the Canadian Health Measures Survey (CHMS) 2007–2009 databases. Data from participants aged 16–65 were classified by industry and occupational group. Values below the detection limit (< LOD) were replaced by LOD/ $\sqrt{2}$. Descriptive statistics are presented.

A total of 813 and 1100 serum samples were respectively available in NHANES and CHMS. Proportions of values < LOD varied by survey: BDE47 (NHANES, 1.4%; CHMS, 22.7%), BDE99 (32.3%; 73.9%), BDE100 (4.7%; 72.9%), BDE153 (5.7%; 55.9%). BDE47 was the congener with the highest lipid-adjusted mean concentrations (NHANES, 44.5 ng/g; CHMS, 22.8 ng/g). Higher BDE47 means were found in Agriculture, Forestry and Fishing industries (61.9 ng/g) and in Protective Services and Armed Forces occupations (48.5 ng/g) for NHANES, whereas in CHMS they were higher in Manufacturing of Durable Goods industries (40.3 ng/g) and in Construction and Extraction occupations (44.7 ng/g). BDE153 means were higher for men than for women in both databases (NHANES-men: 16.8 ng/g, NHANES-women: 11.4 ng/g; CHMS-men: 11.0 ng/g, CHMS-women: 7.8 ng/g). In NHANES, non-working men had higher mean concentrations than workers for all congeners.

In the absence of occupational exposure limits, population surveys can be useful to establish reference levels, but careful interpretation is required for chemicals such as flame retardants that are ubiquitous in both the general and work environments.