Poster Presentation

Psychosocial

0430 FEMALE WORKERS SUFFERED HIGHER PSYCHOLOGICAL SYMPTOMS AFTER OCCUPATIONAL INJURIES THAN MALE WORKERS

¹Judith Shiao*, ^{2,3}Wei-Shan Chin, ³Sherri Yeh, ^{2,3}Leon Guo. ¹Nursing, National Taiwan University (NTU) College of Medicine and NTU Hospital, Taipei, Taiwan; ²National Institute of Environmental Health Sciences, National Health Research Institutes, Zhunan, Taiwan; ³Environmental and Occupational Medicine, National Taiwan University (NTU) and NTU Hospital, Taipei, Taiwan

10.1136/oemed-2017-104636.355

Introduction This study aimed to examine the prevalence rates of psychological symptoms and risk factors between female and male workers at 12 months after their sustaining occupational injury. Demographic and injury-related risk factors for psychological symptoms were evaluated.

Methods Our study candidates were injured workers in Taiwan who were hospitalised for 3 days or longer and received hospitalisation benefits from the Labour Insurance program. A self-reported questionnaire including the Brief Symptom Rating Scale was sent to workers at 12 months after injury.

Results A total of 1233 workers (response rate 28.0%) completed the questionnaire, including 356 women and 877 men. A higher percentage (30.1%) of women had elevated BSRS-5 score of 6 or higher than men (22.5%). The risk factors for elevated psychological symptom scores for female workers were lower education level (odds ratio, OR=1.8, 95% confidence interval, CI=1.1–3.0), main income contributor of the household (OR=1.8, CI=1.1–3.0), severely affected physical appearance due to injury (OR=2.8, CI=1.3–5.9), and having adverse life event after injury (OR=2.0, CI=1.1–3.6) after mutual adjustment. Whereas the risk factors for elevated psychological symptom scores for female workers were loss of consciousness after the injury (OR=2.0, CI=1.3–3.1), severely affected physical appearance due to injury (OR=3.7, CI=2.3–6.0), having adverse life event after injury (OR=2.5, CI=1.6–3.8), not return-to-work (OR=3.2, CI=2.0–5.1), and reduced salary as compared to that before injury (OR=2.4, CI=1.3–4.1).

Conclusions After occupational injury, women suffered from higher rate of psychological symptoms. Risk factors were different between men and women who sustained occupational injuries.

Poster Presentation

Methodology

0431 HOW CAN WE AVOID RE-IDENTIFICATION RISK IN BIG-DATA ANALYSIS? PROPOSITION OF A NEW STRATEGY OF GEOGRAPHICAL SUBDIVISIONS USING GIS TOOLS

^{1,2}Charlotte Maugard*, ³Christophe Cancé, ¹Pauline Achard, ²Olivier François, ^{1,4}Vincent Bonnetterre, ^{1,2}Delphine Bosson-Rieutort. ¹Grenoble-Alpes University (UGA)/TIMC-IMAG Laboratory (UMR CNRS 5525)/EPSP Team (Environment and Health Prediction of Populations), Grenoble, France; ²Grenoble-Alpes University (UGA)/TIMC-IMAG Laboratory (UMR CNRS 5525)/BCM Team (Computational and Mathematical Biology), Grenoble, France; ³Grenoble-Alpes University (UGA)/UMS GRICAD, Grenoble, France; ⁴Grenoble-Alpes Teaching Hospital ("CHU Grenoble-Alpes")/Occupational Medicine and Health Department, Grenoble, France

10.1136/oemed-2017-104636.356

In order to look for relevant signals for detection of emerging occupational diseases among agricultural workers, we developed a data-mining approach applied on health-insurance data (see C. Maugard communication). Applied on the databases of the French dedicated social security system (MSA), this approach first aims to look for associations between chronic diseases and occupational activities (recorded as activity sector codes in the MSA contributors database).

To avoid re-identification, workers location has not been provided, although it is recognised as closely related to cultural practices. Therefore, it was not possible to directly estimate individuals involvement in specific cultures (through existing parcel register and agricultural census for instance) and finally use cultures x pesticides to estimate pesticides exposures.

To deal with this issue, we used an innovative approach to cut off the national territory into "meshes", to obtain a geographical variable accurate enough to assess cultures types while respecting a sufficient number of agricultural workers per meshes to avoid re-identification. This approach consists of an iterative process dividing each geographical unit into 4 parts while respecting a minimum threshold of workers in each mesh. The process continues until each mesh contains a homogeneous number of individuals. Taking into account the prevalence of the chronic diseases of interest, and typology of cultures, we defined a minimum number of individuals per mesh (n=1500). This methodological development allows us to get indirect information about location by MSA at a level interesting to identify cultures (proxy for pesticides use), but restricting the possibilities of individuals re-identification.

Poster Presentation

Cancer

0432 PARENTAL EXPOSURE TO PAINTS AND RISK OF CHILDHOOD CANCER

³Julie Volk*, ³Johnni Hansen, ²Julia Heck, ¹Kjeld Schmiegelow. ¹Department of Paediatrics and Adolescent Medicine, University Hospital Rigshospitalet, Copenhagen, Denmark; ²Department of Epidemiology, UCLA School of Public Health, California, USA; ³Department of Diet, Genes and Environment, Danish Cancer Society Research Centre, Copenhagen, Denmark

10.1136/oemed-2017-104636.357

Objective Even though childhood cancer is a rare disease, it is one of the main causes of death among children in the Western world. Not much is known about the causes of childhood cancers but parental occupational exposures have been suggested by a number of epidemiological studies, including exposure to paints.

Methods and materials All childhood cancer cases (0–15 years) in Denmark from 1968–2012 (n=5,711) were retrieved from the Danish Cancer Registry and population controls (1:100) were randomly selected and matched by age and sex. Maternal and paternal occupational history was retrieved by the Supplementary Pension Fund. Potential confounders were retrieved through the Medical Birth Registry. Register linkages were conducted using the unique identification number assigned to all Danish residents.

Results Preliminary results for cancer of all sites show an OR of 0.89 (95% CI: 0.71–1.01) and 0.86 (95% CI: 0.73–1.01) for maternal and paternal exposure to paint, respectively, after controlling for potential confounders, including SES, maternal smoking, birth order, previous miscarriage, malformation and parental age. Increased but insignificant ORs were found for acute lymphatic leukaemia, non-Hodgkin lymphoma, ependymoma, astrocytoma, Burkitt lymphoma, central nervous system cancers, Ewing sarcoma, melanoma and hepatoblastoma for maternal exposure and acute myeloid leukaemia, glioma, melanoma, neuroblastoma and hepatoblastoma for paternal exposure.

Conclusion Preliminary results have shown little and insignificant effect of parental paint exposure in relation to childhood cancer risk.

Poster Presentation

Disease Surveillance

0433 TOWARDS AN IDEAL NATIONAL WORK-RELATED ILL HEALTH SURVEILLANCE SYSTEM IN GREAT BRITAIN

Yiqun Chen*, David Fishwick, Andrew Curran. *Health and Safety Executive, Buxton, UK*

10.1136/oemed-2017-104636.358

Population-based occupational health surveillance includes work-related ill health surveillance and workplace health risk exposure surveillance. It is an important intelligence gathering system at the national level, which supports the planning, monitoring and evaluation of measures to prevent work-related ill health. The

intelligence is essential for targeted intervention, prioritisation, tracking progress and evaluation of impact.

To make best use of resources for intelligence gathering, the approaches used should be regularly assessed and monitored to ensure they remain fit-for-purpose, cost-effective and forward looking. In the context of the strategic approach for research planning and prioritisation in the Health and Safety Executive (HSE), a series of workshops were developed.

One of the HSE internal workshops was organised in January 2017. Some, 26 (90%) of the 29 invited stakeholders have participated to develop a common vision for a population-based work-related ill health surveillance system that will continue to meet HSE's intelligence needs now and in the future;

Following detailed assessments of the gaps in the current system, a wide range of innovative approaches were explored. Some practical first steps to improve the system were recommended with an emphasis on more systematic and strategic data collection. The key characteristics of an ideal system were also identified, including new features on case investigation and detecting new/emerging work-related ill health risks to inform timely preventative actions. The outputs of the workshop are presented. They have informed HSE priorities in the continued development of the system to support its mission to prevent work-related ill health.

Oral Presentation

Dusts and Fibres

0434 A META-ANALYSIS OF OCCUPATIONAL SILICA EXPOSURE AND RISK OF AUTOIMMUNE RHEUMATIC DISEASES: DOES STUDY QUALITY MATTER?

¹Henrik A Kolstad*, ²Christiane Beer, ³David Sherson, ⁴Anne Trolldborg, ⁴Berit Dalsgaard Nielsen, ⁵Anne Braae Olesen, ⁶Gitte Jacobsen, ⁴Søndergaard Klaus, ^{2,7}Schlünssen Vivi. ¹Department of Occupational Medicine, Danish Ramazzini Centre, Aarhus University Hospital, Aarhus, Denmark; ²Department of Public Health, Danish Ramazzini Centre, Aarhus, Denmark; ³Department of Occupational and Environmental Medicine, University of Southern Denmark, Odense, Denmark; ⁴Department of Rheumatology, Aarhus University Hospital, Aarhus, Denmark; ⁵Department of Dermatology, Aarhus University Hospital, Aarhus, Denmark; ⁶Department of Occupational Medicine, Danish Ramazzini Centre, Hospital Region Vest, Herning, Denmark; ⁷National Research Centre for the Working Environment, Copenhagen, Denmark

10.1136/oemed-2017-104636.359

Objectives Increased risks of rheumatoid arthritis, small vessel vasculitis, systemic lupus erythematosus, and systemic sclerosis have been observed following crystalline silica exposure. Our aims are to estimate pooled risk estimates and assess the impact of study quality.

Methods We followed the PRISMA criteria, identified 1162 articles, and included 21 studies that we classified according to eight quality parameters (high vs. low). We estimated pooled overall and disease specific odds ratios (ORs) with random effects meta-regressions.

Results We observed an increased overall OR of 2.3 (1.7–3.1, 21 studies) and for rheumatoid arthritis (OR 1.7, 95% CI 0.8–3.41, 6 studies), small vessel vasculitis (OR 2.4, 95% CI 1.2–4.7, 6 studies), systemic lupus erythematosus (OR 2.8, 95% CI 0.5–14.7, 3 studies), and systemic sclerosis (OR) 2.9, 1.7–4.9, 6 studies). The following high-quality characteristics