respirable crystalline silica (RCS) and the risk of lung cancer in pooled analyses of community-based studies.

Methods RCS was estimated using a quantitative general population job-exposure matrix ('SYNJEM') for 16 786 lung cancer cases and 20 818 matched population or hospital controls. 34.2% of the men were ever exposed to RCS, while 8.6% of the women were ever exposed. Odds ratios (ORs) and 95% confidence intervals (95% CI) were estimated stratified by gender using unconditional logistic regression models adjusted for age, study centre, cigarette pack-years, time-since-quitting smoking, and ever employment in an occupation with known lung cancer risk.

Results We observed a monotonic increase in risk of lung cancer associated with occupational exposure to RCS among men (unexposed versus 4th quartile among exposed 1.45 (95%CI, 1.31–1.60). Result did not differ by smoking status and remained significantly elevated among non-smokers. The association was stronger in squamous cell carcinoma and small cell lung cancer as compared to adenocarcinoma of the lung especially among former and current smokers. The effect of RCS on lung cancer in women was not detectable likely related to small numbers.

Conclusions The SYNERGY results show that occupational silica exposure is associated with an increased risks of all lung cancer types in a pooled analyses of community-based studies.

Poster Presentation Cancer

0476 THE BRAZIL-ITALY PROJECT ON ASBESTOS-RELATED DISEASES IN CURITIBA

¹Dario Consonni*, ²Luciana Puchalski Kalinke, ³Marco Aurélio Kalinke, ²Fernanda Moura D'Almeida Miranda, ²Larissa Marcondes, ²Tatiana Hallfeld, ²Christiane Brey, ²Shirley Boller, ^{4,1}Angela Cecilia Pesatori, ¹Carolina Mensi, ²Leila Maria Mansano Sarquis. ¹Department of Preventive Medicine, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; ²Universidade Federal do Paraná, Curitiba, Brazil; ³Universidade Tecnológica Federal do Paraná, Curitiba, Brazil; ⁴Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy

10.1136/oemed-2017-104636.394

Brazil is one of the major producers of chrysotile worldwide, but few studies evaluated its effects on the health of Brazilians. Moreover, there are concerns of miscoding of mesothelioma cases. A joint project was started between the Federal University of Paraná (UFPR), Curitiba, Brazil, and the Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy to evaluate effect of chrysotile exposure. The project, supported by CAPES, Brasilia, include two main areas of research. The first is a cohort mortality study among asbestos-cement workers. From the Ministerio Público do Trabalho, Curitiba, we recently received lists of about 10 000 workers employed since the beginning of production (1974, 1974, and 1993) in three asbestos-cement factories in Curitiba and its Metropolitan Region. Data editing and evaluation of sources of information to perform the mortality follow-up are in progress. The second area involves the implementation of a registry of malignant mesothelioma cases in Curitiba based on the Italian model. For this reason, two post-docs are in Milan to study procedures of the Italian and Lombardy Region Mesothelioma Registry, to translate the Italian standardised questionnaire on asbestos exposure, and to

adapt the database. As a complementary study, we are analysing the records (1998–2012) related to mesothelioma of the Curitiba Population-Based Cancer Registry. Clinical documentation of 269 adult subjects with ICD-10 codes C45 (mesothelioma), C38 (mediastinal and pleural cancers), and C48 (peritoneal cancers) will be examined to identify potential false negatives (i.e., primary pleural/peritoneal cancers classified under incorrect cancer sites like mediastinum and retroperitoneum).

Poster Presentation Exposure Assessment

0478 INHALATION AND DERMAL EXPOSURE TO TOLUENE AMONG PRINTING WORKERS IN A PLASTIC BAG FACTORY

¹Wantanee Phanprasit*, ²Kanchanok Songpek, ¹Vorakamol Boonyayothin, ¹Dusit Sujirarat. ¹Mahidol University, Bangkok, Thailand; ²Reckitt Benckiser Healthcare Manufacturing (Thailand) Ltd, Bangkok, Thailand

10.1136/oemed-2017-104636.395

Introduction The study was conducted in a plastic bag factory to explore the inhalation and dermal exposure to toluene of the printing workers who wore no PPE and the potential of a charcoal cloth pad (CCP) as a dermal sampler to assess the skin permeation of liquid toluene.

Methods Twenty-seven stationary air samples were collected on the same 9 days with the urine and dermal samples from 11 printing workers. The CCPs were wrapped on each of the workers' fingers. Air samples were collected and analysed according to NIOSH # 1501 and 65 post-shift urine samples were collected and analysed for toluene using GC-HS/FID. Multiple linear regression was employed to analyse the association of the variables.

Result T three urine samples contained toluene exceeding the BEI of 30 mg/L. Toluene on the CCP (TolCCP) is a meaningful predictor for the UTol (p-value=0.027), with r and r² of 0.441 and 0.195 respectively. The absorbed dose of toluene determined from TolCCP ranging in between 1.05–91.94 mg and counting for the maximum of 12.3% TLV.

Discussion The mean of TWA was above the TLV while that of the UTol was well below the BEI. This indicated that the TWA concentrations could be overestimated due to the size of the room and good general ventilation. The dermal exposure was not significant if the workers wear respirators, but if not the dermal absorption could contribute to the overall uptake and may cause the exposure above the TLV.

Oral Presentation Dusts and Fibres

0480 MEASUREMENT OF ASBESTOS FIBRE RELEASE DURING REMOVAL WORKS IN A VARIETY OF DIY ASBESTOS REMOVAL SCENARIOS

¹Geza Benke*, ²Jeremy Trotman. ¹Monash University, VIC, Australia; ²JTA Health, Safety and Noise Specialists, VIC, Australia

10.1136/oemed-2017-104636.396

Background/Aim In Australia the source of non-occupational mesothelioma cases is poorly described. Although some cases are due to living with former asbestos industry workers or living near former asbestos industries, many cases are not exposed to these risk factors. The incidence of cases due to do-it-yourself (DIY) home renovation or maintenance is unknown, primarily due to lack of exposure data. The aim of this simulation study was to measure asbestos fibre release during removal in a variety of DIY asbestos removal scenarios. **Methodology** Nine different exposure scenarios commonly undertaken in DIY home renovation were simulated. Asbestos fibre exposure was monitored for each location with static samplers located in the work area together with personal samples, using high flow rate pumps operating at 6 to 7 Litres per minute, with the analysis of filters undertaken by SEM and PCM. Simulations were designed to be representative of worst case exposure scenarios.

Results Personal sampling resulted in higher fibre release levels compared to static sampling in all nine scenarios. All static sample scenarios were below 0.15 f/ml. However, for personal sampling removal of asbestos cement (AC) sections (as would be required to accommodate a domestic air-conditioning unit) using an angle grinder, resulted in exposure of 13.23 f/ml. Dry cutting of holes for installation of flues in AC roofing was 2.79 f/ml.

Conclusions Exposure levels in DIY removal were found to be low for most scenarios. Use of power tools without wetting in a confined area was found to be the most exposed scenario.

Oral Presentation

Cancer

0482 EXPOSURE TO HEXAVALENT CHROMIUM AND NICKEL AND LUNG CANCER RISK: A POOLED ANALYSIS OF CASE-CONTROL STUDIES FROM EUROPE AND CANADA

¹Thomas Behrens*, ¹Beate Pesch, ²Roel Vermeulen, ³Ann Olsson, ³Joachim Schüz, ²Lützen Portengen, ¹Benjamin Kendzia, ²Hans Kromhout, ³Kurt Straif, ¹Thomas Brüning, ¹on behalf of The SYNERGY Study Group. ¹Institute for Prevention and Occupational Medicine of the German Social Accident Insurance (IPA), Institute of the Ruhr-Universität, Bochum, Germany; ²Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands; ³International Agency for Research on Cancer (IARC), Lyon, France

10.1136/oemed-2017-104636.397

Background There is limited evidence regarding the exposure-effect relationship of exposure to hexavalent chromium (CrVI) and nickel (Ni) with lung cancer. We estimated the cumulative exposure for CrVI and Ni and assessed exposure-effect relationships for lung cancer risk by sex, smoking status, and histological subtypes.

Methods Fourteen case-control studies (1985–2010) from Europe and Canada were pooled, including 16 901 lung cancer cases (80% men) and 20 965 controls (78% men). Cumulative exposure to CrVI and Ni were estimated. Unconditional logistic regression models were fitted to estimate odds ratios (OR), 95% confidence intervals (CI), and exposure-effect trends adjusted for smoking and occupations with recognised lung cancer risk.

Results The OR for the highest quartile ($>98.95 \mu\text{g}/\text{m}^3\text{-years}$) of cumulative CrVI-exposure was 1.33 (95% CI 1.20–1.48) in men and 1.04 (95% CI 0.48–2.23) in women. In never

smokers, the OR for ever CrVI-exposure was 1.37 (95% CI 1.09–1.73) in men, and OR=1.09; 95% CI 0.70–1.69 in women.

The OR for the highest quartile of cumulative Ni-exposure ($>77.53 \mu\text{g}/\text{m}^3\text{-years}$) was 1.30 (95% CI 1.16–1.45) in men and 1.29 (95% CI 0.59–2.81) in women. The OR for ever Ni-exposure was 1.22 in never smokers for both sexes.

Conclusions Our results showed an exposure-dependent excess risk of lung cancer by occupational exposure to Ni in both sexes, and for CrVI in men. The pattern for CrVI in women was less clear. Analysis of an interaction between Cr- and Ni-exposure was impaired by a high correlation of these agents in metal fumes.

Invited

Methodology

0483 INVITED KEYNOTE: SIFTING THE WHEAT FROM THE CHAFF IN OCCUPATIONAL EPIDEMIOLOGY – NOVEL APPROACHES TO CAUSAL UNDERSTANDING

Debbie Lawlor. University of Bristol, Bristol, UK

10.1136/oemed-2017-104636.398

A key aim of epidemiology is aetiology – determining causal effects of variation in population health and wellbeing. However, the ability to distinguish causal effects from non-causal association in all branches of epidemiology, including occupational epidemiology, is difficult. In this talk I will outline some novel approaches to testing causality, including using genetic instrumental variables (Mendelian randomization) and negative control studies, and how these can be integrated with more conventional approaches (multivariable regression analyses in observational data, randomised controlled trials and natural experiments) in a triangulation framework. I will use examples from perinatal and cardio-metabolic health (the main area in which I work) but also from occupational health to illustrate how these can be used to improve causal understanding of how occupational exposures affect health.

Oral Presentation

Cancer

0484 EXPOSURE TO IONISING RADIATION AND RISK OF LYMPHOMA SUBTYPES: ANALYSIS OF THE EPILYMPH RESULTS

¹Pierluigi Cocco*, ²Matteo Ioi, ¹Qi Zhi, ¹G Satta, ¹Maris G Enna, ³Nickolaus Becker, ⁴Yolanda Benavente, ⁴Sylvia De Sanjosé, ⁵Lenka Foretova, ⁶Anthony Staines, ⁷Marc Maynadié, ⁸Alexander Nieters, ²Lydia Zablotska. ¹University of Cagliari, Cagliari, Italy; ²University of California, San Francisco, USA; ³German Cancer Research Centre, Heidelberg, Germany; ⁴Catalan Institute of Oncology, Barcelona, Spain; ⁵Department of Cancer Epidemiology and Genetics, Brno, Czech Republic; ⁶Dublin City University, Dublin, Ireland; ⁷Dijon University Hospital, Dijon, France; ⁸University of Freiburg, Freiburg, Germany

10.1136/oemed-2017-104636.399

Introduction The association between ionising radiation and risk of solid tumours and leukaemia is well established;