Oral Presentation

Other

0150 IS A POSSIBLE RELATIONSHIP BETWEEN OCCUPATIONAL NOISE EXPOSURE AND VESTIBULAR SCHWANNOMA (ACOUSTIC NEUROMA) CAUSAL OR DUE TO BIAS? A SYSTEMATIC REVIEW AND META-ANALYSIS

Ingrid Sivesind Mehlum*, Håkon A Johannessen, Karl-Christian Nordby, Marit Skogstad, Tore Tynes, Arve Lie. *National Institute of Occupational Health, Oslo, Norway*

10.1136/oemed-2017-104636.118

Objectives Vestibular schwannoma (acoustic neuroma) is a benign, slow-growing tumour, arising from the nerve sheath of the vestibular branch of the eighth cranial nerve. Some studies show a possible relationship between noise exposure and schwannoma, while others not. Our objective was to assess possible associations between occupational noise exposure and schwannoma.

Methods We performed a systematic critical literature review of original articles from key literature databases of associations between workplace noise and health, completed in January 2017. For the meta-analysis, we selected studies of sufficient quality with a measure of association between occupational noise exposure and schwannoma.

Results Six studies were selected, all matched case-control studies, with a total of 1487 cases, five population-based and one registry study. Five studies had self-reported exposure information from interview (4) or questionnaire (1), and three had data from job-exposure-matrices (JEMs). Two studies had both self-report and JEM information. Self-reported occupational noise exposure was consistently associated with schwannoma, odds ratio (OR) 1.70 (95% CI 1.32–2.18). JEM results gave OR 1.16 (0.75–1.81), with more heterogeneity between studies. Long noise exposure ($\geq 15/18$ years; 2 self-report, 2 JEM studies) was also consistently associated with schwannoma, OR 1.76 (1.04–2.99).

Conclusions The results indicate that noise exposure might be associated with schwannoma. We found strong and consistent associations for self-reported noise exposures and exposures of longer duration. JEM-based associations were weaker, maybe due to non-differential misclassification. However, reasons other than a causal relationship, such as recall bias, detection bias or publication bias, cannot be excluded.

Poster Presentation

Intervention Studies

0151 SYSTEMATIC REVIEW OF WORKPLACE INTERVENTIONS FOR REDUCING OCCUPATIONAL SITTING

Lotte Röder, Eva Backé, Ute Latza*, b a u a : *Federal Institute for Occupational Safety and Health, Berlin, Germany*

10.1136/oemed-2017-104636.119

Objectives Based on a review of reviews that included studies until December 2015, we demonstrated a reduction in occupational sitting for multi-component workplace interventions, and interventions addressing the work environment and the individual. This systematic review gives an overview about the rapidly accumulating subsequent evidence and identifies research needs.

Methods A literature search was conducted in Pubmed, PsycInfo, and Sportdiscus. English and German references from 01.01.2016 until 31.07.2016 were included if they evaluated workplace interventions targeting a reduction of occupational sitting or an increase of physical activity. Quality was assessed with a modified version of the Cochrane Risk of Bias Tool.

Results Out of 380 identified references 17 were included (9 randomised controlled trials (RCTs), and 8 non-randomised interventions). Six out of 7 interventions (4 RCTs) addressing the individual, 3 out of 6 studies (1 RCT) targeting the physical work environment (sit-to-stand/dynamic workstations, active building design), and 2 out of 2 multi-component interventions (RCTs) showed statistically significant beneficial changes in objectively measured occupational sitting time and/or physical activity.

Conclusions Beyond inconsistent results for dynamic workstations and further evidence for multi-component interventions, individualised interventions (e.g. personal feedback elements) seem to be beneficial. The effect of new components (e.g. active building design) is still unclear. Future studies are warranted emphasising organisational interventions, and the mechanisms behind single components of multi-component interventions (e.g. social support) with their long-term effects. Further, the interrelationship of prolonged sitting on and off work, and timing of sitting interruptions for cardiometabolic effects need to be further elucidated.

Oral Presentation

Cancer

0152 CANCER MORTALITY OF DMF EXPOSED WORKERS IN KOREA

¹Yeon-Soon Ahn*, ¹Jin-Ha Yoon. ¹Dongguk University Ilsan Hospital, Goyang, Republic of Korea; ²Yeonsei College of Medicine, Seoul, Republic of Korea

10.1136/oemed-2017-104636.120

This study is to identify the association of urinary N-methylformamide level (UNMFL) with cancer mortality in N,N-dimethylformamide (DMF)-exposed male workers in Korea. A cohort was composed of 11,953 DMF exposed workers working between January 1, 2000, and December 31, 2004. These cohort members were matched with the mortality data of the Korean National Statistical Office to follow-up for cancer mortality between 2000 and 2011. Standardised Mortality Ratios (SMRs) of DMF exposed workers with reference to Korean men were calculated. Also controlling age, other carcinogen exposure including hepatitis B and C, the Adjusted Hazard Ratios (AHRs) of workers categorised by the 3 groups of UNMFL with reference to workers with zero UNMFL were calculated. There were no significantly increased or decreased SMRs except for stomach cancer (SMR=0.38, 95% CI=0.10–0.98). The AHRs of overall cancer mortality were significantly increased in workers with 7.5 to 15 mg/L (SMR=2.72, 95% CI=1.09–6.81) and 15 and over 15 mg/L (SMR=2.41, 95% CI=1.03–5.66) compared with workers with 0 UNMFL. Hepatocellular carcinoma mortality (AHR=3.73, 95% CI=1.05–13.24) of workers with 15 and over 15 mg/L and lung cancer mortality (SMR=14.36, 95% CI=1.41–146.86) in workers with 7.5 to 15 mg/L were significantly increased compared with workers with 0 UNMFL. In this study workers with high UNMFL showed increased mortalities for overall, liver and lung cancer comparing to those of workers with zero UNMFL, which suggests DMF might be caused cancer, especially hepatocellular carcinoma which was approved carcinogenicity on liver in animal experiments.

Oral Presentation

Cancer

0153 RISK OF BLADDER CANCER IN A COHORT OF CHEMICAL WORKERS

¹Dirk Taeger*, ¹Beate Pesch, ²Bierfreund Kay-Gerald, ³Christoph Oberlinner, ²Gabriele Leng, ⁴Stenzl Arnulf, ⁵Matthias Kluckert, ¹Thomas Brüning. ¹Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Institute of the Ruhr-Universität Bochum (IPA), Bochum, Germany; ²Department of Health Protection and Occupational Safety, Currenta GmbH and Co. OHG, Leverkusen, Germany; ³Occupational Medicine and Health Protection, BASF SE, Ludwigshafen, Germany; ⁴Institute of Urology, Eberhard Karls University, Tübingen, Germany; ⁵German Social Accident Insurance's Institution for the Raw Materials and Chemical Industry (BG RCI), Heidelberg, Germany

10.1136/oemed-2017-104636.121

Objective Bladder cancer is the fourth most frequent cancer among men in Germany. Aromatic amines can cause bladder cancer and therefore cancerogenic aromatic amines have been

banned a long time ago from workplace, but a long latency can still increase the risk of former exposed workers. In this study, we will assess the risk of bladder cancer in a cohort of chemical workers with earlier use of aromatic amines compared to the general population.

Methods In the prospective cohort study UroScreen, 1800 former chemical workers exposed to aromatic amines were offered to participate in an early detection of bladder cancer by means of tumour markers. In 2003–10, 1609 people were examined at least once. The exposure to aromatic amines was determined by means of questionnaires. The observed bladder cancer incidence was compared with the expected incidence in the general population. Bladder cancer risk was estimated as standardised incidence ratio.

Results Nine incidence urinary bladder carcinoma occurred during the study. Eight cases were ex-smokers and one case was non-smoker. All cases were exposed for at least 10 years, including six cases longer than 20 years. Compared to the general population, the risk of bladder cancer was 2.94 (95% CI 1.35–5.59).

Conclusions Since almost all cases have both smoked and were highly exposed, reliable risk detection is difficult. Nevertheless, in view of a threefold increased risk, the early detection of urinary bladder carcinomas were promising.

Poster Presentation

Other

0154 OCCUPATIONAL DISEASES ATTENDED AT PARC DE SALUT MAR (BARCELONA): CARE COSTS IN A SERIES OF CONFIRMED CASES (2010–2014)

¹M Valinho-Carrete, ^{2,4}FG Benavides, ^{1,3}A Beltran-Fonollosa*, ^{2,4}JM Ramada, ²F Cots, ⁵M Trapero-Bertran, ^{4,6}GL Deldos, ^{1,4}C Serra. ¹CISAL-Centre for Research in Occupational Health, University Pompeu Fabra, Barcelona, Spain; ²IMIM Hospital del Mar Medical Research Institute, Barcelona, Spain, Barcelona, Spain; ³Occupational Health Service, Parc de Salut Mar, Barcelona, Spain; ⁴CIBER de Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain; ⁵Facultad de ciencias económicas y sociales. Universitat Internacional de Catalunya, Barcelona, Spain; ⁶The University of Texas School of Public Health, Houston, Texas, USA

10.1136/oemed-2017-104636.122

Background The Spanish National Health System (NHS) covers non-work-related diseases. The Social Security system, mainly through collaborating insurance companies, covers both medical and wage-related costs of occupational diseases; however, the actual exchange of resources has not been well studied until now. The objective of this study was to evaluate the economic cost of medical care generated by a series of confirmed cases of occupational disease (OD) treated at Parc Salut Mar (PSMAR) in Barcelona, an NHS health system.

Methods Economic study of 40 cases of suspected OD by the PSMAR Occupational Disease Unit (ODU). Between 2010 and 2014, information on the care received (stays, visits, emergencies, diagnostic tests, medical and surgical treatments, etc.) was independently reviewed by three experts, and discrepancies resolved by one of them. The economic value of the care received in relation to the underlying suspected OD was expressed in terms of actual cost and billing rate, according to age, sex, diagnostic group and type of care received.