Union. The original platform (developed for Microsoft Access) allowed for assignment of exposure proportions by 55 industry categories, but was a breakthrough at the time in terms of amalgamating exposure measurement data and occupational hygiene knowledge surrounding carcinogen exposure. Recognising the importance of CAREX in occupational cancer prevention, several other countries around the world have since adapted the original system for use in their own countries, with a few making large improvements to the model. A notable example is Costa Rica, with their TICAREX adaptation that estimated pesticide exposure for the first time, and considered sex as an exposure-defining feature in workplaces. In Canada, the system was expanded further to consider exposure by hundreds of detailed industry and occupation codes, sex, sub-geographical regions within a country, and level of exposure where possible. In addition, an entirely new system for considering community environmental carcinogen exposures was added. The Canadian team has been working with the Pan American Health Organisation and other partners to expand the use of the enhanced CAREX to other countries, in particular those of lower and middle income, where capacity for new research and data structures may be difficult. After 25 years, the CAREX model continues to evolve and improve to meet current needs.

Oral Presentation

Intervention Studies

0285 ASSESSING THE IMPACT OF A GROUP RANDOMISED CONTROLLED INTERVENTION STUDY IN SUPERMARKET BAKERIES WITH A HIGH BAKER'S ALLERGY AND ASTHMA BURDEN

^{1,2}Roslynn Baatjes, ¹Mohamed Jeebhay*. ¹Division of Occupational Medicine and Centre for Environmental and Occupational Health Research, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa; ²Department of Environmental and Occupational Studies, Faculty of Applied Sciences, Cape Peninsula University of Technology, Cape Town, South Africa

10.1136/oemed-2017-104636.235

Introduction Previous studies of supermarket bakery workers in South Africa demonstrated a high prevalence (13%) of baker's allergy/asthma and a 50% decrease in mean flour dust levels after an intervention. This study reports on the incidence of baker's allergy/asthma one year after the intervention.

Methods A group-randomised design assigned 31 bakeries to two intervention groups (IG) (bakery mixer lid and training) (n=244) and a control group (CG) (n=93). Health data prior to and after the intervention included symptom questionnaires; serum wheat IgE; and FeNO during the work shift using NIOX MINO.

Results Overall, the prevalence of work-related chest symptoms in the pre (25%) and post (27%) IGs were similar, but a 13% increase in the CG. Wheat sensitisation prevalence in IG (2% decrease) compared to the CG (3% increase). FeNO (geometric mean) in the IG decreased from 17.7 ± 2.3 ppb to 16.5 ± 2.2 ppb. There was a lower incidence of chest symptoms in IG (16%) compared to CG (23%). IG also had

significantly higher (14%) conversion to negative wheat sensitisation status compared to CG (0%). The mean FeNO decline in IG (-2.2 ppb) was similar to CG (-1.7 ppb). FeNO decline was greater in IG (-25.2 ppb) compared to CG (-13.6 ppb) when baseline FeNO >25 ppb. In a multivariate model for $>10\%$ FeNO decline, significant predictors included FeNO >25 ppb (OR:2.9) and atopic status (OR:2.0).

Conclusion This study demonstrates some evidence of an intervention effect one year after the intervention. The lack of a demonstrably strong effect can be attributable to the short follow up period.

Oral Presentation

Cancer

0286 OCCUPATIONAL EXPOSURE TO GASOLINE AND DIESEL EXHAUSTS AND THE RISK OF KIDNEY CANCER IN CANADIAN MEN

¹Cheryl Peters*, ²Marie-Elise Parent, ^{3,4}Shelley Harris, ^{3,4}Lidija Latifovic, ^{3,4}Linda Kachuri, ⁴Laura Bogaert, ¹Canadian Cancer Registries Group, ¹Paul Villeneuve. ¹Carleton University, Ottawa, Canada; ²Institut Armand-Frappier, Institut national de la recherche scientifique, Montreal, Canada; ³Cancer Care Ontario, Toronto, Canada; ⁴University of Toronto, Toronto, Canada

10.1136/oemed-2017-104636.236

Objectives The objective of this study was to investigate whether occupational gasoline and/or diesel exhaust exposure contribute to kidney cancer risk.

Methods The National Enhanced Cancer Surveillance System is a population-based case-control study conducted from 1994–1997 in Canada. Incident kidney cancer cases were identified using provincial registries, while the control series were identified through random digit dialling, or provincial datasets. Self-reported questionnaires obtained information on lifetime occupational history and cancer risk factors. Hygienists coded occupational histories for diesel and gasoline exhaust exposures using concentration, frequency, duration and reliability. Logistic regression was used to estimate odds ratios (OR) and 95% confidence intervals separately by exhaust type. Models were adjusted for age, province, BMI, and cigarette smoking.

Results Complete occupational data were available for 652 cases and 2368 controls. The majority of workers had been exposed to diesel (53%) or gasoline (55%) exhausts, respectively; most exposures were at low concentrations. Workers who had ever been exposed were significantly more likely to have kidney cancer than those who were never exposed (OR diesel: 1.4; 1.1–1.6, OR gasoline: 1.7; 1.4–2.1). When examining duration of exposure and tertiles of cumulative exposure, diesel and gasoline exposures were both linked to a significantly increased risk of kidney cancer ($p < 0.05$), but no exposure-response pattern was evident. Exposure to gasoline exhaust showed stronger positive relationships with kidney cancer than diesel.

Conclusions This study provides evidence that occupational gasoline and to a lesser extent, diesel exhaust exposure increases the risk of kidney cancer.