ORIGINAL ARTICLE

Examining lung cancer risks across different industries and occupations in Ontario, Canada: the establishment of the Occupational Disease Surveillance System

James K H Jung,^{1,2} Saul G Feinstein,¹ Luis Palma Lazgare,¹ Jill S Macleod,¹ Victoria H Arrandale,^{1,2} Christopher B McLeod,³ Alice Peter,⁴ Paul A Demers^{1,2,3}

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/oemed-2017-104926>).

¹Occupational Cancer Research Centre, Cancer Care Ontario, Toronto, Ontario, Canada

²Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada

³School of Population and Public Health, University of British Columbia, Vancouver, British Columbia, Canada

⁴Population Health and Prevention, Cancer Care Ontario, Toronto, Ontario, Canada

Correspondence to

Dr Paul A Demers, Occupational Cancer Research Centre, Cancer Care Ontario, Toronto, ON M5G 2L3, Canada; Paul.Demers@cancercare.on.ca

Received 30 November 2017

Revised 28 March 2018

Accepted 19 April 2018

Published Online First

7 May 2018

ABSTRACT

Background The Occupational Disease Surveillance System (ODSS) was established in Ontario, Canada by linking a cohort of workers with data created from Workplace Safety and Insurance Board (WSIB) claims to administrative health databases. The aim of this study was to use ODSS to identify high-risk industry and occupation groups for lung cancer in Ontario.

Methods Workers in the WSIB lost time claims database were linked to the Ontario Cancer Registry using subjects' health insurance numbers, name, sex, birthdate and death date (if applicable). Several occupations and industries known to be at increased risk were outlined a priori to examine whether ODSS could replicate these associations. Age-adjusted, sex-stratified Cox proportional hazard models compared the risk of lung cancer within one industry/occupation versus all other groups in the cohort. Workers with a lung cancer diagnosis prior to cohort entry were excluded for analysis, leaving 2 187 762 workers for analysis.

Results During the 1983 to 2014 follow-up, 34 661 workers in the cohort were diagnosed with lung cancer. Among expected high-risk industries, elevated risks were observed among workers in quarries/sand pits and construction industries for both sexes, and among males in metal mines, iron foundries, non-metallic mineral products industries and transportation industries. Excess risk was also observed among occupations in drilling/blasting, other mining/quarrying, mineral ore treating, excavating/grading/paving, truck driving, painting, bus driving and construction.

Conclusions This current surveillance system identified several established high-risk groups for lung cancer and could be used for ongoing surveillance of occupational lung cancer in Ontario.

INTRODUCTION

In Canada, lung cancer is fourth in cancer incidence for men and third in women and is the leading cause of cancer mortality in males and females.¹ Although lung cancer incidence is no longer increasing in Canada, the 5-year survival rate remains low (17%).¹ As lung cancer is primarily caused by smoking, prevention efforts have largely focused on tobacco control. However, while approximately 15% of lung cancer cases have been attributed to occupational exposures,² such hazards have

Key messages

What is already known about this subject?

- About 15% of lung cancer cases can be attributed to workplace exposures, and a number of occupations are exposed to known carcinogens such as asbestos, silica dust and diesel engine exhaust.

What are the new findings?

- The Occupational Disease Surveillance System (ODSS) confirmed many well-established high-risk groups for lung cancer such as those in construction, mining and transportation.
- Other high-risk groups included workers in industries of defence services and occupations such as metal forging workers, janitors/charworkers/cleaners and motor vehicle fabricating/assembling positions.

How might this impact on policy or clinical practice in the foreseeable future?

- ODSS can provide quantitative risk estimates for occupational diseases at a population level that are necessary to support the targeting of work-related disease prevention programmes.

typically not been a priority in cancer prevention policies.³ A focus on lung cancer prevention beyond smoking could further reduce future incidence.⁴

The International Agency for Research on Cancer (IARC) has identified many occupational exposures such as asbestos, silica dust, chromium VI compounds, diesel engine exhaust, nickel, paint and welding fumes as lung carcinogens with sufficient evidence in humans.⁵ As such, there is strong evidence linking excess lung cancer risk to employment in occupations including construction⁶; quarries, sand pits and mining⁷; transportation^{8,9}; painting¹⁰ and welding,¹¹ even after adjusting for smoking.

As ongoing monitoring is needed for disease prevention, some attempts have been made to establish surveillance systems for occupational cancers. Internationally, the best models have been based on large-scale record linkages. An example is the Nordic Occupational Cancer (NOCCA) surveillance system,¹² where occupational information from

► <http://dx.doi.org/10.1136/oemed-2018-105189>

To cite: Jung JKH, Feinstein SG, Palma Lazgare L, et al. *Occup Environ Med* 2018;**75**:545–552.

five Nordic countries (Denmark, Finland, Iceland, Norway and Sweden) was obtained through census data, which was then linked to national cancer registries. However, census linkages are only possible in a small number of countries where complete occupational information is collected in linkable form. In North America, the surveillance of occupational cancers has largely been based on cause-specific mortality patterns by occupation and industry using death certificate records.¹³ However, this form of surveillance can be limited by the lack of accuracy in recording occupational information (ie, they reflect the most recent occupation at time of death and not at time of exposure). Moreover, death records cannot be used to study non-fatal cancers.

In Canada, the existence of a single payer health system and the maintenance of comprehensive administrative health databases helps facilitate health surveillance for the population. Yet despite the availability of high-quality information on new disease diagnoses, the initiation of occupational disease surveillance continues to be limited by the absence of information on occupation and industry of employment in the administrative health data. Koehoorn *et al*¹⁴ addressed this limitation by identifying industry of employment using data from the provincial health registry for British Columbia, Canada, which identify the name of employers for people whose premiums are employer paid. This database was linked to administrative health data to explore high-risk industries for asthma. Cherry *et al*¹⁵ used an alternative approach to overcome this challenge by linking workers' compensation claims—which contained data on occupation and industry of employment at the time of injury or illness resulting in the claim—to physician billing records for asthma in Alberta, Canada. Using unique personal health numbers to match the claims to the provincial administrative health database, occupations with higher than expected rates of asthma were identified.

These linkage approaches created an avenue of performing occupational disease surveillance that could be used as a model for other jurisdictions. The Occupational Disease Surveillance System (ODSS) was developed in the province of Ontario, Canada to link workers' information on occupation and industry (based on workers' compensation claims) to administrative health databases. While ODSS was developed for the surveillance of a variety of occupational diseases, the first focus was to examine occupational cancers through a linkage to the provincial cancer registry. Specifically, lung cancer was the first occupational disease examined using ODSS as it is one of the most common occupational cancers with a well-established set of occupational risk factors.¹⁶

The objective of this study was to use ODSS to identify high-risk industry and occupation groups for lung cancer in Ontario, Canada. We first examined the effectiveness of ODSS in identifying known high-risk groups for lung cancer, which could provide support for this project as an ongoing surveillance system for occupational disease. Additionally, ODSS was used to investigate other suspected occupational associations for lung cancer, which could serve as a basis for future research.

METHODS

The ODSS was established through the linkage of a cohort of workers created with data from Ontario Workplace Safety and Insurance Board (WSIB) claims with administrative health databases (see figure 1). The system acts as a prospective open occupational cohort since additional workers can enter the cohort through new WSIB claims over time. This study obtained ethics approval from the University of Toronto Health Sciences Research Ethics Board.

Creation of the ODSS cohort

The WSIB claims database includes all accepted lost time claims (applications submitted when employees suffer a work-related injury or disease that can lead to permanent disability and/or loss of wages) by workers in Ontario, providing the occupation and industry data from 1983 to 2014. Claims preceding 1983 were more likely to have missing data for key linkage variables and were excluded. The database includes: occupation/industry code (specific to employment related to the compensation claim), claim date, the nature of work-related injuries and/or illnesses, worker's name, sex, birthdate and death date (if applicable). In this database, industry was coded according to the Canadian Standard Industry Classification 1970 (SIC-70) and 1980 (SIC-80) codes. For analytical purposes, all industry codes were converted to SIC-70, where industries could be classified by division,¹⁰ major (49) or minor (248) groups. Occupations were coded according to the Canadian Classification Dictionary of Occupation (CCDO) at the 4-digit level. Using these codes, occupations could be classified by division,¹⁷ major (88) or minor (487) groups.

An internal linkage of the WSIB claims database was completed to link multiple claims records belonging to the same worker using the AutoMatch V4.2 (MatchWare Technologies, Inc., Silver Spring, Maryland, USA) probabilistic record linkage software. This application was used to group multiple records for a single worker within a file and generate a unique identifier. Of 3 833 772 WSIB claims between 1983 and 2014, we identified 2 253 734 unique workers.

Exclusion criteria

Workers were excluded due to:

- ▶ Missing sex
- ▶ Missing/invalid birthdate
- ▶ Age at claim under 15 years old
- ▶ Invalid claim date
- ▶ Missing/invalid both the SIC-70 and CCDO codes.

This resulted in 2 190 246 unique workers included as the ODSS worker cohort.

Linkage with administrative health data

Workers from the WSIB claims database were linked to the Registered Persons Database (RPDB; 1991–2015). This database contains information for individuals who are registered for insured health services in the province through the Ontario Health Insurance Plan (OHIP). The RPDB contains information on sex, residence location, birthdate, death date (if applicable), date of emigration from province (if applicable) and health insurance number (HIN). Information on name, sex, birthdate and death date was used to link workers from the WSIB claims database (n=2 253 734) with the RPDB (n=16 162 277).

OHIP-registered persons are assigned a unique 10-digit HIN, and this identifier can be used to match individuals to other administrative health databases. Using a combination of the identifying information listed above, approximately 82% of all workers in the ODSS cohort (1 796 731 unique subjects) were linked to HINs.

Ontario Cancer Registry (OCR)

The OCR maintained by Cancer Care Ontario (n=2 050 907)¹⁸ contains information on incident cancer cases in Ontario (1964–2016), excluding non-melanoma skin cancers. The registry collects data from several sources including hospital discharge and day surgery records, pathology reports from hospital and

*Abbreviations:

- HIN = Health Insurance Number
- SIC = Standard Industry Classification (1970) CCDO = Canadian Classification Dictionary of Occupation

Figure 1 Establishing the ODSS worker cohort through the linkage of administrative health databases. ODSS, Occupational Disease Surveillance System.

community laboratories, records from regional cancer centres and death certificates received from the Registrar General. Incident cancers were coded according to the International Classification of Disease 10th edition.¹⁹

Following the linkage to the RPDB, the workers were linked to the OCR deterministically using the subjects' HIN, if available. Where no HIN was available, the linkage was probabilistic, using name, sex, birthdate and death date. In total, 214 821 unique workers in the ODSS cohort with at least one cancer diagnosis were linked to the OCR. Only the first tumour at the primary site (malignant cases only) was considered.

Statistical analysis

For cohort members, the date of entry into the cohort was the date of their first accepted lost time WSIB compensation claim. Follow-up continued until cancer diagnosis, year of death, emigration out of province or administrative censoring (31 December 2016), whichever occurred first. Workers with a lung cancer diagnosis prior to cohort entry were excluded from the analysis (leaving 2 187 762 workers for analysis). Disease risk was estimated with Cox proportional hazard models using a rolling cohort analysis, where the risk of cancers within one industry or occupation group was compared with the rest of the cohort.⁸ HRs and 99% CIs were calculated for each occupation and industry group, adjusted for year of birth. Sex-stratified analyses were also conducted. Results were reported for only the groups that met the reporting threshold of at least six cases. All analyses were completed using SAS V.9.4 (SAS Institute).

A priori selection of groups for analyses

Known associations between workplace exposures and lung cancer risks were identified a priori. These served as a guideline to examine whether ODSS could accurately identify established high-risk groups and demonstrate its utility as a valid surveillance tool. The senior author (PAD), an occupational epidemiologist with expertise in industrial hygiene, performed the selection of the groups. Groups with known occupational exposures to carcinogens classified by IARC as Group 1 (sufficient evidence in humans) were identified as a priori groups. Other groups of interest included those with suspected increased risk due to occupational exposures, based on evidence from scientific literature and/or expert knowledge. In total, 37 industries and 49 occupations were preidentified for this analysis, selected from the major and minor levels. Additionally, the results for all industries and occupations at the division level were provided.

RESULTS

Among the 2 187 762 ODSS cohort workers (66% males, n=1 440 683), 34 661 workers with lung cancer were identified during the 1983 to 2014 follow-up. The mean age at cohort entry was 37.4 years (SD 12.3). Table 1 provides the risk estimates for all industries and occupations at the division level. Tables 2 and 3 provide the risk estimates for the a priori groups and other suspected groups of interest, and table 4 summarises the a priori groups that showed positive associations for lung cancer as expected. The results for all industry and occupation

Workplace

Table 1 HRs for lung cancer among all industries and occupations, division level

Industry/occupation	Cases (no of workers)	HR (99% CI)
Industry		
Agriculture	379 (35 120)	0.81 (0.71 to 0.92)
Forestry, fishing and trapping	249 (10 721)	1.15 (0.97 to 1.35)
Mines (including milling), quarries and oil wells	798 (23 204)	1.43 (1.30 to 1.57)
Manufacturing	12 692 (694 723)	0.98 (0.95 to 1.01)
Construction	3648 (211 549)	1.09 (1.04 to 1.14)
Transportation, communication and other utilities	3533 (197 529)	1.14 (1.09 to 1.19)
Wholesale and retail trade	5396 (429 870)	0.99 (0.95 to 1.03)
Finance, insurance and real estate	479 (24 020)	1.15 (1.02 to 1.30)
Community, business and personal service	7266 (600 256)	0.95 (0.92 to 0.99)
Public administration and defence	3311 (191 092)	1.04 (1.00 to 1.09)
Occupation		
Managerial, administrative and related	347 (31 028)	1.03 (0.90 to 1.18)
Natural sciences, engineering and mathematics	287 (26 366)	0.85 (0.73 to 0.99)
Social sciences and related fields	250 (30 721)	1.04 (0.88 to 1.22)
Teaching and related	335 (48 506)	0.61 (0.53 to 0.71)
Medicine/health	1462 (135 434)	0.85 (0.79 to 0.91)
Artistic, literary, recreational and related	131 (15 003)	1.11 (0.88 to 1.39)
Clerical and related	2824 (197 515)	1.09 (1.04 to 1.15)
Sales	1515 (148 237)	1.06 (0.99 to 1.14)
Service	5389 (371 247)	1.06 (1.01 to 1.10)
Farming, horticultural and animal husbandry	555 (50 265)	0.88 (0.79 to 0.98)
Fishing, hunting, trapping and related occupations	11 (558)	1.16 (0.53 to 2.52)
Forestry/logging	216 (10 707)	1.04 (0.87 to 1.24)
Mining/quarrying including oil/ gas field	441 (13 034)	1.47 (1.29 to 1.66)
Processing—metal/clay, glass, stone/chemicals	1427 (79 369)	1.18 (1.10 to 1.26)
Processing—food/wood/pulp/ textile	1428 (99 358)	1.00 (0.93 to 1.07)
Machining and related	3615 (189 751)	1.11 (1.06 to 1.16)
Product fabricating, assembling and repairing	5816 (328 720)	1.01 (0.98 to 1.05)
Construction trades	4076 (216 079)	1.09 (1.05 to 1.14)
Transport equipment operating	3672 (168 429)	1.39 (1.33 to 1.46)
Materials handling and related, NEC	2522 (153 276)	1.11 (1.05 to 1.17)
Other crafts and equipment operating	572 (28 350)	1.12 (1.00 to 1.25)
Occupation NEC	3410 (215 574)	1.10 (1.05 to 1.16)

NEC, not elsewhere classified.

groups, stratified by sex, can be found in online supplementary tables 1–6.

Industry groups

At the division level, the highest risks were observed in mining/quarries, transportation/communication/other utilities,

Table 2 HRs for lung cancer among a priori industry groups and other suspected industries of interest (at the major level unless otherwise noted)

Industry	Cases (no of workers)	HR (99% CI)
A priori groups		
Metal mines	578 (13 859)	1.46 (1.31 to 1.63)
Non-metal mines	35 (1089)	1.44 (0.93 to 2.22)
Quarries/sand pits	110 (4025)	1.60 (1.25 to 2.05)
Rubber products*	227 (13 041)	0.97 (0.82 to 1.15)
Iron/steel mills*	641 (21 695)	1.07 (0.97 to 1.19)
Iron foundries*	168 (6168)	1.28 (1.05 to 1.56)
Smelting/refining*	38 (983)	1.16 (0.77 to 1.77)
Shipbuilding/repair*	24 (907)	1.05 (0.62 to 1.78)
Non-metallic mineral products	585 (24 915)	1.22 (1.10 to 1.36)
General contractors	1448 (74 814)	1.17 (1.09 to 1.25)
Special-trade contractors	2367 (145 370)	1.05 (0.99 to 1.11)
Transportation	2523 (136 038)	1.21 (1.15 to 1.28)
Blacksmithing/welding shops*	71 (3750)	1.24 (0.91 to 1.68)
Other suspected groups of interest		
Agriculture	379 (35 120)	0.81 (0.71 to 0.92)
Forestry	242 (10 300)	1.15 (0.98 to 1.36)
Services incidental to mining	87 (4643)	1.13 (0.86 to 1.49)
Tobacco products	35 (1117)	1.27 (0.82 to 1.96)
Plastics fabricating*	502 (40 883)	0.91 (0.81 to 1.03)
Leather	79 (5685)	0.71 (0.53 to 0.95)
Textile	369 (19 316)	1.06 (0.93 to 1.21)
Wood	542 (38 173)	1.03 (0.92 to 1.15)
Furniture/fixture	357 (30 257)	0.69 (0.60 to 0.79)
Paper	512 (25 284)	1.03 (0.92 to 1.16)
Printing/publishing	450 (25 996)	1.13 (1.00 to 1.28)
Metal fabricating	2038 (127 741)	1.01 (0.95 to 1.07)
Machinery (except electrical)	856 (41 851)	1.04 (0.95 to 1.14)
Electrical products	824 (38 553)	1.08 (0.99 to 1.18)
Petroleum/coal products	17 (1 158)	0.74 (0.40 to 1.38)
Chemical/chemical products	357 (21 553)	0.96 (0.84 to 1.10)
Paint/varnish manufacturers*	37 (2054)	0.90 (0.59 to 1.38)
Storage	136 (10 628)	1.23 (0.98 to 1.53)
Electric power, gas and water utilities	415 (22 415)	1.03 (0.91 to 1.17)
Wholesale trade	2036 (151 240)	1.05 (0.99 to 1.11)
Motor vehicle repair shops*	476 (36 454)	1.05 (0.93 to 1.19)
Health/welfare services	2381 (178 587)	0.94 (0.89 to 0.99)
Restaurants, caterers and taverns*	1042 (125 953)	0.88 (0.81 to 0.96)
Defence services*	187 (5736)	1.38 (1.14 to 1.66)

*Minor industry group.

construction and finance/insurance/real estate industries. Agriculture was the only industry to show significantly reduced risks (table 1).

A priori industry groups that showed significantly increased risks included quarries/sand pits, metal mines, iron foundries, non-metallic mineral products, transportation and general contractors (table 2). Based on sex-stratified results, metal mines, non-metallic mineral products industries, transportation industries and general contractors reached significance only in males (although the first three still showed excess risk in females) (Supplementary file). Among male general contractors, highway/bridge/street construction (HR 1.61, 99% CI 1.39 to 1.87) and

Table 3 HRs for lung cancer among a priori occupation groups and other suspected occupations of interest (at the major level unless otherwise noted)

Occupation	Cases (no of workers)	HR (99% CI)
A priori groups		
Bartenders*	72 (4431)	1.62 (1.19 to 2.19)
Waiters, hostesses and stewards*	321 (32 038)	0.96 (0.83 to 1.11)
Drilling/blasting	165 (3482)	1.61 (1.32 to 1.97)
Other mining/quarrying including oil/gas field	333 (10 522)	1.53 (1.33 to 1.76)
Mineral ore treating	42 (983)	1.54 (1.04 to 2.29)
Metal smelting, converting and refining*	70 (2310)	1.15 (0.84 to 1.56)
Clay, glass and stone processing/forming	255 (10 187)	1.24 (1.05 to 1.46)
Welding/flame cutting*	904 (47 110)	1.08 (0.99 to 1.18)
Boilermakers, platers and structural metal workers*	98 (4432)	1.85 (1.43 to 2.40)
Painting/decorating, except construction*	221 (11 466)	1.25 (1.05 to 1.48)
Excavating/grading/paving	536 (18 394)	1.52 (1.36 to 1.70)
Construction trades	3057 (165 414)	1.09 (1.03 to 1.14)
Railway transport operating	97 (4033)	1.23 (0.95 to 1.60)
Bus drivers*	239 (7733)	1.23 (1.04 to 1.46)
Truck drivers	3011 (122 525)	1.48 (1.41 to 1.56)
Stationary engine/utilities equipment operating	207 (7596)	1.24 (1.04 to 1.48)
Other suspected groups of interest		
Nursing therapy and related assisting	1369 (124 139)	0.85 (0.79 to 0.91)
Radiological technologists/technicians*	10 (1305)	0.88 (0.39 to 1.99)
Firefighting	174 (12 076)	0.92 (0.76 to 1.12)
Police/detectives	185 (23 360)	0.67 (0.56 to 0.82)
Guards/watchmen*	426 (23 879)	1.29 (1.14 to 1.46)
Chefs/cooks*	522 (46 128)	0.98 (0.87 to 1.09)
Barbers, hairdressers and related*	12 (1304)	0.87 (0.41 to 1.82)
Laundering/dry cleaning	119 (7115)	1.02 (0.81 to 1.30)
Janitors/charworkers/cleaners	2179 (103 022)	1.25 (1.18 to 1.32)
Farmers/farm management	45 (4389)	0.69 (0.47 to 1.01)
Farm, nursery and related	498 (43 787)	0.92 (0.82 to 1.03)
Forestry/logging	216 (10 707)	1.04 (0.87 to 1.24)
Chemicals, petroleum, rubber, plastic processing	556 (41 570)	1.09 (0.97 to 1.21)
Wood processing except paper pulp	113 (6664)	1.18 (0.92 to 1.50)
Pulp/papermaking and related	100 (5710)	0.92 (0.71 to 1.19)
Textile processing	169 (9782)	1.11 (0.91 to 1.35)
Tobacco processing*	11 (433)	1.11 (0.51 to 2.41)
Metal machining	808 (41 797)	1.07 (0.98 to 1.18)
Forging*	63 (2529)	1.64 (1.19 to 2.27)
Wood machining	105 (8749)	0.87 (0.67 to 1.11)
Engravers, etchers and related*	12 (995)	1.85 (0.88 to 3.89)
Patternmakers/mouldmakers*	23 (1537)	1.31 (0.77 to 2.24)
F/A: motor vehicle*	917 (53 476)	1.19 (1.09 to 1.30)
F/A: electrical/related equipment	673 (38 180)	1.11 (1.00 to 1.22)
F/A: wood products	304 (24 588)	0.76 (0.65 to 0.88)
F/A: textile, fur and leather products	417 (26 683)	0.74 (0.65 to 0.84)

continued

Table 3 continued

Occupation	Cases (no of workers)	HR (99% CI)
F/A: rubber, plastic and related products	259 (15 244)	0.92 (0.78 to 1.08)
Motor vehicle mechanics*	946 (59 343)	1.02 (0.93 to 1.11)
Paper product F/A*	129 (6840)	1.12 (0.89 to 1.40)
Water transport operating	44 (2694)	0.98 (0.66 to 1.44)
Taxi drivers/chauffeurs*	82 (3427)	1.18 (0.89 to 1.57)
Printing press*	140 (7853)	1.09 (0.88 to 1.36)
Bookbinders and related*	68 (3626)	1.25 (0.91 to 1.71)

*Minor occupation group.
F/A, fabricating/assembling.

other construction (HR 1.32, 99%CI 1.13 to 1.55) showed significantly increased risks. Among males in the mining industries, employment in uranium mines (HR 1.93, 99%CI 1.50 to 2.47) showed the highest risk. Although employment in asbestos mines showed significantly elevated risks, the number of total cases was below the reporting threshold.

Among other suspected industry groups of interest, the only industry that showed significantly increased risks was defence services.

Occupation groups

The division level results show that the highest risks were observed among occupations in mining/quarrying, transport equipment operating, metal/clay/glass/stone/chemicals processing, construction trades and machining (table 1). Teaching and medicine/health occupations showed the lowest risks.

A priori occupation groups that showed significantly increased risks included boilermakers/platers/structural metal workers, drilling/blasting, mineral ore treating, other mining/quarrying, excavating/grading/paving, bartending, truck driving, painting/decorating, stationary engine/utilities equipment operating, bus driving and construction. Based on sex-specific results, boilermakers/platers/structural metal workers reached significance only among males, bartenders and painting/decorating positions reached significance only in females and neither sex reached significance when stratified for mineral ore treating positions and bus drivers (Supplementary file). However, both sexes still showed excess risk in all of these groups. Among the construction trades, those with significantly increased risks included insulating occupations (HR 2.52, 99%CI 1.71 to 3.71), painters/paperhangers (HR 1.41, 99%CI 1.18 to 1.69), structural metal erectors (HR 1.37, 99%CI 1.05 to 1.79) and plasterers (HR 1.30, 99%CI 1.03 to 1.65) for males. Although glaziers showed significantly elevated risks among females, the number of cases in these groups was below the reporting threshold.

Other occupations of interest with significantly elevated risks included metal forging workers, janitors/charworkers/cleaners and motor vehicle fabricating/assembling positions.

DISCUSSION

The ODSS identified positive associations for the majority of industries (12/13) and occupations (15/16) selected a priori to have an increased risk of lung cancer, although not all reached statistical significance. Pukkala and colleagues reported similar findings in their paper on the NOCCA surveillance system,¹² which obtained occupational information from census data. Similar to ODSS, NOCCA observed significantly elevated risks for lung cancer among miners, quarry workers, construction

Table 4 Summary of industry and occupation groups selected a priori expected to be at high risk

Industry/occupation	Exposure	Overall		Male		Female	
		Association	Probability	Association	Probability	Association	Probability
Industry							
Metal mines	Silica, asbestos, diesel exhaust, nickel	+	**	+	***	++	.
Non-metal mines	Silica, asbestos, diesel exhaust, radon	+	***	+	*	X	X
Quarries/sand pits	Silica, asbestos, diesel exhaust	++	***	++	***	X	X
Rubber products industries	Dust/fumes/chemicals from rubber production	Null	.	—	.	+	.
Iron/steel mills	Silica, PAHs	+	**	+	.	+	.
Iron foundries	Silica, PAHs	+	***	+	**	X	X
Smelting/refining	Cadmium, arsenic, nickel	+	.	+	.	X	X
Shipbuilding/repair	Asbestos	+	.	+	.	X	X
Non-metallic mineral products	Silica	+	***	+	***	+	.
General contractors	Silica, asbestos, diesel exhaust	+	***	+	***	Null	.
Special-trade contractors	Silica, asbestos, diesel exhaust	+	***	Null	.	+	*
Transportation	Diesel exhaust	+	***	+	***	+	.
Blacksmithing/welding shops	Welding fumes	+	*	+	.	X	X
Occupation							
Bartenders	Environmental tobacco smoke	++	***	+	.	+++	***
Waiters, hostesses, stewards	Environmental tobacco smoke	Null	*	—	*	Null	.
Drilling/blasting	Silica	++	***	++	***	X	X
Other mining/quarrying	Silica, asbestos, radon	++	***	++	***	+++	***
Mineral ore treating	Silica, asbestos, radon	++	**	+	*	X	X
Metal smelting, converting, refining	Cadmium, arsenic, nickel	+	.	+	.	X	X
Clay, glass, stone processing/forming	Silica	+	***	+	**	+	.
Welding/flame cutting	Welding fumes	+	**	+	.	+	.
Boilermakers, platers, structural metal workers	Asbestos	++	***	++	***	X	X
Painting/decorating	Paint pigments, filaments, resin	+	***	+	.	+++	***
Excavating/grading/paving	Silica	++	***	++	***	+++	**
Construction trades	Silica, asbestos, diesel exhaust	+	***	+	**	++	***
Railway transport operating	Diesel exhaust	+	*	+	.	X	X
Bus drivers	Diesel exhaust	+	**	+	*	+	*
Truck drivers	Diesel exhaust	+	***	+	***	+	***
Stationary engine/utilities equipment	Diesel exhaust	+	***	+	**	X	X

Association scale based on HRs ---<0.50, --0.50–0.66, –0.67–0.94, Null 0.95–1.04, +1.05–1.49, ++1.50–1.99, +++>2.00.

Probability scale based on p values. Not significant. *<0.05, **<0.01, ***<0.0001.

x, data not provided as they were under reporting threshold of >5 cases.

PAH, polycyclic aromatic hydrocarbon.

workers, transportation workers, painters and bartenders—all of which have known exposures to Group 1 carcinogens as outlined by IARC.

Work related to mining, quarrying and construction has consistently shown increased risks due to workplace exposures to silica dust, asbestos, radon and diesel engine exhaust.^{20–22} Transport workers are also exposed to diesel engine exhaust and thus have shown increased risk of lung cancer in other studies in North America.^{8,9} The increased risk in painting occupations in ODSS and NOCCA can be attributed to their exposures to various hazardous chemicals in paint pigments, resins and filaments.²³

Historically, employees working in bars and restaurants such as bartenders and waiters have shown increased risk of lung

cancer partially due to exposures to secondhand smoking.¹⁷ Smoking in public settings such as bars was not banned in Ontario until 2006,²⁴ which may explain the elevated risk among bartenders (although only females reached significance) in ODSS. Increased risks were not observed in waiters and waitresses in ODSS, although it has been suggested that bartenders have a twofold to fourfold increase in secondhand smoke exposure compared with table waiting staff.²⁵ Exposures to silica dust likely explain the significantly elevated risks in the other a priori groups, including iron foundries, non-metallic mineral products industries, clay/glass/stone processing and forming occupations and excavating/grading/paving positions.^{21,23} Finally, male boilermakers showed significantly increased risks in ODSS, which

can be explained by their known occupational exposures to asbestos.²⁶

The results from this study add to the current understanding that workers in these fields are likely at higher risk of lung cancer due to occupational exposures. However, there were certain industries and occupations that did not show significantly elevated risks in ODSS despite having known exposures to IARC Group 1 carcinogens for lung cancer. Although construction workers overall had an excess risk of lung cancer, some specific roles (eg, brick and tile setting, concrete finishing, roofing) did not show increased risk. The discrepancy may be due to the possible low WSIB coverage in these construction occupations, as complete coverage across all construction industries has only recently become mandatory.²⁷

In addition, higher risks were also expected in smelting occupations and industries due to metals such as copper, arsenic and nickel; welders due to the welding fumes that may comprise carcinogens such as nickel, chromium VI and cadmium; rubber products industries due to dust, fumes and chemicals from rubber production (which involve N-nitrosamines and polycyclic aromatic hydrocarbons (PAHs)); iron/steel mills due to silica dust and shipbuilding industries due to asbestos.^{21 28} While the results did not reach statistical significance, ODSS was able to identify trends in increased risk in these established high-risk groups.

Several groups with suspected work-related associations for lung cancer based on prior research were also observed to be at increased risk. For example, the potential exposures to metals, PAHs and asbestos among forging workers²⁹ could likely explain the increased risk observed in ODSS. The increased risks in janitors may be due to asbestos exposure in the buildings of their workplace. Furthermore, motor vehicle assembly workers may be at elevated risk due to exposures to paint products and asbestos from brake installation.³⁰ The increased risk among the defence services industry was more challenging to understand, as active military personnel are not part of the provincial workers' compensation plan used to create the cohort. While occupational associations for lung cancer are possible, exposures in defence services are diverse, and thus, it is difficult to identify the exact exposure agent(s) responsible. Lastly, while the results did not reach statistical significance, elevated risks were observed among workers in storage and printing industries, possibly due to exposures to diesel exhaust³¹ and chromium VI compounds,³² respectively. Future epidemiological studies could help ascertain whether the high risks in these groups may indeed be attributable to exposures to specific agent(s) at the workplace.

The overall division level results showed that generally, the highest risks were observed among blue-collar occupations (eg, mining, construction and transportation), as expected. The reduced risk among the agriculture industry has been well observed previously and is likely due to the lower rates of smoking among workers in this field.^{33 34} The reduced risks in occupations such as teaching and medicine/health may also reflect the effects of socioeconomic status on risk for lung cancer,³⁵ as well as differences in smoking habits. However, discrepancies existed, such as the significantly elevated risks observed among the finance/insurance/real estate industries. Lastly, there were some differences between sexes in the results, where certain industries or occupations showed increased risk in males but not for females, and vice versa. In some cases, the number of female workers may have been too small to detect a difference for some groups. The possibility that males and females had different roles or exposures cannot be dismissed.³⁶ These discrepancies demonstrate further avenues for investigating work-related risks.

Limitations and strengths

One limitation of the study was that only the industry and occupation specific to the time of the compensation claim was available, and employment at this time may not necessarily correspond to lifetime work history. While approximately one-third of the ODSS worker cohort had multiple claims, this was not sufficient data to reconstruct their full-work history. The dates or the duration of employment in a particular job could also not be determined. As a surveillance system, measurements of workplace exposures to potential carcinogens could not be obtained, which can vary between and within jobs and industries, as well as over time.

Information on smoking status, which may be more common among blue-collar workers,³⁷ was also not available. However, it has been suggested that the absence of smoking data does not make a large difference in the risk estimates derived for associations between various occupation groups and lung cancer, particularly when there is a large and heterogeneous reference population.^{38 39} Another limitation was that the WSIB covers 70 to 75 per cent of workers in Ontario.⁴⁰ Thus, we were unable to capture relationships among non-covered industries such as trade unions, private day cares and travel agencies. Lastly, the analyses may have been susceptible to finding spurious associations between industries/occupations and lung cancer due to multiple testing. However, we applied a strict significance threshold of 99% to try to reduce these associations and primarily did not observe significantly elevated risks in industries and occupations without any plausible occupational risk for lung cancer.

A key strength of this study was the large sample of both male and female workers in Ontario. The occupational information was obtained before disease diagnosis, reducing information bias from data collection. Moreover, the authors believe that the occupational information on WSIB claims is collected carefully resulting in high data quality. Lastly, this surveillance system compared workers to workers, rather than workers to the general population, minimising the healthy worker effect and providing a comparison group that is more similar in social class and overall health.

CONCLUSION

This current study identified various high-risk occupation and industry groups for lung cancer in Ontario, supporting previous findings and demonstrating that ODSS can identify groups that are known to be at increased risk of occupational diseases such as lung cancer. The quantitative disease risk estimates obtained by this system can provide useful data to support the implementation of prevention programmes for lung cancer caused by workplace exposures. In the future, ODSS will be used to explore patterns of increased risk of other occupational diseases, including different cancers but also conditions such as silicosis, asthma, carpal tunnel syndrome and Raynaud's syndrome.

This work further demonstrates that the usage of workers' compensation claims, in conjunction with other administrative health databases, is a valid and feasible approach for occupational disease surveillance. Other jurisdictions with access to similar types of administrative health databases can look to implement similar approaches to conduct occupational disease surveillance.

Acknowledgements The authors would like to thank Nelson Chong for conducting the data linkage.

Contributors JKHJ drafted the manuscript, interpreted the findings and produced the figures/tables. SGF assisted in the manuscript writing and provided methodological expertise regarding the development of ODSS. JKHJ, LPL and JSM

conducted the analyses. PAD, VHA, CBM and AP provided expertise in occupational health and administrative health data. PAD performed the selection of the a priori groups, conceived the study and provided overall supervision. All authors assisted in editing the manuscript.

Funding This project was funded by the Ministry of Labour (no 14-R-29) and the Public Health Agency of Canada (no 1516-HQ-000066). The Occupational Cancer Research Centre is supported from core funding from the Canadian Cancer Society and the Ontario Ministry of Labour.

Competing interests None declared.

Patient consent Not required.

Ethics approval Health Sciences Research Ethics Board, University of Toronto.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Canadian Cancer Society, Statistics Canada, Public Health Agency of Canada. *Canadian cancer statistics 2017*. Ottawa: Canadian Cancer Society, 2017.
- 2 Rushton L, Hutchings SJ, Fortunato L, et al. Occupational cancer burden in Great Britain. *Br J Cancer* 2012;107(Suppl 1):S3–7.
- 3 Occupational and Environmental Working Group for the Toronto Cancer Prevention Coalition. *Preventing occupational and environmental cancer: a strategy for Toronto*. Toronto: City of Toronto, 2001.
- 4 Canadian Cancer Society, Statistics Canada, Public Health Agency of Canada. *Canadian cancer statistics 2015*. Ottawa: Canadian Cancer Society, 2015.
- 5 International Agency for Research on Cancer. Agents classified by the IARC monographs, volumes 1–118 2017. <http://monographs.iarc.fr/ENG/Classification/> (accessed 29 May 2017).
- 6 Consonni D, De Matteis S, Pesatori AC, et al. Lung cancer risk among bricklayers in a pooled analysis of case-control studies. *Int J Cancer* 2015;136:360–71.
- 7 Taeger D, Pesch B, Kendzia B, et al. Lung cancer among coal miners, ore miners and quarrymen: smoking-adjusted risk estimates from the synergy pooled analysis of case-control studies. *Scand J Work Environ Health* 2015;41:467–77.
- 8 Garshick E, Laden F, Hart JE, et al. Lung cancer and vehicle exhaust in trucking industry workers. *Environ Health Perspect* 2008;116:1327–32.
- 9 Pintos J, Parent ME, Richardson L, et al. Occupational exposure to diesel engine emissions and risk of lung cancer: evidence from two case-control studies in Montreal, Canada. *Occup Environ Med* 2012;69:787–92.
- 10 Ramanakumar AV, Parent ME, Richardson L, et al. Exposures in painting-related occupations and risk of lung cancer among men: results from two case-control studies in Montreal. *Occup Environ Med* 2011;68:44–51.
- 11 Kendzia B, Behrens T, Jöckel KH, et al. Welding and lung cancer in a pooled analysis of case-control studies. *Am J Epidemiol* 2013;178:1513–25.
- 12 Pukkala E, Martinsen JI, Lynge E, et al. Occupation and cancer - follow-up of 15 million people in five Nordic countries. *Acta Oncol* 2009;48:646–790.
- 13 Dubrow R, Sestito JP, Lulich NR, et al. Death certificate-based occupational mortality surveillance in the United States. *Am J Ind Med* 1987;11:329–42.
- 14 Koehoorn M, Tamburic L, McLeod CB, et al. Population-based surveillance of asthma among workers in British Columbia, Canada. *Chronic Dis Inj Can* 2013;33:88–94.
- 15 Cherry N, Beach J, Burstyn I, et al. Data linkage to estimate the extent and distribution of occupational disease: new onset adult asthma in Alberta, Canada. *Am J Ind Med* 2009;52:831–40.
- 16 World Health Organization. Environmental and occupational cancers 2011. <http://www.who.int/mediacentre/factsheets/fs350/en/> (accessed 19 June 2017).
- 17 Jensen JA, Schillo BA, Moilanen MM, et al. Tobacco smoke exposure in nonsmoking hospitality workers before and after a state smoking ban. *Cancer Epidemiol Biomarkers Prev* 2010;19:1016–21.
- 18 Cancer Care Ontario. Ontario Cancer Registry 2015. <https://www.cancercare.on.ca/ocs/csurv/ocr/> (accessed 19 May 2017).
- 19 World Health Organization. Classification of Diseases (ICD) 2016. <http://www.who.int/classifications/icd/en/> (accessed 19 May 2017).
- 20 Driscoll T, Nelson DL, Steenland K, et al. The global burden of disease due to occupational carcinogens. *Am J Ind Med* 2005;48:419–31.
- 21 International Agency for Research on Cancer. IARC monographs on the evaluation of the carcinogenic risks of chemicals to humans: Arsenic, metals, fibres, and dusts. Volume 100C. Lyon: IARC; 2012. <http://monographs.iarc.fr/ENG/Monographs/vol100C/index.php>.
- 22 International Agency for Research on Cancer. IARC monographs on the evaluation of the carcinogenic risks of chemicals to humans: Radiation. 2012. Lyon: IARC. Volume 100D. <http://monographs.iarc.fr/ENG/Monographs/vol100D/>.
- 23 International Agency for Research on Cancer. IARC monographs on the evaluation of the carcinogenic risks of chemicals to humans: Chemical agents and related occupations. Volume 100F. Lyon: IARC; 2012. <http://monographs.iarc.fr/ENG/Monographs/vol100F/index.php>.
- 24 Government of Ontario. Smoke-free Ontario 2014. <https://www.ontario.ca/page/smoke-free-ontario> (accessed 28 June 2017).
- 25 Maskarinec MP, Jenkins RA, Counts RW, et al. Determination of exposure to environmental tobacco smoke in restaurant and tavern workers in one US city. *J Expo Anal Environ Epidemiol* 2000;10:36–49.
- 26 O'Reilly KM, McLaughlin AM, Beckett WS, et al. Asbestos-related lung disease. *Am Fam Physician* 2007;75:683–8.
- 27 The Workplace Safety and Insurance Board. FAQ - Mandatory coverage in construction 2017. http://www.wsib.on.ca/WSIBPortal/faces/WSIBDetailPage?cGUID=WSIB013886&rDef=WSIB_RD_ARTICLE&_afLoop=1363692685201000&_afWindowMode=0&_afWindowId=null#%40%3FcGUID%3DWSIB013886%26_afrWindowId%3Dnull%26_afrLoop%3D1363692685201000%26rDef%3DWSIB_RD_ARTICLE%26_afrWindowMode%3D0%26_afrCtrl-state%3D4xyuj5uu_29 (accessed 16 Oct 2017).
- 28 Guha N, Loomis D, Guyton KZ, et al. Carcinogenicity of welding, molybdenum trioxide, and indium tin oxide. *Lancet Oncol* 2017;18:581–2.
- 29 Moulin JJ, Clavel T, Roy D, et al. Risk of lung cancer in workers producing stainless steel and metallic alloys. *Int Arch Occup Environ Health* 2000;73:171–80.
- 30 Cole DC. Cancer in the auto industry: A map of the literature for the Industrial Disease Standards Panel 1994. <http://www.canoshweb.org/odp/html/rp2.htm> (accessed 19 Jul 2017).
- 31 CAREX Canada. Lung carcinogens: occupational exposure summary package 2014. <http://www.occupationalcancer.ca/wp-content/uploads/2014/07/CAREX-Canada-Lung-carcinogens.pdf> (accessed 25 July 2017).
- 32 Centers for Disease Control and Prevention. *Criteria for a recommended standard: Occupational exposure to hexavalent chromium* Atlanta: CDC, 2013.
- 33 Alavanja MC, Sandler DP, Lynch CF, et al. Cancer incidence in the agricultural health study. *Scand J Work Environ Health* 2005;31(Suppl 1):39–45. discussion 5–7.
- 34 Blair A, Zahm SH, Pearce NE, et al. Clues to cancer etiology from studies of farmers. *Scand J Work Environ Health* 1992;18:209–15.
- 35 Ekberg-Aronsson M, Nilsson PM, Nilsson JA, et al. Socio-economic status and lung cancer risk including histologic subtyping—a longitudinal study. *Lung Cancer* 2006;51:21–9.
- 36 Kennedy SM, Koehoorn M. Exposure assessment in epidemiology: does gender matter? *Am J Ind Med* 2003;44:576–83.
- 37 Gaudette LA, Richardson A, Huang S. Which workers smoke? *Health Rep* 1998;10:35–45.
- 38 Axelson O. Aspects of confounding and effect modification in the assessment of occupational cancer risk. *J Toxicol Environ Health* 1980;6:1127–31.
- 39 Blair A, Steenland K, Shy C, et al. Control of smoking in occupational epidemiologic studies: methods and needs. *Am J Ind Med* 1988;13:1–4.
- 40 Association of Workers' Compensation Boards of Canada. Canadian workers' compensation system - Year at a glance 2017. http://awcbc.org/?page_id=11803 (accessed 16 Oct 2017).