2.71; above TLV 1.95, 95% CI: 1.27 to 3.00). About one third of CTS cases were attributable to exposure levels above the AL.

Conclusions The ACGIH TLV method predicted the risk of CTS, but the dose-response was flat above the AL: a fine-tuning of the proposed thresholds should be considered.

1631b RISK FACTORS FOR CARPAL TUNNEL SYNDROME – FINDINGS FROM THE NIOSH UPPER EXTREMITY MUSCULOSKELETAL DISORDER CONSORTIUM

BA Evanoff*, E Eisen, F Gerr, S Burt, K Hegmann, B Silverstein, A Garg, AM Dale, S Bao, C Harris-Adamson, J Kapellusch, L Merlino, D Rempel. *Division of General Medical Science, Washington University School of Medicine, Saint Louis, MO, USA*

10.1136/oemed-2018-ICOHabstracts.738

Starting in 2000, six research groups from the USA were supported by NIOSH to perform large, prospective epidemiologic studies examining associations between workplace physical risk factors and upper limb musculoskeletal disorders. A total of 4321 workers at 55 employers/plants across a variety of hand-intensive industries were followed for up to six years. Individual workplace exposure data included direct observation and video analysis. Health data included self-report, physical examination, and nerve conduction measures; our case definition for carpal tunnel syndrome (CTS) required both typical symptoms and nerve conduction abnormalities. Those performing the physical examinations and the video analyses were blinded to exposure and medical condition, respectively. Pooled analyses of consortium data controlled for personal factors (age, body mass index, gender, co-morbid diseases) and non-overlapping physical exposures (force, posture repetition) to study the association between work exposures and carpal tunnel syndrome. We found no independent effects of wrist posture or total repetition rate on the incidence of CTS. In contrast, strong dose-dependent associations were found between incident CTS and peak hand force (Borg CR10 ≥ 3), *forceful* repetition rate (>3 exertions per minute of ≥ 9 N pinch force or 45N power grip), and the proportion of time spent in forceful exertion ($>11\%$). We also found that the ACGIH Threshold Limit Value for Hand Activity (TLV for HAL) predicted CTS, and that that current 'action limit' is too high to adequately protect workers. Varying the formula of the TLV to emphasise force over repetition better predicted incident CTS. Study findings suggest that efforts to reduce workplace exposures should focus on jobs requiring high hand force and repeated or prolonged forceful exertions. Our study also suggests that the TLV for HAL and other less labour intensive assessment methods are valid and usable tools for workplace prevention.

1631c RECENT CHANGES TO THE ACGIH HAND ACTIVITY LEVEL TLV

D Rempel*. *Division of Occupational and Environmental Medicine, University of California, San Francisco, USA*

10.1136/oemed-2018-ICOHabstracts.739

The ACGIH Hand Activity Level (HAL) Threshold Limit Value (TLV) is a risk assessment tool designed to protect

workers, who perform repetitive hand exertions for 4 or more hours per day, from distal upper extremity disorders. Recent large, longitudinal studies, provide strong evidence that repetitive forceful hand exertions increase risk for occupational wrist tendinosis and carpal tunnel syndrome.^{1,2,3} In the Harris-Adamson study,³ forceful hand exertions (e.g., >9 N pinch or >45 N power grip force) and the %time performing forceful hand work increased risk. Based on these and psychophysics studies,⁴ hand exertions should be considered in risk assessment models if they are above 10% of posture specific strength. In addition, in these large studies, the prior HAL TLV action limit (0.56) was not sufficiently protective⁵ and, therefore, has been revised. The name of this TLV was changed to Hand Activity (HA) TLV. Other changes to the TLV will also be presented.

REFERENCES

1. Harris C, Eisen E, Goldberg R, et al. Workplace and individual factors in wrist tendinosis among blue-collar workers: The San Francisco study. *Scand J Work Environ Health* 2011;**37**(2):86–98.
2. Bonfiglioli R, Mattioli S, Armstrong T, et al. Validation of the ACGIH TLV for hand activity in the OCTOPUS cohort: A two-year longitudinal study of carpal tunnel syndrome. *Scand J Work Environ Health* 2013;**39**:155–63.
3. Harris-Adamson C, Eisen EA, Kapellusch J, et al. Biomechanical risk factors for carpal tunnel syndrome: A pooled study of 2474 workers. *Occup & Environ Medicine* 2015;**72**:33–41.
4. Potvin JR. Predicting maximum acceptable efforts for repetitive tasks an equation based on duty cycle. *Human Factors* 2012;**54**(2):175–188.
5. Kapellusch JM, Gerr F, Malloy EJ, et al. Exposure-response relationships for the ACGIH TLV for hand activity level: Results from a pooled data study of carpal tunnel syndrome. *Scand J Work Environ Health* 2014;**40**:610–20.

1631d QUANTIFYING COMPLEX TASKS USING THE REVISED STRAIN INDEX

JM Kapellusch*, A Garg. *University of Wisconsin – Milwaukee, Milwaukee, USA*

10.1136/oemed-2018-ICOHabstracts.740

Modern occupational tasks are often complex – consisting of multiple, discrete sub-tasks, each with its own unique combination of force, duration of exertion, posture, and frequency. Quantifying the physical stress from these complex tasks using simple, 'mono-task' assessment tools, such as the 1995 Strain Index (SI), can be difficult and assumes a large degree of ergonomics and MSD knowledge and training. Compared to the 1995 SI, the Revised Strain Index (RSI) minimises complicated measurement decisions and improves upon the 1995 SI by: (1) using percent maximum voluntary contraction (or Borg CR-10 equivalent) for applied hand force, (2) using duration per exertion (in seconds) rather than duty cycle, and (3) distinguishing between flexed and extended wrist postures. Thus the stress from each effort of a task (i.e., each sub task) can be individually quantified by the RSI and compared to other efforts in a cycle, or alternative efforts in the case of task intervention. By incorporating frequency of exertions (i.e., efforts per minute) and duration of task (i.e., hours per day), the RSI summarises stress associated with a simple task in a manner similar to the 1995 SI. For complex tasks, the RSI incorporates the Composite Strain Index (COSI) algorithm which integrates individual sub-tasks into an aggregate COSI score. The algorithm rank-orders subtasks from most to least stressful, thus ensuring that total task-stress is not lower than that of the highest stress sub-task alone. Sub-tasks are then incrementally added to one another by integrating over the frequency of exertion multiplier to produce the final COSI