Introduction The organophosphates (OPs) are a group of insecticides that have been widely used worldwide for the past 50 years. It is estimated that 3,000,000 people are exposed to OPs yearly, with the associated death rate being 3,00,000 deaths/year. OPs are cholinergic inhibitors and their toxicity is possible through acute or chronic exposure, with severe consequences for different organs and systems. The main objective of this study was to identify signs and symptoms of exposure to OPs on women with chronic exposure.

Methods A prospective study (1994–2014) followed up 43 women exposed to OPs, and evaluated signs and symptoms described as being associated to OPs exposure, as well as analytical parameters associated to asthenia and cellular protection, namely pyruvate (PA) and lactate (LA).

Results During the first appointment, 98% of the women had asthenia. Among the evaluated signs and symptoms, musculoskeletal injuries (78%) and menstrual cycle changes (36%) occurred with the highest frequency. Slightly less frequent were the changes observed in the peripheral nervous system (11%). Regarding the evaluated biochemical parameters, women generally showed a decrease in ferritin and an increase in TSH. Throughout the study a progressive drop in AChE was observed. In 64% and 92% of the women a change in PA and LA values respectively was observed.

Discussion The results suggest that prolonged exposure to OPs can chronically affect different human organism systems, namely parameters related to mitochondrial dysfunction

855

DERMAL EXPOSURE TO DIISOCYANATES: DEVELOPMENT AND VALIDATION OF AN ANALYTICAL METHOD FOR ACCURATELY ASSESSMENT OF VERY LOW LEVELS OF EXPOSURE

¹Radu-Corneliu Duca*, ¹Matteo Creta, ¹Katrien Poels, ^{1,2}Lode Godderis, ¹Jeroen Vanoirbeek. ¹*Environment and Health, Department of Public Health and Primary Care, University of Leuven, Leuven, Belgium;* ²*External Service for Prevention and Protection at Work, IDEWE, Haasrode, Belgium*

10.1136/oemed-2018-ICOHabstracts.1185

Introduction Dermal exposure to sensitizers such as diisocyanate have been described to promote the development of asthma in later stages when respiratory occupational exposure occurs. Therefore, we developed a reliable, sensitive and validated methodology based on dermal patches to assess skin exposure to diisocyanates.

Methods An UPLC-Unispray-MS/MS method was established and validated in order to reach very low levels of detection. Custom-made dermal patches were developed in order to allow optimal sampling of diisocyanates. Their sampling capability was evaluated in a controlled environment test-chamber where patches were exposed to increasing concentrations of diisocyanates.

Result The UPLC-MS/MS method using a Unispray ionisation source, based on supercritical fluids ionisation and Coanda effect, allowed reaching very low levels of detection (LoD=1 pg/mL) for all the targeted compounds (i.e. 4,4-MDI, 2,4-MDI, 2,6-TDI, 2,4-TDI, 1,6-HDI, and IPDI). Due to the high sensitivity of the analytical method, very low levels of diisocyanates (i.e. 25 pg/patch) are detected on the custom-made dermal patches. Furthermore, the patches allowed the sampling of a broad range of concentration levels (from 5 pg/cm² to 5 ng/cm²), which have been correlated with the air levels from the controlled environment chamber-test.

Discussion We succeeded to develop a method to assess dermal exposure to diisocyanates. Field studies are now necessary to further evaluate the suitability of the custom-made patches, as well as to relate low levels of exposure and potential health outcomes.

Primary Care and Work

1701

SCALING UP WORKERS' HEALTH COVERAGE THROUGH PRIMARY HEALTH CARE

¹Garth Manning, ²Frank van Dijk, ²Peter Buijs. ¹*Chief Executive Officer, World Organisation of Family Doctors (WONCA); Bangkok, Thailand;* ²*Learning and Developing Occupational Health (LDOH) foundation, The Netherlands*

10.1136/oemed-2018-ICOHabstracts.1186

Aim To explore ways of upskilling primary health care to offer essential occupational health to a broader population; some country examples.

Presenters: ¹Prof Jukka Takala, ²Dr Ramnik Parekh, ³Dr Orrapan Untimanon, ⁴Dr Hanifa M Denny, ⁵Dr Ezequiel Lopez, ⁶Dr Muzimkhulu Zungu, ⁷Dr Ahmet Özlüh, ⁸Dr Paul Smits

¹President of International Commission on Occupational Health (ICOH); Senior Consultant, Ministry of Manpower Services Centre, Singapore

²Chair- Mission BOHS: Indian Association of Occupational Health (IAOH), Mumbai, India

³Bureau of Occupational and Environmental Diseases, Department of Disease Control, Ministry of Public Health, Thailand

⁴Diponegoro University, Semarang, Indonesia

⁵Clinica Belgrano Family Medicine and Occupational Health Department, Quilmes, Argentina

⁶National Institute for Occupational Health School of Health Systems and Public Health, University of Pretoria

⁷Republic of Turkey Ministry of Health, General Directorate of Public Health, Ankara, Turkey

⁸Centre for evidence-based education, Academic Medical Centre (AMC), Amsterdam, The Netherlands

1701a

GLOBAL BACKGROUND OF INSUFFICIENT COVERAGE BY OCCUPATIONAL HEALTH SERVICES: WHAT CAN WE DO TO SUPPORT PREVENTION AND CARE

Jukka Takala. Senior Consultant, Ministry of Manpower Services Centre, Singapore

10.1136/oemed-2018-ICOHabstracts.1187

Background According to latest estimates by ILO 2.8 million men and women die every year caused by problems at work. Health disorders cause 2.4 million deaths. Costs are 3.9% of the global GDP or 3 trillion (million millions) USD. Problems include: poor legal and enforcement coverage, poor or no workers' compensation systems, poor or no occupational health services – some 15% or less of the global workforce is covered. According to WHO 93% of global health resources go for treatment and 7% for prevention. Lack of knowledge, policies, systems and resources is evident.

Methods and processes A number of solutions and good practices have been identified to be useful and successful, although largely in developed countries. Simple methods to increase coverages and enhance services are needed. The ILO