effectiveness of treatments. The use of ionising radiation in radiology, however, is not without risks for health professionals directly involved in radiation work.

Methods The study was cross-sectional and prospective, including medical imaging technicians and engineers selected radiology departments, regularly hired and assigned to radiation work. Data were reported on a self-administered questionnaire.

Results Five hospitals were selected with 59 participants, including 6 (9%) women, 54 (92%) senior technicians and 5 (8%) engineers. The cumulative age group of 30–50 years constituted 95% of the participants; 16 (27%) participants had been exposed for 5–10 years, 2 (4%) had been exposed for more than 20 years; The risk of cancer (96.6%), radiodermatitis (31%) and infertility (71.4%) were recognised by the participants. The main PPEs identified by the participants as radio-protectors were the lead apron (96.6%), the leaded glove (68.6%), the shells (31%), the anti-RX goggles (57.8%); 33.2% of the participants wore them regularly, 60.6% were irregular, 6.2% did not wear them. PPE was available for 37.8% of the presentations. The interest of the dosimeter was known to 94.4% of the participants. Apparatus was revised in 19% of cases; 91% of participants received IR training and were qualified to work under radiation. Pictograms existed in 40% of hospitals, light signals in 80%; 58.2% of the presentations knew their meaning.

Discussion The low availability and irregular wearing of PPE, and the ignorance of hazard indicators are more likely to expose them to IR.

Conclusion Strengthen protection measures through the availability of PPE and training

931 OCCUPATIONAL SKIN CANCER IN OUTDOOR WORKERS IN ITALY: EXPECTED NUMBER VS CASES RECOGNISED BY THE ITALIAN NATIONAL COMPENSATION AUTHORITY (INAIL)

¹Alberto Modenese*, ²Leena Korpinen, ¹Fabriziomaria Gobba. ¹Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy; ²Clinical Physiology and Neurophysiology Unit, North Karelia Central Hospital and Honkalampi, Joensuu, Finland

10.1136/oemed-2018-ICOHabstracts.1229

Introduction Solar Ultraviolet Radiation (UV) is one of the main risk factors for Non Melanoma Skin Cancer (NMSC) and Malignant Melanoma (MM). In Italy, only considering agriculture, fishery and construction sectors, the approximate number of workers exposed to solar UV (Outdoor workers – OW-) is 2 million (1.6 million males, 4 00 000 females).

Our aim is to compare the number of skin cancers (SCs) expected in OW to the number recognised by the Italian National Compensation Authority (INAIL).

Methods We collected data of Italian National Cancer Registries and the INAIL database of occupational diseases, including cancer, respectively available on the websites www.registri-tumori.it and www.bancadaticsa.inail.it

Results In Italy the yearly incidence of MM is 14,2 per 1 00 000 in males and 13 per 1 00 000 in females, that of NMSC 119.4 per 1 00 000 in males and 90.7 per 1 00 000 in females.

Applying these incidence rates to the Italian OW number, the expected SCs per year are approximately 2561 (279 MMs and 2282 NMSCs). INAIL recognised as occupational disease n. 246 cases (20% MM vs 80% NMSCs) in the last 5 years, i.e. less than 50 cases per year.

Discussion and conclusions Our results show that, in Italy, the National Compensation Authority recognises less than the 2% of the cases expected to occur in OW each year: 50 vs 2500. Main limitations of these data are that the incidence rates applied to OWs were not standardised, that the number of solar UV related SCs calculated is possibly under-estimated, considering that, e.g., not all OW groups were included, and that the data from cancer registries were quite outdated while the SCs incidence is increasing.

In conclusion, our data suggest a large under-estimation of occupational SCs in Italy, and that a better recognition of these diseases in OW is a relevant, and urgent, problem.

934 GLIOMAS INCIDENCE IN ITALY

¹Alberto Modenese*, ²Leena Korpinen, ¹Fabriziomaria Gobba. ¹Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy; ²Clinical Physiology and Neurophysiology Unit, North Karelia Central Hospital and Honkalampi, Joensuu, Finland

10.1136/oemed-2018-ICOHabstracts.1230

Introduction The International Agency for Research on Cancer has classified Radio Frequency in the Group 2B, ‘possibly carcinogenic to humans’, based on an association found between exposure related to mobile phones use and risk of gliomas and acoustic neuromas, even if this classification was discussed for some inconsistencies. Considering the large increase in mobile phone users from the ‘80 s, an increase in incidence of these tumours should be expected. The aim of this work is to follow the incidence of gliomas in Italy.

Methods The national incidence of central nervous system (CNS) cancers and that of gliomas from 2006 was obtained from the Italian Cancer Registries

Result Considering CNS cancers, in the period 1996–2009 the yearly age-standardised incidence per 1 00 000 has decreased from 10.8 to 9.8 in males, and from 8.0 to 7.0 in females. The incidence of gliomas, available for years 2006–2009 only, shows a slight increase from 5.7 to 6.3 in males, while is substantially stable (around 4) in females.

Discussion and conclusion Since the late ‘90 s, mobile phone use in Italy has largely increased: according to World Bank data, rose from 11.3 mobile phones per 100 Italians in 1996, to 107.7 in 2004 (exceeding the number of inhabitants), and 142.1 in 2015. If RF exposure related to mobile phones is associated with gliomas, considering a possible latency of 10–20 years an increase of the incidence of this tumours should be expected, at least from the second half of the 2000s

Considering all CNS tumours, during the period 1996–2009 data show a decreasing trend in Italy. For glioma, currently available standardised incidence rates, covering only the period 2006–2009, suggest a slight increase, but limited to males. Data available at this moment do not adequately support any firm conclusion on the trends of these tumours in Italy.

Reproductive Hazards

1727 REPRODUCTIVE HAZARD IN THE WORKPLACE AND ENVIRONMENT

Pau-Chung Chen*. *Department of Public Health and Institute of Occupational Medicine and Industrial Hygiene, National Taiwan University College of Public Health, Taipei, Taiwan*

10.1136/oemed-2018-ICOHabstracts.1231

Aim of special session

1. Provide new knowledge to prevent reproductive hazards.
2. Build mutual networks among safety and health experts

¹Soile Jungewelter, ²Imane EL Amri, ^{3,4}Andrea Hiddinga-Schipper, ⁵Ayman Fahim, ⁶Teus Brand, ⁷Ching-Chun Lin

¹Finnish Institute of Occupational Health, Helsinki, Finland

²Faculty of Medicine and Pharmacy of Casablanca, Casablanca, Morocco

³National Institute for Insurance against Accidents at work-Department of Occupational and Environmental Medicine, Epidemiology and Hygiene, Monteporzio Catone (RM), Italy

⁴Arbo Unie, Utrecht, The Netherlands

⁵Faculty Of Medicine, Suez Canal University, Ismailia, Egypt

⁶Netherlands Centre for Occupational Diseases, Coronel Institute of Occupational Health, Academic Medical Centre, Amsterdam, The Netherlands

⁷Institute of Occupational Medicine and Industrial Hygiene, National Taiwan University College of Public Health, Taipei, Taiwan

1727a IS OCCUPATIONAL NOISE EXPOSURE DURING PREGNANCY RELATED TO LANGUAGE ACQUISITION OF THE CHILD?

^{1,2}Soile Jungewelter*, ^{1,2}Helena Taskinen, ³Eira Jansson-Verkasalo, ^{1,4}Marja-Liisa Lindbohm, ¹Jouko Remes, ³Minna Huottilainen, ¹Markku Sallmén. ¹Finnish Institute of Occupational Health, Helsinki, Finland; ²University of Helsinki, Helsinki, Finland; ³University of Turku, Department of Psychology and Speech, Turku, Finland; ⁴University of Tampere, Tampere, Finland; ⁵University of Helsinki, Cognitive Brain Research Unit, Helsinki, Finland

10.1136/oemed-2018-ICOHabstracts.1232

Introduction We examined the impact of noise exposure during pregnancy on the child's language acquisition at the age of one and two years.

Methods We conducted a cohort study among women working in the food industry, as kindergarten teachers, musicians, dental nurses and pharmacists with a new born child. At the age of one year the analyses included 408 mother-child pairs and at the age of two years 288. The mother filled the baseline questionnaire before the child was 12 months old, and the language acquisition questionnaire when the child was 12 months (Infant-Toddler Checklist, ITC) and 24 months (MacArthur Communicative Development Inventories, MCDI) old. An occupational hygienist assessed the noise exposure in three categories: no exposure, low exposure, moderate/high exposure. We analysed the data using linear regression.

Results At the age of one there were no statistically significant differences among the adjusted mean language acquisition ITC-scores of the children in different exposure categories. The adjusted scores among boys were 30.1 (95% CI: 28.3 to 31.8) for no exposure, 29.7 (27.4–32.0) for low noise, and 29.3 (26.7–31.9) for moderate/high exposure. Among girls these were 33.7 (31.9–35.5), 33.8 (31.3–36.4), and 33.6 (31.3–36.0), respectively. No associations were found in analyses of social communication, speech production and language comprehension. Noise exposure was associated with lower scores among kindergarten teachers. At the age of two mean MCDI-scores did not differ significantly between the noise exposure groups. The adjusted mean scores for expressive vocabulary among girls were 295 (95% CI: 254 to 336) for no exposure, 303 (243–362) for low exposure, and 269 (212–326) for moderate/high exposure. Among boys the scores were 200 (154–246), 178 (111–246), and 225 (153–298), respectively.

Discussion We found no clear association between noise exposure during pregnancy and language acquisition among one-year- or two-year-old children.

1727b PREMATURE BIRTH AND WORKING CONDITIONS: A PROSPECTIVE STUDY ABOUT 68 PREGNANT WOMEN

¹I El Amri*, ²F Benbrahim, ¹B Benali, ¹A El Kholti. ¹Faculty of Medicine and Pharmacy, Casablanca, Morocco; ²Faculty of Medicine and pharmacy, Rabat, Morocco

10.1136/oemed-2018-ICOHabstracts.1233

Introduction The question about mediation between work and pregnancy is a real problem in industrialised countries, but in a country like ours, the issue is far away from being taken as a priority. Thus, we've chosen to conduct this prospective study about premature birth and working conditions trying to find a correlation between the two, while removing medical and obstetrical risk factors.

Methods The study was conducted in between January and December 2014. We've included women who had a job 6 months or more before the conception. To evaluate working conditions we've used the fatigue score developed by Mamelle. To evaluate the impact of working conditions on premature birth, we've monitored cervical length changes.

Results At first, we've gathered a selection of 68 pregnant women debuting their follow up between 11 and 13 weeks of amenorrhea plus 5 days. An acceleration of the modification of the uterine cervix was noted within the group of women having a «Mamelle» score of 3.

Discussion The first epidemiological study bringing up the part of tiring working conditions in premature birth was conducted by Mamelle in France. After that, more authors, such as Mac Donald, got interested by the subject and implicated long working hours, rotating shifts and carrying heavy loads in the occurrence of premature birth. We took 18 mm as a limiting value for the length of the cervix under which intervention measures should be taken.

Conclusion Risk assessment of working conditions is the only deciding factor when it comes to the pursuit of work while being pregnant. The decision needs to be taken by both the attending and occupational physicians.