1737a OCCUPATION AND COPD

David Fishwick. *University of Sheffield, UK*

10.1136/oemed-2018-ICOHabstracts.1302

Occupational exposures are able to cause a variety of occupational lung diseases that effect either predominantly the airway or the lung parenchyma. Exposures to agents at work that are able to cause COPD remain globally common, although the evidence supporting each of these is variable. Better evidence is available to support the roles of silica, grain dust, coal and cadmium, for example, as specific causes of occupational COPD, whilst multiple epidemiologic al studies also support the harmful effect of exposures to vapours, gases dusts and fumes.

This presentation will summarise the current view of occupational COPD, to include its definition, common causes and it will also include discussion of the uncertainties around occupational COPD. The talk will also include specific mention of the harmful effects of inhaled respiratory crystalline silica, and the continuing global problems associated with silicosis.

1737b OCCUPATION AND ASTHMA

JR Feary. *Royal Brompton Hospital, London, UK*

10.1136/oemed-2018-ICOHabstracts.1303

Asthma is widely recognised to be a heterogenous disease with several 'endotypes'; differentiation of these is not necessarily straightforward and requires assessment from several different angles. Moreover there is increasing evidence that the label is (over-) applied inappropriately.

The often close relationships between occupation and asthma are equally well recognised. Employees, for example, may develop a work-related exacerbation of their pre-existing disease or, *de novo*, 'occupational asthma', usually as a result of an acquired hypersensitivity to a, airborne, workplace sensitising agent; differentiation between these two conditions is often, but not always, straightforward.

However, not infrequently, respiratory symptoms are erroneously attributed to asthma when they are in fact, a reflection of other conditions originating in the upper airway or in relation to disordered breathing patterns. Correct identification of these conditions allows individuals to remain in their current job and avoids the use of protracted polypharmacy; novel approaches are available to assist in making these diagnoses and help to direct appropriate management.

In addition to assessment of current symptoms, a key question often posed is how an individual's asthma will affect their future employment? There is good evidence that variation *between* individuals is greater than that *within* an individual; this allows the prediction of risk of future exacerbations and of the risks of recurrence of remitted childhood disease.

This presentation will review modern approaches to diagnosis and management of asthma and related conditions, including the appropriate role of novel, 'biologic, treatments.

572 WORK-RELATED ASTHMA ASSOCIATED WITH HIGH PREVALENCE OF RESPIRATORY SYMPTOMS AND REDUCED WORK-ABILITY IN A CROSS-SECTIONAL POPULATION BASED STUDY FROM NORWAY

^{1,2}G Klepaker*, ¹MV Svendsen, ^{2,3}J Kongerud, ¹AKM Fell. ¹Department of Occupational and Environmental Medicine, Telemark Hospital, Skien, Norway; ²Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway; ³Department of Respiratory Medicine, Oslo University Hospital, Oslo, Norway

10.1136/oemed-2018-ICOHabstracts.1304

Introduction Some studies have indicated that patients with occupational asthma have more severe disease than other asthma patient. Indicators of severity may be reduced quality of life and higher levels of sick-leave and disability. There are, however, few studies assessing respiratory symptom prevalence and work-ability. The aim of our study was to estimate the differences in respiratory symptom prevalence and work-ability between patients with indication of work-related asthma compared with other asthma patients.

Methods The association between work-relation and respiratory symptoms and work ability was assed among 16 099 subjects from the general population, aged 16 to 50 years, included in a cross-sectional study form Telemark, Norway. 828 subjects that reported having physician-diagnosed asthma, ever being employed and currently using asthma medication were included. Of these 171 (21%) had respiratory symptoms at work that improved during weekends, holydays, other absence from work or change of work. These were compared with the remaining patients with physician-diagnosed asthma (n=657). Self-reported work-ability was assessed by the first single item question of the work ability index (WAI). Analyses where adjusted for age, gender, education and occupational group.

Result The patients with work-related asthma reported significantly more respiratory symptoms in the past 12 months; wheezing (OR 2.6, 95% CI: 1.5 to 4.5), woken by dyspnoea during the night (OR 2.0, 95% CI: 1.4 to 2.8), woken by coughing (OR 1.8, 95% CI: 1.3 to 2.6), asthma attack (OR 2.2, 95% CI: 1.5 to 3.1) and reduced work-ability (OR 2.3, 95% CI: 1.5 to 3.3).

Discussion Our study indicates that work-related asthma is associated with higher prevalence of respiratory symptoms and reduced work-ability (score <8) compared with asthma without work-relation. These findings support the hypothesis suggesting that occupational asthma may be more severe than non-occupational asthma.

573 OCCUPATIONAL RESPIRATORY DISORDERS IN A 'CHEMICAL' INDUSTRY IN TOGO

¹Antoine Vikkey Hinson*, ²Gbehomilo Edhorh, ³Koffi Atsu Aziagbe, ⁴Badiro Aguemou, ⁵Fabien Gounongbe, ¹Menonli Adjibimey, ¹Benjamin Fayomi. ¹Unity of Teaching and Research in Occupational Health and Environment, Faculty of Sciences and Health of Cotonou (University of Abomey-Calavi Benin); ²Service des Urgences médicales du Centre Hospitalier Universitaire Sylvanus Olympio de Lomé-Togo; ³Service de Pneumo-ptisiologie du Centre Hospitalier Universitaire Sylvanus Olympio de Lomé-Togo; ⁴Département de Santé Publique, unité de biostatistique Faculté des Sciences de la Santé de Cotonou (Université d'Abomey-Calavi Bénin); ⁵Département de Médecine et Spécialités Médicales, Faculté de Médecine, Université de Parakou Bénin

10.1136/oemed-2018-ICOHabstracts.1305

Introduction Occupational asthma (OA), obstructive chronic bronchopneumonia (OCBP) and pneumoconiosis are occupational respiratory diseases having important impact on morbidity and mortality. Some population studies show that 15% of the cases of asthma with the adults can be attributed to work environment. The aims of this study is to evaluate the respiratory function especially the occupational asthma of workers in a plastic bag manufacturing factory in Togo.

Material and methods By a cross sectional study, we included 182 workers in the sample. They filled a questionnaire and the workers with asthma complains was submitted to spirometry test.

Results The mean age of subjects was $37,13 \pm 5,71$ years with a male predominance (99,5%). The mean of the work experience was $7,66 \pm 2,88$ years. 1,1% of workers were smokers. Rhinitis and conjunctivitis were reported with 79,7%. The frequency of occupational rhinitis was 62,6%. Subjects that presented asthma complains were 34,1%. In 25,8% of cases asthma complains are associated to the work. The abnormal pulmonary function was overlooked by reversible smalls airways pattern (6,6%) and reversible obstructive pattern (4,9%). Spirometry was compatible with asthma in 11,5% of cases. The prevalence of occupational asthma was 7,7%. Occupational asthma only concerns the posts of Thermo-welding and Extrusion. All the workers in these posts did not use respiratory protection.

Conclusion Occupational asthma is a reality in a plastic bag manufacturing factory in Togo. This fact needs preventives and correctives actions to avoid that disease.

612

A FREELY-AVAILABLE TUBERCULOSIS SCREENING TOOL FOR ADULT WORKERS EXPOSED TO RESPIRABLE CRYSTALLINE SILICA

¹NK Reul*, ²B Braid, ³Z Gray. ¹Associate Medical Director for Occupational Disease, Washington State Department of Labour and Industries, Olympia, Washington, USA; ²Clinical Nurse Specialist, Washington State Department of Labour and Industries, Olympia, Washington, USA; ³Epidemiologist, Washington State Department of Labour and Industries, Olympia, Washington, USA

10.1136/oemed-2018-ICOHabstracts.1306

Introduction Exposure to respirable crystalline silica is known to increase the risk of progression to active tuberculosis (TB) disease in those who have latent TB infection. Both latent TB infection and active TB disease are relevant components of screening examinations in silica-exposed workers.

Methods Health care providers screening silica-exposed workers for TB must generally:

- identify for further evaluation and treatment any employees with suspected active TB disease
- adhere to safety and health regulatory requirements, when applicable
- even when regulations do not require screening, exercise sound screening practices in workers who stand to benefit from medical evaluations
- be familiar with current peer-reviewed publications; specialty society recommendations; national and state public health policy approaches
- coordinate with local approaches to TB screening

- incorporate both occupational and non-occupational health indications into screening decisions.

Using a publicly-funded, transparent, and democratic process, the Department of Labour and Industries has developed a TB risk assessment and testing algorithm for adult workers exposed to respirable crystalline silica.

The screening tool integrates these varied needs and resources into a cohesive document for use at the point of health care delivery.

Result The tool is a clinical adjunct that helps health care providers make two key decisions when evaluating patients with exposure to respirable crystalline silica:

1. Which workers should undergo a comprehensive evaluation for active tuberculosis disease?
2. Which workers should receive testing for latent TB infection?

Discussion Widespread availability of the risk assessment and testing algorithm is assured through direct inclusion in Washington State regulatory documents. The tool integrates medical content and public policy for clinicians performing the TB screening in silica-exposed workers through a single, accessible form. As a freely-available clinical aid, it is available to clinicians performing TB screening in silica-exposed workers throughout the United States, and worldwide.

690

BREATH TEST FOR PNEUMOCONIOSIS USING AN ELECTRONIC NOSE: A CASE-CONTROL STUDY

¹Hsin-Yi Peng*, ¹Che-Jui Chang, ^{1,2,3,4}Pau-Chung Chen, ^{1,2,3}Hsiao-Yu Yang. ¹Institute of Occupational Medicine and Industrial Hygiene, National Taiwan University College of Public Health, Taipei, Taiwan; ²Department of Public Health, National Taiwan University College of Public Health, Taipei, Taiwan; ³Department of Environmental and Occupational Medicine, National Taiwan University Hospital, Taipei, Taiwan; ⁴Department of Environmental and Occupational Medicine, National Taiwan University College of Medicine, Taipei, Taiwan

10.1136/oemed-2018-ICOHabstracts.1307

Introduction Current medical surveillance program has flaw that may result in poor detection of early pneumoconiosis around the world. Breath analyses have attracted substantial attention in the screening of occupational environmental lung disease. Pneumoconiosis could generate specific volatile organic compounds (VOCs) that may constitute a specific breath print for diagnosis. The objective of this study was to develop a breath test for pneumoconiosis by analysing VOCs using sensor array technique.

Methods We conducted a case-control study and enrolled study subjects from stone workers in Hualien between October 2016 and November 2016. One litter of breath air was collected after five minutes of tidal breathing through a non-rebreathing valve with inspiratory VOC-filter, and storage by a Tedlar bag. The air was analysed by a 32 nanocomposite sensor array electronic nose within 30 min. Using the ILO/ICRP profusion category $\geq 1/1$ in chest X-ray as the reference standard, we assessed the diagnostic accuracy of the breath test. Data were randomly split into 80% for model building and 20% for validation.

Result After excluding three subjects with poorly controlled diabetes mellitus with fasting serum glucose level >200 mg/dl and one subjects of asthma under medication, a total of 98 subjected were used in final analysis that included 34 cases of pneumoconiosis and 64 healthy controls. By linear