fingers. Such a suggestion may apply to transmission to even more far district, due to the fact that blood vessels can work as waveguide for pressure, because they are designed to be so. **Methodology** Can be summarised as following. Effects of vibrations on the hearing function will be assessed by stimulated otoacoustic emission method. Vibration elicitation will be strictly controlled (shaker and 6 DOF vibrating plate). Overall vibration will be measured by accelerometers on joints and head. Blood vessel vibration propagation will be measured by high resolution, dual frequency echography^{4,5} on main vessels (coronary) and small vessels. The frequency spectrum of hearing loss and blood vessels vibrations, deducted by the heart pumping effects, will be compared to look for coincidence.

REFERENCES

1. Sisto R, Botti T, Cerini L, Di Giovanni R, Marchetti E, Lunghi A, Sacco F, Sanjust F, Tirabasso A, Moleti A. Synergistic effects of noise and hand-arm vibration on distortion product otoacoustic emissions in healthy subjects. *Int. J. Ind. Ergon* 2016;1–7. <http://dx.doi.org/10.1016/j.ergon.2016.10.006>
2. Adewusi SA, Rakheja S, Marcotte P, Boutin J. Vibration transmissibility characteristics of the human hand–arm system under different postures, hand forces and excitation levels. *Journal of Sound and Vibration* 2010;**329**:2953–2971. <http://dx.doi.org/10.1016/j.jsv.2010.02.001>
3. IFA Report 5/2017e, DGVV, Hand-arm vibration: Exposure to isolated and repeated shock vibrations – Review of the International Expert Workshop 2015 in Beijing. <http://www.dguv.de/ifa/publikationen/reports-download/reports-2017/index-2.jsp>
4. Papadopoulou V, Balestra C, Theunissen S, Germonpré P, Obeid G, Boutros A, Dayton PA, Eckersley RJ, Cosgrove D, Tang MX. *Can contrast mode echocardiography help estimate bubble population dynamics post-dive?* EUBS 43rd annual scientific meeting, Ravenna, Italy, 12–16 September 2017.
5. *Self-assessment of the jugular venous pulse from space and special environments* – Paolo Zamboni, EUBS 43rd annual scientific meeting, Ravenna, Italy, 12–16 September 2017.

510 THE IMPORTANCE OF NOISE IN THE REQUEST OF EARLY RETIREMENT IN BRAZIL

RV Brito*, VR Batalini, AES Zafalon, LR Ferreira. *Centro Universitário das Faculdades Associadas de Ensino – UNIFAE, São João da Boa Vista, Brazil*

10.1136/oemed-2018-ICOHabstracts.1481

Introduction In Brazil, workers exposed to harmful agents are benefited by a public policy denominated Early (Special) Retirement. The main reason for this early retirement is to benefit workers who performed activity that can cause harm to health and include agents such as noise, biological, chemistry, heat, and others. This exposure has to be continuous, permanent and non-intermittent. In this context, due to the large number of companies that expose their workers to noise, the present study aims to analyse the prevalence of noise's requests granted in the request of Early Retirement and characterise the profile of workers who are exposed to this harmful agent.

Methods This was a retrospective study based on the analyses of 118 Profissiographic Profile Documents (PPD), a Brazilian National Social Security document use for granting early retirement for the worker exposed to harmful agents. The research was performed from November 2016 until May 2017, characterising the noise as the main harmful agent for the concession of the early retirement.

Results The noise accounted for 50,79% of requests for Early Retirement, in addition to 74,19% of all benefits granted. Considering merely the noise's requests, 44,74% were accepted. Considering the analysis of gender, 97,67% of noise's requests were made by men. Furthermore, the more

prevalent employment in the analysis were production assistant (26,07%), production operator (21,79%) and mechanic (13,62%). In addition, considering the noise's requests granted, production assistant represented 36,52%, followed by production operator (27,83%) and mechanic (4,35%).

Discussion The study's findings appointed the need for a special attention of government and companies to look for new strategies that aim to reduce the noise intensity in the work environment, especially on the secondary sector (Manufacturing and Industry), contributing for a better occupational health and worker production

1433 EFFECTIVENESS OF 'HEARING CONSERVATION PROGRAM' IN THE LNG INDUSTRY – A QATAR PERSPECTIVE

Nandakumar Gopakumar Pillai*, Rhee Costan. *Ras Gas Company Ltd, Qatar*

10.1136/oemed-2018-ICOHabstracts.1482

Background Noise, or unwanted sound, is one of the most pervasive occupational health problems. Ras Gas hearing conservation program aims to prevent hearing deterioration of its employees working in Ras Gas Company Ltd (RG) owned or operated properties that have potential exposure to high levels of noise and is based on the hearing conservation guidelines from OSHA.

Methods This Prospective study analysed the accumulated hearing evaluation data of the employees working with Ras Gas for a minimum of 10 years in the 'Similar exposure group' (SEG) within the operations group to test the effectiveness of the hearing conservation program (HCP) which was implemented in 2009. The data was extracted from the Medical Director (MD) Software of the Medical Services Department and includes the original audiograms of the 70 selected employees belonging to the operations group. This study planned to analyse:

- Pre – employment audiograms
- Audiograms taken at the start of the hearing conservation program (2009) and
- The audiograms done in 2014 (i.e; the hearing assessment of these employees from the day they started working with Ras Gas to the time the Hearing Conservation program was started and up to the 2014 hearing status) of the selected employees.

Results A total of 210 audiograms (3 × 70) were reviewed. The analysis of the audiograms displayed an improvement in employees hearing in line with the introduction of a hearing conservation program when the age correction factor was applied.

Discussion This study showed that, if properly executed, a hearing conservation program can prevent, reduce and even improve noise induced hearing deterioration. The study is particularly important in industries like mining, quarrying and oil and gas extraction which has the highest prevalence estimates of hearing loss.

1260 RELEVANCE OF AUDIOMETRIC HSE CATEGORIZATION REFERENCE VALUES AND MODERNIZATION

Sujil Jacob*, Martin Hogan. *Corporate Health Ireland Ltd, Cork, Ireland*

10.1136/oemed-2018-ICOHabstracts.1483

Introduction Audiometry plays a vital role in both a preventative and screening aspect in Occupational Medicine. It serves to quickly triage and identify those individuals at risk for progressive hearing loss in noisy environments and adherence to requirements in specific professions. To facilitate appropriate referral pathways, a system of categorisation HSE/HSA has been developed ranging from 1 (acceptable hearing) to 3 (Poor hearing) based on comparisons for age and gender – this allows for trained professionals who are not specialist *per se* to be involved in testing.

This paper examines series of audiograms performed for the purpose of pre-employment to establish whether the expected distribution of 80th, 20th, 5th percentile of the normal population respectively for category 1 to 3, is in keeping with the actual values of the tested populations and, if at all there is a ‘Healthy worker effect’ which refers to the working population having a favourable morbidity profile.

Methods Sequential Analysis of 1000 Pre-employment Audiograms and their computer generated HSE Categorization

Result The results are compared to the population sample used for standardisation of the HSE categorization.

Discussion The question arises of whether there is a need to update current age- gender adjusted reference values of the categorization tables, taking in to account the increased sound exposure to both environmental and Personal devices resultant of Modernization.

718 TREMOR MEASUREMENTS IN A TWENTY-TWO YEAR LONGITUDINAL STUDY OF WORKERS EXPOSED TO HAND-HELD VIBRATING TOOLS

¹Rita Bast-Pettersen, ¹Karl-Christian Nordby, ²Gunilla Wastensson, ¹Raymond Olsen, ¹Lisa Aarhus. ¹Department of Occupational Medicine and Epidemiology, National Institute of Occupational Health, Oslo, Norway; ²Department of Occupational and Environmental Medicine, Sahlgrenska University Hospital and University of Gothenburg, Sweden

10.1136/oemed-2018-ICOHabstracts.1484

Background Use of hand-held vibrating tools may lead to hand-arm vibration syndrome (HAVS), which is composed of vascular, neurological and muscular components. Few studies have investigated possible long-term effects on tremor related to hand-held vibrating tools.

Objectives The objectives were to evaluate postural and rest hand tremor in workers with current and previous exposure to vibrating hand tools, taking into account the possible effects from medical conditions and life-style habits such as tobacco and alcohol consumption.

Subjects and methods Forty workers previously employed in a specialised engineering and construction company were examined with a test-battery and a clinical examination in 1994 and 2017. Age at last examination was 60.7 years (44.6 to 77.8 years).

Hand tremor was measured in 2016/2017 with the Tremor Pen from CATSYS.

Their exposure to hand-held vibrating tools was assessed as acceleration x lifetime exposure. Biological samples of Carbohydrate-deficient transferrin (CDT), glycated haemoglobin (HbA1c), cotinine, nicotine and caffeine were collected on the day of examination.

Results The data collection was finished by ultimo March 2017. Preliminary results indicate that tobacco consumption assessed by

cotinine in serum and diabetes assessed by HbA1c were more closely associated with tremor parameters than lifetime exposure to hand-held vibrating tools. For all subjects the magnitude of the tremor, the tremor Intensity, was 0.15 and 0.13 ms⁻² for the dominant and non-dominant hands respectively.

The hand position is important when testing for tremor. When postural and rest tremor measures were compared, the frequency of the tremor increased significantly, from 7.4 to 9.8 Hz and from 7.6 to 9.3 Hz for the dominant and non-dominant hands respectively.

424 WHOLE BODY VIBRATION AMONG PROFESSIONAL BUS DRIVERS – EVALUATION OF AN INTERVENTION STUDY TO REDUCE LOW BACK PAIN

¹K Lauwers*, ¹S Acke, ^{1,2}M Verbrugghe, ¹G Dumoulin, ¹M-N Schmickler, ²L Braeckman. ¹Mensura Occupational Health Services, Belgium; ²Department of Public Health, University of Ghent, Belgium

10.1136/oemed-2018-ICOHabstracts.1485

Introduction Bus drivers are exposed to whole body vibration (WBV) and postural stress, both risk factors for low back pain (LBP).

The study aimed to (1) assess WBV exposure and (2) evaluate the effect of an intervention to reduce LBP.

Methods

- Exposure to WBV was assessed by measuring vibration levels (i.e., weighted root mean square acceleration [Aw] and vibration dose value [VDV]) in nine buses.
- For the intervention study, a toolbox presentation including ergonomic and lifestyle advice and an exercise program was instructed interactively to bus drivers. Two questionnaires (pre- and post-intervention) were constructed based on the VIBRISKS questionnaire to evaluate the effect of the intervention on LBP prevalence and to assess associated risk factors (e.g., manual material handling, seating posture, and WBV). The bus drivers' motivation to change driving behaviour was also evaluated.

Results

- The vibration levels never exceeded the exposure limits [Aw: 1.15 m/s², VDV: 21 m/s^{1.75}]. Only in one bus, the Aw (0.56 m/s²), were the levels slightly higher than the action limit [0.5 m/s²]. Four buses had a VDV (9.56 m/s^{1.75}, 13.0 m/s^{1.75}, 10.7 m/s^{1.75} and 11.0 m/s^{1.75}) that exceeded the action limit [9.1 m/s^{1.75}].
- Sixty and 47 bus drivers participated in the pre- and post-intervention, respectively. The prevalence of LBP in the previous seven days decreased significantly (p=0.02). The score on correct lifting techniques was a significant determinant (p=0.046) in predicting LBP in the previous seven days (OR=0.32, 95% CI: 0.10 to 0.98). There was a significant post-intervention increase in knowledge about correct seating posture (p=0.001), and 95.5% of the bus drivers were willing to maintain their behavioural change.

Discussion WBV exposure should be frequently evaluated to maintain below action-limit levels. Although this study revealed a significant reduction in LBP prevalence in the previous seven days, further research is needed in a randomised controlled trial design.