the Asbestos and Kgalagadi Relief Trusts that compensate qualifying miners and environmental claimants.

Results Women comprised 2.47% (n=394) of 15 940 cases. The women were older (56.6 ± 17.11 years) than men (53.63 ± 14.44 years) and had shorter mining-related exposures (7.21 ± 7.71 versus 18.18 ± 18.20 years). Most women had asbestos mining (46.19%) or environmental (14.72%) exposure; 87 (22.08%) were gold and 37 were platinum (9.39%) miners. Among men, there were 64.28% gold, 18.47% platinum and 5.55% asbestos miners, and 0.30% had environmental asbestos exposure. Disease proportions in women and men were: emphysema, 16.00% and 27.73%; silicosis, 3.30% and 23.13%; tuberculosis, 17.77% and 23.13%; lung cancer, 4.31% and 3.67%; asbestosis, 16.75% and 4.28% and mesothelioma, 17.26% and 2.00%.

Discussion The burden of asbestos-related diseases was high in women. The few cases of silicosis in women are an alert to high silica dust exposures. The study highlights the importance of autopsies for disease diagnosis, and education of women on mining and environmental OLD-related risks and their right to access statutory compensation.

1452 POWERFUL WOMEN'S HANDS

¹I Bello*, ²A Salmen-Navarro. ¹Simon Bolívar University, Venezuelan Society on Occupational Health, Caracas, Venezuela; ²New York University, New York, USA

10.1136/oemed-2018-ICOHabstracts.1498

Introduction In 2016 a research on musculoskeletal disorders and gender was completed, especially on the capacity of women to perform repetitive tasks involving a prehensile effort of less than 15 n or manual loads of less than 50 g per repetition, in relation to male workers.

Methods The sample was formed by 150 women and 150 men from three industrial plants located in Valencia (Venezuela), Santa Cruz (Venezuela) and Tijuana (Mexico) in fan, snacks and hydraulic connexions manufacturing sectors. The study correlated three variables: job risks (biomechanics: repetition and postures, psychosocial: quantitative psychological requirements), health valuation and average labour productivity.

Results Women had a lower biomechanical involvement in upper limbs in the presence of a similar exposure to male, having, on average, a higher rate of productivity, especially in the case of tasks of low force demand. The male group always started the days (daily and weekly) with peaks of productivity, but this was decreasing throughout the day and week; However, the productivity of the female group remained constant and was, on average, 18.3% higher than that of the male sample, taking into account the effect of absences, medical rest and turnover.

Conclusion The study does not show a greater biomechanical resistance of the female articular systems in relation to the masculine ones, but it does suggest a greater 'muscular intelligence', which acts as a protective factor and enhances productivity. From a macroergonomic point of view, the female population was more resilient, which allows maximising the learning curve, increasing labour availability and minimising turnover. These protective factors can be incorporated as part of a plan for the prevention of the ergonomic and psychosocial risks from which both genders can benefit.

1519 NIGHT WORK OF WOMEN ABOUT 50 CASES

W Allouche, M Lgharbi, B Benali, K Chati, A El Kholti. *Casablanca Faculty of medicine and pharmacy Hassan II University, Morocco*

10.1136/oemed-2018-ICOHabstracts.1499

Introduction The work in atypical and cyclical schedules is becoming more and more current with the authorisation of night work of women and the continuous work of many companies. Regular and prolonged night work can have important repercussions, also on the health of employees and on their living conditions.

Methods This is a cross-sectional descriptive study that aims to analyse the constraints, conditions and effects of night work on women using a questionnaire distributed on 50 production operators all working in shift at night (23 hour 00–07 hour 00) with bearing system 3×8 in an automotive cabling company in Kéniterra Morocco.

Results Of the 50 operators questioned the average age is 37 years, Having 6 years of average professional seniority. 62% start work between 00:00 and 05:00 in the morning and only 24% finish their work before 03:00 in the morning. 92% of the operators report difficulty falling asleep, 82% have a disturbed sleep while 88% say that the rhythm of life impedes their social life. 90% of women have digestive disorders, 60% of which eat at irregular hours.

Discussion Many studies show that women would have more sleep deficit than men due to home bonds these results are identical to our study, since there is a high incidence of sleep disorders this can be explained that all the interviewees are female. According to Knutsson A, gastrointestinal disorders such as abdominal pain and problems of diarrhoea and constipation are more common in irregular workers than in day workers, these are similar with our study, since 90% of women present digestive disorders.

Conclusion In the light of new scientific contributions, a legal framework for night work in general is particularly necessary for women to improve working conditions and to prevent their harmful effects.

1754 INTEGRATED HEALTH-BASED RISK ASSESSMENT FRAMEWORK FOR SINGAPORE'S WORKPLACES

¹Jeff Hwang Yi-Fu, ²Cheng Yue Pan, ³Associate Professor Chia Sin Eng, ⁴Judy SNG. ¹Lecturer, Saw Swee Hock School of Public Health, National University of Singapore; ²Senior Manager (Total Workplace Safety and Health), Workplace Safety and Health Council (Singapore), Tripartite Alliance Limited; ³Senior Consultant (Total Workplace Safety and Health), Workplace Safety and Health Council (Singapore), Tripartite Alliance Limited and Associate Professor, Saw Swee Hock School of Public Health, National University of Singapore; ⁴Senior Lecturer, Saw Swee Hock School of Public Health, National University of Singapore

10.1136/oemed-2018-ICOHabstracts.1500

Introduction Despite significant global improvement in occupational health and safety over the past few decades, workplace deaths and injuries are still occurring. At the same time, many countries are experiencing an increasing prevalence of chronic diseases due to ageing population and lifestyle factors. Currently, risk assessment does not commonly take into account employees' personal health risks. Our paper aims to review the association between chronic diseases and risk of workplace injuries, and recommend a health-based risk assessment