from 2002 to 2016 were analysed with joinpoint regression and the magnitudes of occupational class differences were estimated with the relative index of inequality (RII).

Results Most of the trends first increased and turned to decrease in 2007/2010. Managers and professionals had the least amount of self-certified SA, but steadily increasing trends were observed among men. Self-certified SA followed only partially the typical socioeconomic gradient, as routine non-manuals had the highest levels of SA. The magnitude of occupational class differences in self-certified SA was stable during the study period only among women. Self-certified SA and occupational class differences have increased in recent years among men in the lower occupational classes.

Conclusion Socioeconomic differences exist in self-certified SA among young employees, but gradient is only partial. Overall, high amounts of self-certified SA especially in the lower occupational classes are in need for further studies and preventive measures.

131 BEATING BURNOUT, BEING KIND – OAK

¹Zakiah Amir*, ²Anna McHugh, ³Lynda Sisson. ¹Faculty of Occupational Medicine, Royal College of Physicians of Ireland, Dublin, Ireland; ²Letterkenny University Hospital, Letterkenny, Ireland; ³Workplace Health and Wellbeing Unit, Health Service Executive, Dublin, Ireland

10.1136/oemed-2018-ICOHabstracts.1722

Introduction There is increasing evidence of burnout among healthcare workers (HCW). Workers engagement is one solution to burnout with positive outcomes towards patient care and work satisfaction. Simple, low key positive activities have been shown to make differences to people in a pressurised environment. Our intervention aims to establish whether acts of kindness at work can affect workers engagement.

Methods Occasional Acts of Kindness (OAK) was a pilot intervention programme trialled in a large tertiary hospital in December 2016. All HCWs were invited to attend the 2 hours session when participants take a break from work in a friendly atmosphere, sit for a while, engage with one another and ask colleagues about how they were coping with work and life. The effectiveness of this intervention was evaluated using a validated staff engagement questionnaire, Utrecht Work Engagement Scale. Participants were also encouraged to leave their own comments regarding the intervention.

Result 65 HCWs of varying grades and departments participated. 97% of participants agree that the intervention boosts collegiality and strengthens relationships at work, 90% agree it would make them more likely to take the initiative to help a struggling colleague, 80% agree it boosts their energy at work, 83% agree it increases their mental resilience at work, 90% agree it enhances their enthusiasm at work, 87% agree it helps them persevere even if things are not going well, 63% agree it helps them continue working even for long period, 84% agree it helps instil a sense of pride in workplace and 93% agree it boosts overall morale at work. Their additional comments regarding the intervention were also cohesively positive.

Discussion Our intervention proved that acts of kindness at work have multitudes positive effects on workers and work. A culture of kindness is one solution to the growing issue of burnout among HCW.

478

DOCPASS: A TOOL TO ASSESS READINESS FOR WORK IN SURGEONS AND ANAESTHETISTS TO IMPROVE AND ENSURE SAFETY OF PATIENTS

BM de Niet*, MHW Frings-Dresen, JK Sluiter. Academic Medical Centre, Coronel Institute of Occupational Health, Amsterdam Public Health research institute, Amsterdam, The Netherlands

10.1136/oemed-2018-ICOHabstracts.1723

Introduction To protect patient safety in the operating room it is important to create awareness among surgeons and anaesthetists on their own work readiness and its influence on their performance. The purpose of the study was to design a tool (DOCpass), study its feasibility in daily operating room practice and redesign it based on suggestions made by future end users.

Methods The content of the DOCpass prototype was selected on the basis of aspects of health and work that can influence performance as was found in literature. A suitable platform was subsequently chosen to meet pre-defined requirements such as ease of use and accessibility of the tool. To determine the feasibility in the operating room practice and collect points of improvement for redesign, post-task walkthroughs were organised with surgeons and anaesthetists at a university medical centre in the Netherlands.

Result DOCpass was developed as a webpage in JavaScript and HTML which makes the tool accessible from local hospital computers or mobile devices (e.g. mobile phone, tablet). It screens for personal aspects (e.g. fatigue and mood states) and work-related aspects (e.g. shift and sleeping time duration) that have been found to affect performance and patient safety in practice. A total of 13 improvement points were suggested by future end users and the prototype was adjusted according to these. The improvements concerned the content of the introduction, the relevance of screening methods and applicability of feedback.

Conclusion DOCpass assesses and provides feedback on work readiness in colour coding through digital screening of work and personal health aspects. Participating surgeons and anaesthetists in the feasibility study reported that the tool is potentially useful in the operating room were the content to be improved. Feedback that is provided should be easy to act upon and could in this way add practical value to work.

853

DEVELOPMENT OF AN INTERVENTION FOR NURSES' SUSTAINABLE EMPLOYABILITY: A FOCUS GROUP STUDY

¹LAM Brouwers*, ²JA Engels, ²YF Heerkens, ³AJ Van der Beek. ¹PHD student, HAN University of Applied Sciences, Nijmegen, The Netherlands; ²Professor of HAN University of Applied Sciences, Nijmegen, The Netherlands; ³Professor, Head of Department of Public and Occupational Health, VU University Medical Centre, Amsterdam, The Netherlands

10.1136/oemed-2018-ICOHabstracts.1724

Introduction In today's Western world, people have to work longer before they can retire and the working population is ageing. In healthcare, due to an increase in complexity, severity and extent of the demand for health care, it is of great importance to invest in retention of nurses, who are considered particularly vulnerable in staying employable.

A structured dialogue between employee and supervisor is seen as promising opportunity to improve sustainable employability of nurses and the corresponding required working behaviour. In order to accomplish so-called 'joint ownership',

composing an intervention programme will be planned in joint effort by a constructive dialogue between employee and supervisor. The aim of this qualitative study was assess the needs of employees and supervisors towards the aforementioned approach to improve sustainable employability of nurses.

Method A structured needs assessment was performed to answer the question of what are the realistic, attainable and useful characteristics of an intervention, in which nurses and their supervisors have a constructive dialogue on sustainable employability. The so-called Vitality Scan is filled out beforehand and used as a starting point of the dialogue, during which customised follow-up actions are arranged if appropriate. We performed six focusgroup interviews, three amongst employees (n=13) and three amongst supervisors (n=15). Qualitative data analysis was performed using Atlas.ti.

Results Both employees and supervisors perceived a safe *psychosocial working environment* as most essential. We found a positive *attitude* and *intention* towards the characteristics of the Vitality Scan and dialogue. The (time) effort and amount of *self-efficacy* amongst supervisors were considered potential barriers as well as facilitators. Thorough training of the supervisors was considered necessary.

Conclusion Nurses and supervisors are positive towards working on sustainable employability of nursing staff using the designed intervention. Several existing facilitators and barriers in the organisation were addressed and practical improvements were suggested.

750

PSYCHOSOCIAL RISKS IN TRADE UNION REPRESENTATIVES

¹S Ronsmans*, ^{2,3}L Godderis. ¹KU Leuven – University of Leuven, student Master of Occupational Medicine, Leuven, Belgium; ²KU Leuven – University of Leuven, Department of Public Health and Primary Care, Environment and Health, Leuven, Belgium; ³IDEWE, External Service for Prevention and Protection at Work, Heverlee, Belgium

10.1136/oemed-2018-ICOHabstracts.1725

Introduction Quantitative studies reveal that trade union representatives are dealing with a number of *specific* psychosocial risk factors. The existing literature leaves many questions unanswered on the underlying reasons for these psychosocial risks and on difficulties in addressing them. A qualitative research could offer an opportunity to tackle this knowledge gap and to provide deeper understanding of this subject.

Methods Eight semi-structured in-depth interviews were conducted with trade union representatives from the 2 largest Belgian unions: ACV (*Confederation of Christian Trade Unions*) and ABVV (*General Federation of Belgian Labour*). A purposeful sample of participants was recruited from a wide range of sectors and types of work organisation. Template analysis was used to examine their unique job demands and job resources.

Results Using the job demands-resources model as a conceptual framework it was possible to map the unique psychosocial risk factors experienced by trade union representatives. Quantitative, qualitative and emotional overload, role conflict, role ambiguity and a conflictual relationship with their employers were found to be the major job demands. Social support from the workers, other representatives and the trade union were found to be the most important job resources, as well as having the knowledge and skills, experiencing success and having

strong trade union beliefs. The findings of this study also suggest that workplace restructuring could profoundly impact the job demands and job resources experienced by trade union representatives.

Conclusions A rich and complex account of the experiences of the participants could be created. In general the results fit the existing literature but they add a deeper understanding of *how* and *why* trade union representatives are experiencing certain job demands and job resources. The study resulted in a wide range of propositions to improve the prevention of psychosocial risks in trade union representatives.

694

EFFICIENCY OF VOCATIONAL REHABILITATION PROGRAMS FOR WORKERS WITH SCHIZOPHRENIA: A SYSTEMATIC LITERATURE REVIEW ACCORDING TO THE PRISMA GUIDELINES

¹A-E Leverich, ¹S Acke*, ^{1,2}M Verbrugghe, ¹M-N Schmickler, ³C De Brouwer. ¹Mensura Occupational Health Services, Belgium; ²Department of Public Health, University of Ghent, Belgium; ³Environmental health and Occupational health, Université libre de Bruxelles, Belgium

10.1136/oemed-2018-ICOHabstracts.1726

Introduction Unemployment is common among people with schizophrenia and contributes most to the economic cost of the condition. This study was aimed at (1) reviewing the available tools to facilitate the vocational rehabilitation of workers with schizophrenia and (2) describing their specific opportunities.

Methods A search of PubMed for original, peer-reviewed articles published from 2006 until January 31, 2016, yielded 643 citations. Exclusion criteria were patients less than 18 years or older than 65 years of age, patients with additional diagnoses, studies conducted in developing countries, lack of an adequate study design, systematic reviews or case studies, no work related studies, and other major analytic inadequacies. The remaining 24 articles were assessed in qualitative synthesis using the Scottish Intercollegiate Guidelines Network (score $\geq 2+$), Critical Appraisal Skills Program or Prediction Model Risk of Bias Assessment Tool checklists.

Results

- Only seven quantitative and one qualitative study could be included. The number of study participants with schizophrenia varied substantially (n=7 in the qualitative study, n=100 in the largest quantitative study). Most studies were conducted in the United States (n=5), as well as two in Canada and one in Japan.
- Cognitive remediation programs (Neurocognitive Enhancement Therapy) combined with supported employment were significantly more effective ($p < 0.005$) than supported employment by itself. Next, Individual Placement and Support (the manual version of supported employment) and cognitive behavioural therapy (Indianapolis Vocational Intervention Program) were, by themselves, more effective than conventional vocational rehabilitation, regardless of whether it was for competitive employment.

Discussion Combining cognitive remediation programs with supported employment programs significantly improved the vocational outcomes of workers with schizophrenia, especially