developed, to investigate the associations between exposure and the presence of skin damage.

Method Based on ICNIRP publications, a detailed questionnaire, considering both working and leisure UV exposure, was developed and applied in group of volunteers and in patients affected by Non Melanoma Skin Cancers (NMSCs), to evaluate whole-life SR exposure and the relevant factors associated, including exposure habits, environmental factors and use of protections.

Results In patients with NMSCs, we found that the co-presence of in situ squamous cell carcinoma (SCC) and invasive NMSCs was more frequent in OWs than in indoor workers (IW), as was also the frequency of multiple skin lesions. The prevalence of lesions in the most exposed body areas, as the face, was significantly higher in OWs, while working in shades was negatively associated with skin lesions. Also working posture was relevant: adopting a bent-over position was associated with lesions on the head. For leisure activities, the questionnaire application confirms the relevant role of tanning beds, in particular for body areas usually not highly exposed to SR, and the relevance of intense exposure in the middle hours of the vacation days ('sunbaths'), significantly associated with NMSCs presence.

Conclusion A detailed questionnaire-based evaluation in subjects with NMSCs can be very useful for estimating the whole-life individual UV exposure and the relevancy of occupation. Furthermore, an integration with personal dosimetric UV measurements can lead to a semi-quantitative evaluation, to set out exposure levels and different exposure modalities (e.g. intermittent vs cumulative) for various occupations and, possibly, for the different SCs.

1616c

INTEGRATED CARE FOR WORKERS WITH OCCUPATIONAL SKIN CANCER IN THE GERMAN SOCIAL ACCIDENT INSURANCE SCHEME – REHABILITATION WITH ALL APPROPRIATE MEANS PROVIDED BY THE STATUTORY SOCIAL ACCIDENT INSURANCE IN GERMANY

Bettina Lindemann. *Institution for Statutory Accident Insurance and Prevention for Health and Welfare Services, Germany*

10.1136/oemed-2018-ICOHabstracts.248

Introduction Skin cancer caused by exposure to certain harmful substances has been a part of the list of occupational diseases in Germany for some time. In 2015 certain types of skin cancer (squamous cell carcinoma or multiple actinic keratosis of the skin caused by natural UV-irradiation) were added as a new occupational disease to the list of occupational diseases. These types of skin cancer are recognised as occupational disease if the insured person has been exposed to certain effective doses of sunlight at his or her workplace. There is a rising number of such cases reported to the statutory social accident insurance and its institutions which are the entities in charge of prevention, rehabilitation and compensation of occupational diseases in accordance with the German social security system. The institutions for statutory social accident insurance have developed concepts to handle the growing numbers of these cases.

Methods Based on research and the discussion with medical experts, standards for the procedure, for the medical

treatment, the rehabilitation of workers affected by occupational skin cancer and the prevention of occupational skin cancer have been established. Concepts for follow-up care have been developed and new methods of therapy have been admitted for the medical treatment of patients suffering from occupational skin cancer.

Results The statutory social accident insurance has established measures to provide effective methods of rehabilitation for the patients concerned in accordance with the basic principle governing this social security scheme in Germany – acting 'with all appropriate means'.

Discussion Apart from providing effective help for the workers affected a major aim for the institutions of statutory accident insurance is the prevention of occupational skin cancer caused by exposure to natural UV-irradiation. The institutions of statutory social accident insurance will continue their efforts to reduce the risk for workers of contracting skin cancer caused by exposure to natural UV-irradiation at the workplace.

1616d

ON THE UV EXPOSURE OF WORKERS IN EUROPE IN GENERAL AND GERMANY IN PARTICULAR: WHAT DO WE KNOW? RESULTS OF THE GENESIS-UV PROJECT

Marc Wittlich. *Institute for Occupational Safety and Health (IFA), Sankt Augustin, Germany*

10.1136/oemed-2018-ICOHabstracts.249

Introduction We are exposed to solar ultraviolet radiation (UVR) every day, during work and leisure time. To date, there is only little knowledge about the level of irradiance. Already in 1992, UVR has been graded a group 1 carcinogen by the International Agency for Research on Cancer (IARC). In an ageing society, along with an ideal of a brown skin, non-melanoma skin cancer (NMSC) incidence rates rapidly increase.

Methods For risk assessment and deduction of protective measures, real measured data serve best. Regarding solar UVR exposure, long-term personal dosimetry measurements is required. With GENESIS-UV, we designed a suitable system. With only little impairment, each test persons measures autonomously. Data are transferred via mobile service or internet to a data server once a week automatically. GENESIS-UV has proven to be feasible in collaborations in the whole world without any changes in the system.

Results Since 2014, about 800 test persons delivered data points representing about 80 000 days of measurement – in Germany. We derived the annual irradiance for more than 100 occupations, and far more occupational activities as well. Interestingly, the irradiance values for the occupations cover a very wide range (www.dguv.de/genesis). Going deeper into the occupational activities, prevention-relevant information could be obtained. Since 2016, measurements have been expanded at European level.

Discussion Outdoor workers have to be protected from UVR exposure. During working hours, their exposure can be up to 4.5 times the leisure dose. Our approach from occupations to occupational activities enables us to suggest very distinct protective measures, and drawing a full picture of workers exposure. Particularly, we showed that already short exposure times may lead to a notable hazard. Thus, the definition of an outdoor worker has to be reviewed.

The study with GENESIS-UV is by far the largest study on UV irradiance of workers worldwide.

Dermatoses

1605 APPLICATION OF BIOMARKERS IN WORK-RELATED CONTACT DERMATITIS

Sanja Kezic*. *Coronel Institute of Occupational Health, Academic Medical Centre, Amsterdam, Netherlands*

10.1136/oemed-2018-ICOHabstracts.250

Aim of special session Understand the role of biomarkers for early interventions in occupational health settings

Work-related contact dermatitis represent a substantive part of occupational diseases in many countries of the World. They cause extensive suffering for the affected workers and represent a huge socio-economic burden. Selection of appropriate biomarkers, including those involved in skin barrier and inflammation, can provide the basis for early, health-protective interventions in occupational health settings. Furthermore, they allow identify new potential therapeutic targets, aid diagnostics and monitoring of therapy and interventions to improve workers' skin health

1605a INDIVIDUAL SUSCEPTIBILITY TO WORK-RELATED CONTACT DERMATITIS

Sanja Kezic. *Coronel Institute of Occupational Health, Academic Medical Centre, Amsterdam, Netherlands*

10.1136/oemed-2018-ICOHabstracts.251

Introduction Contact dermatitis (CD) is the most common cause of occupational skin diseases, especially in occupations such as nursing, hairdressing, metalworking and construction industry. Environmental exposure and personal susceptibility both contribute to its development.

Methods Recent studies on phenotypic and genotypic factors which might modify risk for irritant- and allergic contact dermatitis (resp. ICD and ACD) will be addressed.

Results Polymorphisms in several candidate genes responsible for skin barrier, immune response and metabolism showed to play a role in individual susceptibility to CD. Next to genetic factors, a history of atopic dermatitis and filaggrin loss-of-function mutations have been confirmed as major risk modifying factors for ICD. Association between atopic dermatitis and filaggrin mutations with ACD is less clear which might at least partly be assigned to differences in study design e.g. choice and definition of phenotype, investigated contact allergens, and population size. E.g. the effect of filaggrin mutations seems to be allergen-specific. Several studies suggested polysensitization as a phenotype for increased susceptibility to ACD.

Discussion Advances in our knowledge on the mechanisms underlying CD initiated a large number of studies searching for candidate genes that might modify the risk for ICD and ACD. Future studies are warranted to investigate whether genetic and phenotypic factors shown to be related to individual susceptibility are specific for a certain skin irritant or contact allergen and further evaluate their suitability in clinical practice.

1605b FILAGGRIN DEGRADATION PRODUCTS AS A BIOMARKER OF IRRITANT- AND ALLERGIC CONTACT DERMATITIS

Ivone Jakasa. *Faculty of food technology and biotechnology, University of Zagreb, Zagreb, Croatia*

10.1136/oemed-2018-ICOHabstracts.252

Introduction Filaggrin, a structural protein of the uppermost skin layer, stratum corneum (SC) plays an important role in aetiology of contact dermatitis (CD). Filaggrin contributes to the mechanical strength of the skin while its degradation products are major constituent of natural moisturising factors (NMF) responsible for adequate skin hydration. The major determinant of NMF in the skin are mutations in the filaggrin gene, a known risk factor for occupational contact dermatitis. Next, NMF can be affected by dermal exposure to skin irritants or contact allergens. The levels of NMF might therefore serve as a biomarker of individual susceptibility or to assess skin damaging effect of irritants and allergens.

Methods In experimental studies in humans, the levels of NMF components in the SC were determined before and after exposure to common skin irritants (SLS; n-propanol, acetic acid and NaOH) and contact allergens (Cr, Ni, MCI/MI and PPD).

Results All skin irritants led to significant decrease in NMF levels in the SC while among contact allergens this effect was observed only by MCI/MI.

Discussion Decrease in NMF levels indicates damage of the skin barrier which occurred after dermal exposure to both, contact allergens and irritants. Therefore, NMF might be a useful biomarker to detect early effects associated with dermal exposure to chemicals in occupational settings. Furthermore, NMF might be useful to identify skin damaging properties of contact allergens.

1605c INDIVIDUAL SUSCEPTIBILITY FOR OCCUPATIONAL SKIN DISEASES IN METAL WORKERS

Richard Brans. *Department of Dermatology, Environmental Medicine and Health Theory, University of Osnabrueck, Germany*

10.1136/oemed-2018-ICOHabstracts.253

Introduction Occupational contact dermatitis (OCD) is one of the most frequent occupational diseases worldwide and very common among metal workers. The burden of disease is high due to its often chronic course and a high risk of work absenteeism and unemployment. The prerequisite for the development of OCD is work-related exposure to irritants and/or allergens. However, endogenous factors may increase the individual susceptibility and interfere with preventive interventions.

Methods A prospective controlled intervention study has been initiated in metal apprentices with annual follow-up assessments over three years. The apprentices in the intervention cohort received specific health education on prevention of OCD at the beginning of their training. Saliva and stratum corneum samples were taken for genotyping and analysis of phenotypic biomarkers, including assessment of epidermal