

# Night work and miscarriage: a Danish nationwide register-based cohort study

Luise Moelenberg Begtrup,<sup>1</sup> Ina Olmer Specht,<sup>2</sup> Paula Edeusa Cristina Hammer,<sup>1</sup> Esben Meulengracht Flachs,<sup>1</sup> Anne Helene Garde,<sup>3</sup> Johnni Hansen,<sup>4</sup> Åse Marie Hansen,<sup>5</sup> Henrik Albert Kolstad,<sup>6</sup> Ann Dyreborg Larsen,<sup>3</sup> Jens Peter Bonde<sup>1</sup>

► Additional material is published online only. To view please visit the journal online ([/](#)). Please see the author notes to BMJ

<sup>1</sup>Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark

<sup>2</sup>The Parker Institute, Bispebjerg and Frederiksberg Hospital, Frederiksberg, Denmark

<sup>3</sup>National Research Centre for the Working Environment, Copenhagen, Denmark

<sup>4</sup>Institute of Cancer Epidemiology, Danish Cancer Society, Copenhagen, Denmark

<sup>5</sup>Department of Public Health, University of Copenhagen, Copenhagen, Denmark

<sup>6</sup>Department of Occupational Medicine, Danish Ramazzini Centre, Aarhus University Hospital, Aarhus, Denmark

## Correspondence to

Dr Luise Moelenberg Begtrup, Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg Hospital, København, 2400, Denmark; [luise.moelenberg.begtrup.02@regionh.dk](mailto:luise.moelenberg.begtrup.02@regionh.dk)

The results were presented at the Reproyoung conference, Ystad 25th–26th October.

Received 14 November 2018  
Revised 5 February 2019  
Accepted 17 February 2019  
Published Online First  
25 March 2019



© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Begtrup LM, Specht IO, Hammer PEC, *et al.* *Occup Environ Med* 2019;**76**:302–308.

## ABSTRACT

**Objective** Observational studies indicate an association between working nights and miscarriage, but inaccurate exposure assessment precludes causal inference. Using payroll data with exact and prospective measurement of night work, the objective was to investigate whether working night shifts during pregnancy increases the risk of miscarriage.

**Methods** A cohort of 22 744 pregnant women was identified by linking the Danish Working Hour Database (DWHHD), which holds payroll data on all Danish public hospital employees, with Danish national registers on births and admissions to hospitals (miscarriage). The risk of miscarriage during pregnancy weeks 4–22 according to measures of night work was analysed using Cox regression with time-varying exposure adjusted for a fixed set of potential confounders.

**Results** In total 377 896 pregnancy weeks (average 19.7) were available for follow-up. Women who had two or more night shifts the previous week had an increased risk of miscarriage after pregnancy week 8 (HR 1.32 (95% CI 1.07 to 1.62)) compared with women, who did not work night shifts. The cumulated number of night shifts during pregnancy weeks 3–21 increased the risk of miscarriages in a dose-dependent pattern.

**Conclusions** The study corroborates earlier findings that night work during pregnancy may confer an increased risk of miscarriage and indicates a lowest observed threshold level of two night shifts per week.

## INTRODUCTION

In Europe around 14% of all women report working at night at least once a month.<sup>1</sup> Studies in humans have found lower levels of melatonin mediated by exposure to light-at-night and with no full catch-up during the day among night workers.<sup>2–3</sup> Furthermore, several consecutive night shifts may cause circadian disruption by phase shifting the suprachiasmatic nucleus (master clock) desynchronising with the sleep cycle and the peripheral oscillators throughout the body.<sup>4</sup> Melatonin is primarily synthesised in the pineal gland, but also in peripheral organs such as the placenta and ovaries. It is thought to be an important free radical scavenger and plays a role in preserving the optimal function of the placenta.<sup>5</sup> Furthermore, experimental studies have demonstrated the importance of tightly regulated circadian rhythms, in which melatonin also has a pivotal role, in the maintenance of successful pregnancies.<sup>6</sup> Supporting this is the finding of a

## Key messages

### What is already known about this subject?

- Experimental studies indicate that endogenous melatonin contributes to the maintenance of a successful pregnancy. Night work causes exposure to light at night and circadian disruption, which decreases the release of melatonin.
- Observational studies have indicated an association between working nights and miscarriage, but inaccurate exposure assessment precludes affirmative risk assessment.

### What are the new findings?

- This is the first study to investigate the association between night work and miscarriage using detailed and prospective measurement of exposure to night work.
- Our results indicate that women who work two or more night shifts per week may be at increased risk of miscarriage the following week. Furthermore, both the cumulated number of night shifts and consecutive number night shifts increased the risk of miscarriage in a dose-dependent pattern.

### How might this impact on policy or clinical practice in the foreseeable future?

- The findings increase the knowledge about exposure to night work and have relevance for working pregnant women as well as their employers, physicians and midwives. Moreover, the results could have implications for national occupational health regulations.

lower pregnancy success rate among mice exposed to shifting in the light/dark cycle compared with controls.<sup>7</sup> However, many biological processes of the circadian regulation of reproduction in humans are still unknown.<sup>8</sup>

Around one-third of all human embryos are lost, the majority within 6 weeks from the last menstrual period, most often unnoticed by the pregnant women and only some 10%–14% are recognised as clinical miscarriages.<sup>9</sup> More than half of miscarriages are due to chromosomal abnormalities, which could arise within the sperm, within the egg before a female is born or during the completion

of meiosis shortly before conception. Since only the latter mechanism could possibly be caused by the mother's occupational exposures, miscarriages related to maternal exposures is possibly more easily detected among non-chromosomal late miscarriages.<sup>10</sup>

Meta-analyses addressing the association between night work and miscarriage have reported a moderately increased risk of miscarriage in relation to fixed night work, whereas no or weak associations are reported for rotating shiftwork including night work.<sup>11 12</sup> However, studies are few and exposure assessment primarily based on self-reports and limited by the inability to adjust for important factors such as sick leave and number of working hours. Thus, there is a need for prospective studies with refined exposure assessments making it possible to explore the effect of the intensity of night work and the types of shift schedules used.

The aim of this study was to investigate whether women who worked at night during pregnancy had an increased risk of miscarriage. We investigated the risk of miscarriage after night work the previous week and among women who worked cumulated night shifts, consecutive nights shifts and had quick returns back to work after a night shift (defined as shift return in <11 hours).

## METHODS

### Design and study population

Our register-based cohort study includes all female employees working in the five Danish administrative regions, who became pregnant during the period from 1 January 2007 through to 31 December 2013. As the Danish administrative regions run all public hospitals in Denmark, our cohort consists primarily of hospital-based employees, such as nurses and physicians.<sup>13</sup> Using their civil registration number we identified women who had given birth from the Danish Medical Birth Register (DMBR),<sup>14</sup> and women who had been treated at a Danish hospital for miscarriage, molar or ectopic pregnancy or induced abortion from the Danish National Patient Register (DNPR).<sup>15</sup> The DNPR holds information on all hospital contacts including inpatient, outpatient and emergency contacts, but not on contacts to specialists outside hospitals.<sup>15</sup> Both the DNPR and DMBR provide almost complete information on gestational age (GA) and day at delivery or submission to hospital. In Denmark all women are offered an ultrasound scan around pregnancy weeks 11–14 to screen for Down syndrome. Ninety-five per cent of the Danish women have the scan. Thus, the GA of births are, in most cases, based on ultrasonography, whereas the GA of miscarriages before pregnancy weeks 11–14 are, most often, based on the last menstrual period. We estimated the date of conception of each pregnancy by subtracting the GA from the date of delivery or hospital submission for miscarriage. For three miscarriages and 271 births (1.35%) missing data on the GA were replaced by the median values (8.5 weeks for miscarriages and 40 weeks for births). A number of 21 (1%) miscarriages occurred at 4 weeks. Miscarriages with registered GA less than 4 weeks (n=6), or more than 22 weeks (n=6) were excluded. Only the first registered pregnancy from 1 January 2007 through to 31 December 2013 with at least 28 days of employment after the date of fertilisation was included (the index conception) (online figure 1).

### Exposure assessment

Data on working hours were obtained from the Danish working hour database (DWHD), which is a national database of administrative payroll data.<sup>13</sup> For every working day the DWHD provides

information on the start and end time (date:hours:minutes) of a shift.<sup>13</sup> A night shift was defined according to the 2009 IARC working group on shiftwork, as working at least 3 hours between midnight and 5:00.<sup>4</sup> The sum of night shifts was computed for each consecutive pregnancy week from week 3 through to week 21. For descriptive purposes exposed employees were defined as study participants with one or more night shifts during pregnancy weeks 3–21.

The risk of miscarriage among women who were exposed to night work was examined as a 'short term effect' by the number of night shifts completed the previous week. Moreover a 'cumulated effect' was examined in three ways by adding the number of night shifts, by adding the number of consecutive night shifts with spells of at least two, three, four, five, six or seven night shifts, and by adding the number of quick returns after a night shift (initiating a new shift <11 hours after a night shift). All cumulated effects were calculated from pregnancy week 3 until the week before outcome, censoring or pregnancy week 22, whichever came first.

### Outcome assessment

Data on hospital admissions due to miscarriages, molar or ectopic pregnancies, and induced abortions were retrieved from the DNPR using the ICD-10 codes DO00–DO07. Using the median of registered GA, the miscarriages were categorised in two groups, namely miscarriages in pregnancy weeks 4–8 and miscarriages in weeks 9–22. Because late clinical miscarriages are defined as pregnancies terminating after pregnancy week 12, the association between night work and miscarriages in weeks 13–22 was also explored. The pregnancies were followed from week 4 until miscarriage (the outcome), molar or ectopic pregnancy (censoring), induced abortion (censoring), discontinuance of employment, or pregnancy week 22, whichever came first.

### Covariates

Maternal date of birth was obtained from the DWHD, which enabled calculation of maternal age at the time of the index conception. The DMBR provided information on parity (completeness 97.7%), while this information was not available from the DNPR. However, by linking women admitted for miscarriage to the DMBR it was possible to retrieve data on parity for most of the women who had given birth before or after the time of the miscarriage (93.6%). For nulliparous women with miscarriage this information was missing (6.4%). Baseline smoking and body mass index (BMI) were retrieved from the DMBR and based on the first midwife contact. For the women with miscarriage as index, pregnancy smoking status and BMI reported in relation to the birth closest in time to the hospital admission for the miscarriage was selected (median difference 42.9 months). Information on job title was retrieved from Statistics Denmark (DST) using DISCO-88 and DISCO-08, the Danish version of the International Standard Classification of Occupations in the calendar years 2007–2009 (DISCO-88)<sup>16</sup> and 2010–2013 (DISCO-08),<sup>17</sup> respectively. Classification of socioeconomic status (SES) into high, medium and low was derived from DISCO codes based on Statistics Denmark's categorisation. Covariates were grouped according to the categories presented in table 1.

### Statistical methods

To determine the 'short-term effect' of night work, exposure data were used as both a continuous variable and as categorised into three groups: none, one, or two or more night shifts the

**Table 1** Characteristics of the study population according to exposure of night work (n=22 744)

Characteristics	Exposed ≥1 registered night shift during pregnancy week 3–21 (n=10 047)	Reference group No registered night shifts during pregnancy week 3–21 (n=12 697)
Outcome of pregnancy, n (%)		
Births	9 089 (90)	11 007 (87)
Miscarriages	740 (8)	1 149 (9)
Molar and ectopic pregnancies	44	96 (1)
Induced abortions	174 (2)	445 (3)
Time for miscarriage (pregnancy week)		
Gestational age, median (min, max)	9.0 (4.0, 21.0)	8.0 (4.0, 21.0)
Follow-up weeks at risk, median (Pct 25, 75)	22 (22,22)	22 (22,22)
Work during pregnancy weeks 3–21, median (Pct 25,75)		
Number of day shifts	40 (25, 52)	50 (11, 80)
Number of evening shifts	6 (0, 16)	0 (0, 2)
Number of night shifts	9 (4,16)	
Maternal age at conception		
Mean years (SD)	30.5 (3.9)	30.9 (4.4)
≤25 years, n (%)	512 (5)	1 028 (8)
26–30 years, n (%)	4 531 (45)	4 701 (37)
>30 years, n (%)	5 004 (50)	6 968 (55)
Parity, n (%)		
0	5 948 (59)	6 967 (55)
1	2 442 (24)	3 434 (27)
2+	1 411 (14)	1 963 (15)
Missing	246 (2)	333 (3)
Former miscarriage, yes n (%)	736 (7)	1 104 (9)
BMI before pregnancy Mean (SD)	23.7 (4.3)	23.9 (4.6)
Underweight (<18.5 kg/m <sup>2</sup> ), n (%)	743 (7)	977(8)
Normal weight (18.5–24.9 kg/m <sup>2</sup> ), n (%)	6 646 (66)	8 167 (64)
Overweight (25.0–29.9 kg/m <sup>2</sup> ), n (%)	1 818 (18)	2 382 (19)
Obese (30+kg/m <sup>2</sup> ), n (%)	840 (8)	1 171 (9)
Smoking during pregnancy, n (%)		
Non-smoker	9 252 (92)	11 579 (91)
Smoker	492 (5)	726 (6)
Missing	303 (3)	392 (3)
Socio-economic status (SES), n (%)		
Low	869 (9)	3 563 (28)
Medium	6 939 (69)	6 811 (53)
High	2 224 (22)	2 230 (18)
Missing	15	93 (1)
Most frequent occupation, n (%)*		
Nurse	6 242 (62)	3 405 (27)
Physician	1 732 (17)	955 (8)
Medical secretary	53	1 373 (11)
Physiotherapist/occupational therapist	29	1 175 (10)
Nurse assistant	510 (5)	727 (6)
Laboratory technician	233 (2)	642 (5)
Cleaning/kitchen worker	17	557 (4)
Pedagogue/care helper	230 (2)	383 (3)
Psychologist	<10	418 (3)
Midwife	305 (3)	41
Office worker	10	304 (2)
Teacher/scientist	81 (1)	300 (2)

\*Among occupations with at least 100 employees.

previous week. Data on cumulated night shifts were also used both as a continuous variable and as categorised by 0, 1–10, 11–20, 21–25 and >25 night shifts.

We estimated the risk of miscarriage by the different night work dimensions by discrete Cox regression with time-varying exposure from pregnancy week 4 through to week 22,

corresponding to the time after the implantation of the fertilised egg and until the week after which expulsion of the fetus is defined as a preterm birth or stillbirth. Each week was assigned week-specific exposure levels, and analyses were performed with and without adjustment for maternal age, BMI, smoking, parity, SES and former miscarriages, which were chosen a priori.<sup>18 19</sup>

To ensure only night work prior to a miscarriage was taken into account, a lag of 1 week was used.

Competing risk by induced abortions<sup>20</sup> was examined in sensitivity analyses using the proportional hazard model proposed by Fine and Gray.<sup>21</sup> To account for possible differences between employees working and not working nights we performed sensitivity analyses within employees who had at least one night shift in pregnancy weeks 3–21. We observed a substantial decline in the number of registered miscarriages after 2010 (from 9.7% to 6%) and conducted a sensitivity analysis only including pregnancies registered between 2007 and 2010. Furthermore, we performed sensitivity analyses including only nulliparae and nurses as these represented the largest occupational group in the Danish regions. Effect modification by maternal age, BMI, smoking and SES were explored by adding interaction terms to the regression analyses.

Analyses were undertaken on pseudo-anonymised data at a remote platform at Statistics Denmark by SAS 9.4 software. Cox regressions were executed applying the PHREG procedure. A significance level of 0.05 was used.

## RESULTS

A total of 22 744 pregnant employees and 377 896 pregnancy weeks at risk were included in the final analyses. Baseline characteristics of the study population by exposure to night work are presented in [table 1](#). Nearly half (44%) of the participants were exposed to night work with a median of nine night shifts during pregnancy weeks 3–21. Only 124 employees worked fixed nights with no registered day or evening shifts. A total of 1 889 women (8.5%) had a miscarriage. The exposed group had fewer miscarriages with a higher median for GA and fewer previous miscarriages compared with the reference group. A higher proportion of women in the exposed group were nulliparae, nurses and physicians, and had higher SES compared with the reference group.

We found an increased short-term risk of miscarriage after pregnancy week 8 with an adjusted HR of 1.32 (95% CI 1.07 to 1.62) if the women had  $\geq 2$  night shifts the previous week ([table 2](#)). The adjusted HR of late clinical miscarriage (pregnancy week 13–22) was 1.28 (95% CI 0.70 to 2.34). Only 133 of the miscarriages (7%) were late clinical miscarriages.

Age modified the risk of miscarriage according to night work the previous week ( $P < 0.05$  for multiplicative interaction). Women in the age group 26–30 years had the highest risk of miscarriage after pregnancy week 8 per additional night shift the previous week (HR 1.23 (95% CI 1.11 to 1.37)). Neither SES, maternal BMI nor tobacco smoking modified the association between recent night work and the risk of miscarriage (online appendix table 1).

All the sensitivity analyses were consistent with results from the main analyses (online appendix table 2). Taking competing risk of induced abortions into account did not affect the results.

A cumulated effect of number of night shifts during pregnancy weeks 3–21 was found with adjusted HR for miscarriage of 1.15 (95% CI 1.02 to 1.29) per 10 night shifts corresponding to one night shift every second week. In the categorised data, a dose-dependent risk of miscarriage was observed with an adjusted HR of 2.62 (95% CI 1.30 to 5.29) among those with 26 or more night shifts during pregnancy weeks 3–21 (average of 35 night shifts, ranging from 26 to 79). However, this group had a risk time of only 4 246 pregnancy weeks and eight cases ([table 3](#)).

A total of 6 435 pregnant employees (28%) had consecutive night shifts, the most frequent being two consecutive night

**Table 2** Risk of miscarriage by having night work the previous week

	All miscarriages (pregnancy weeks 4–22)			Miscarriages pregnancy weeks 4–8			Miscarriages pregnancy weeks 9–22		
	Crude HR (95% CI)	Adjusted† HR (95% CI)	Cases*/risk time†	Crude HR (95% CI)	Adjusted† HR (95% CI)	Cases*/risk time†	Crude HR (95% CI)	Adjusted† HR (95% CI)	Cases*/risk time†
Continuous exposures‡	1.06 (1.00 to 1.11)	1.06 (1.01 to 1.12)	1 889/377 896	1.03 (0.96 to 1.11)	1.02 (0.95 to 1.10)	930/110 671	1.09 (1.01 to 1.17)	1.10 (1.03 to 1.19)	959/267 225
Categorical exposure									
No night shift	1	1	1 521/314 511	1	1	741/89 229	1	1	780/225 282
One night shift	1.02 (0.87 to 1.20)	1.00 (0.85 to 1.18)	167/30 822	1.06 (0.85 to 1.32)	1.05 (0.84 to 1.32)	90/9 978	0.99 (0.78 to 1.25)	0.91 (0.71 to 1.17)	77/20 844
Two+night shifts	1.15 (0.99 to 1.33)	1.18 (1.01 to 1.37)	201/32 563	1.06 (0.86 to 1.31)	1.06 (0.85 to 1.31)	99/11 464	1.24 (1.01 to 1.53)	1.32 (1.07 to 1.62)	102/21 099

\*Miscarriage.

†Pregnancy weeks.

‡Adjusted for maternal age, BMI and smoking in the beginning of pregnancy, parity, SES, former miscarriages.

§Mean effect of adding an additional night shift the previous week.



**Table 3** Risk of miscarriage pregnancy weeks 4–22 by cumulated night shifts during pregnancy

	Cases*	Risk time†	Crude HR (95% CI)	Adjusted‡ HR (95% CI)
Continuous exposure§				
Ten night shifts			1.13 (1.01 to 1.27)	1.15 (1.02 to 1.29)
Categorical exposure				
No night shifts	1 149	226 184	1	1
1–10 night shifts	646	113 058	1.03 (0.93 to 1.13)	1.05 (0.94 to 1.16)
11–20 night shifts	78	30 060	1.21 (0.96 to 1.53)	1.20 (0.94 to 1.53)
21–25 night shifts	8	4 348	1.59 (0.79 to 3.19)	1.70 (0.84 to 3.42)
26+night shifts	8	4 246	2.48 (1.23 to 5.00)	2.62 (1.30 to 5.29)

\*Miscarriages.

†Pregnancy weeks.

‡Adjusted for maternal age, BMI and smoking during pregnancy, parity, SES, former miscarriages.

§Effect per additional night shift during pregnancy week 3–21.

shifts. The risk of miscarriage increased for each additional number of consecutive night shifts per spell, however, very few women ( $n=1.163$ ) had  $\geq$ four consecutive night shifts. (online figure 2). Quick return after night shift was registered for 810 pregnant employees during weeks 3–21 with a median of one quick return. No association was found between quick returns and the risk of miscarriage (HR 1.02 (95% CI 0.85 to 1)).

## DISCUSSION

In our nationwide cohort of pregnant women, primarily employed at hospitals, we found an increased risk of miscarriage among women who had night work the previous week, and among women with cumulated numbers of night shifts. Two or more night shifts the previous week increased the risk of miscarriage after pregnancy week 8 with 32% compared with women who had not worked night shifts the previous week. The number of night shifts and number of consecutive night shifts during pregnancy weeks 3–21 showed a dose-dependent increased risk. We found no association between quick returns after a night shift and the risk of miscarriage, but due to the power constraints these results should be interpreted with caution.

### Strengths and limitations of the study

To the best of our knowledge, our study represents the first to use prospective administrative data, which eliminates the risk of recall bias which is a common limitation in previous studies.<sup>12</sup> Furthermore, detailed payroll data accounted for sick leave, which is common among pregnant women,<sup>22,23</sup> and night work intensity.

However, some limitations need to be addressed. While all births in Denmark are registered in the DMBR, only miscarriages treated at hospitals are registered in the DNPR. We lacked information on very early miscarriages, which may be unnoticed by the women or handled in primary care. However, this is a premise in register-based studies and might attenuate the risk estimates if exposures are assumed to be especially harmful in the first weeks of gestation. A Danish pregnancy-planner study using hCG analysis found that 12.4% of conceived pregnancies ended as clinically recognised miscarriages.<sup>24</sup> Reasons for the lower proportion of miscarriages found in our study could be that our population was healthier and had less focus on pregnancy compared with the women in the pregnancy-planner study. However, it is more likely a reflection of organisational changes. In Denmark fewer miscarriages are being evacuated<sup>25</sup> and thus, a higher proportion of women may be treated by a primary care specialist. This may also partly explain the substantial decline in the proportion of registered miscarriages after

2010 relative to births, which is unlikely explained by biological causes. Nonetheless, place of treatment is likely independent of exposure and any potential misclassification would be non-differential with less risk of bias. This is supported by our sensitivity analysis restricted to pregnancies registered between 2007 and 2010, which showed consistent results.

The difference in distribution of SES, occupations, parity and number of previous miscarriages between employees working night shifts and employees never working nights could potentially confound the results in the analyses. We adjusted for SES and parity and our sensitivity analyses including only nurses or nulliparae, respectively, were consistent with the results in the primary analyses. It is disputed whether to adjust for previous miscarriages or not. If previous miscarriages are caused by the exposure of interest, risk estimates might erroneously be attenuated. If previous miscarriages are due to other risk factors with an unbalanced distribution across exposure categories adjustment is needed.<sup>9</sup> However, the risk estimates did not change substantially whether adjustment was performed or not. We also observed a difference between employees having night shifts and employees not working nights regarding number and time of miscarriage. This could be explained by delayed entry in the exposed group (only women with no abortions before the first registered night shift were included) causing survivor bias. In the Cox analyses this was accounted for.

Also, we were unable to account for other work-related exposures such as lifting and non-sitting work posture, which may increase the risk of miscarriage according to some studies.<sup>12,26,27</sup> Our sensitivity analysis only including nurses supported the primary results, but it has been shown in the American Nurses' Health Study cohort that nurses working day shifts have less strenuous work (lifting and standing) compared with nurses working fixed nights or shiftwork including night shifts.<sup>28,29</sup>

In Denmark during the study's time period about 8% of all pregnancies were conceived after fertility treatment. Being in fertility treatment could be a potential confounder due to increased risk of miscarriage and possible changed attitude towards working nights. The same could be the case for women with previous miscarriages. Unfortunately, we had no data on fertility treatment or cancelled night shifts. Because of this potential healthy worker selection our results could underestimate the effect of night work on miscarriage.

We found a stronger association between night work and the risk of miscarriages after pregnancy week 8. This may be explained by the decline in the proportion of chromosomally

abnormal fetuses with GA, which makes an association with environmental exposure more easily detectable among later miscarriages.<sup>30</sup> The association between night shifts and late clinical miscarriage (after pregnancy week 12) was less strong, but with a wide CI because of few cases.

### Findings in relation to other studies

Our findings confirm results in previous studies on fixed night work and risk of miscarriage.<sup>29 31–37</sup> However, studies on shiftwork including night shifts and risk of miscarriage have been inconsistent and have lacked information on number of consecutive shifts.<sup>29 31 32 34 36 38</sup> To date, only three previous studies have been based on prospectively collected data.<sup>32 37 38</sup> An American study, with information on exposure retrieved from interviews before pregnancy week 13, found no effect of working evening/night, but non-significant increased odds of miscarriage if working rotating shifts (OR=1.34 (95% CI 0.77 to 2.34)). The extent to which shiftwork included night shifts was not indicated.<sup>38</sup> In two studies based on the Danish National Birth Cohort (DNBC) night work was measured by asking the women whether they primarily worked 'fixed nights' or 'shiftwork including night shifts'. Both studies reported an increased risk of miscarriage among women who worked fixed nights with corresponding risk estimates of HR 1.27 (95% CI 0.89 to 1.82)<sup>37</sup> and HR 1.81 (95% CI 0.88 to 3.72)<sup>32</sup> respectively. For shiftwork including night shifts the HR was 1.21 (95% CI 1.06 to 1.39)<sup>37</sup> and 1.10 (95% CI 0.78 to 1.57),<sup>32</sup> respectively. The crude assessment of exposure in the earlier studies could result in misclassification and bias towards the null, especially in the group who had shiftwork. However, it is noteworthy that the pregnant women were included in the DNBC in pregnancy weeks 11–25 (median 16)<sup>37</sup> and thus primarily addressed late miscarriages.<sup>32</sup> In our study we only observed a few late miscarriages. The stronger association between fixed night work and miscarriages could be explained by the intensity of night shifts, including a higher number of cumulated and consecutive night shifts, with a higher risk of circadian disruption and decrease in melatonin levels. This is consistent with our results which showed a dose-related effect of the cumulated number of night shifts.

Although our population was based on a nationwide cohort, it primarily consisted of women working in public hospitals, who may have more health-promoting behaviour compared with the general Danish population. This was indicated in our data showing a lower prevalence of smoking in early pregnancy<sup>39</sup> and a lower proportion of obese women.<sup>40</sup> However, we found no modifying effect of BMI and smoking.

### CONCLUSION

The study corroborates earlier findings that night work during pregnancy may confer an increased risk of miscarriage and it indicates a lowest observed threshold level of two night shifts per week.

The new knowledge has relevance for working pregnant women as well as their employers, physicians and midwives. Moreover, the results could have implications for national occupational health regulations.

**Acknowledgements** The authors thank Clara Helene Glazer for proof-reading the final manuscript.

**Contributors** LMB, JPB, PECH, EMF and IOS conceived and designed the study. AHG, JH, ÅMH, HAK established and provided data from the DWHD. JPB analysed the data and EMF gave statistical support. LMB drafted the manuscript and all authors interpreted the data and revised the manuscript.

**Funding** This work was supported by the Danish Working Environment Research Fund grant 31-2015-03 2015001705.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** The study was approved by the Danish Data Protection Agency (though the notification system in the Capital region of Denmark, j.nr.: 2012-58-0004). By Danish law, no informed consent is required for a register-based study.

**Provenance and peer review** Not commissioned; externally peer reviewed.

### REFERENCES

- 1 Eurofond. Sixth European Working Conditions Survey – Overview report (Eurofond 2016): publications office of the European Union. 2016.
- 2 Daugaard S, Garde AH, Bonde JPE, et al. Night work, light exposure and melatonin on work days and days off. *Chronobiol Int* 2017;34:942–55.
- 3 Gómez-Acebo I, Dierssen-Sotos T, Papantoniou K, et al. Association between exposure to rotating night shift versus day shift using levels of 6-sulphatoxymelatonin and cortisol and other sex hormones in women. *Chronobiol Int* 2015;32:128–35.
- 4 Stevens RG, Hansen J, Costa G, et al. Considerations of circadian impact for defining 'shift work' in cancer studies: IARC Working Group Report. *Occup Environ Med* 2011;68:154–62.
- 5 Reiter RJ, Tan DX, Korkmaz A, et al. Melatonin and stable circadian rhythms optimize maternal, placental and fetal physiology. *Hum Reprod Update* 2014;20:293–307.
- 6 Kennaway DJ, Boden MJ, Varcoe TJ. Circadian rhythms and fertility. *Mol Cell Endocrinol* 2012;349:56–61.
- 7 Summa KC, Vitaterna MH, Turek FW. Environmental perturbation of the circadian clock disrupts pregnancy in the mouse. *PLoS One* 2012;7:e37668.
- 8 Boden MJ, Varcoe TJ, Kennaway DJ. Circadian regulation of reproduction: from gamete to offspring. *Prog Biophys Mol Biol* 2013;113:387–97.
- 9 Wilcox A. *Fertility and Pregnancy. an epidemiologic perspective*. Oxford: Oxford University Press, 2010.
- 10 Simpson JL. Causes of fetal wastage. *Clin Obstet Gynecol* 2007;50:10–30.
- 11 Stocker LJ, Macklon NS, Cheong YC, et al. Influence of shift work on early reproductive outcomes: a systematic review and meta-analysis. *Obstet Gynecol* 2014;124:99–110.
- 12 Bonde JP, Jørgensen KT, Bonzini M, et al. Miscarriage and occupational activity: a systematic review and meta-analysis regarding shift work, working hours, lifting, standing, and physical workload. *Scand J Work Environ Health* 2013;39:325–34.
- 13 Garde AH, Hansen J, Kolstad HA, et al. Payroll data based description of working hours in the Danish regions. *Chronobiol Int* 2018;35:795–800.
- 14 Bliddal M, Broe A, Pottegård A, et al. The Danish Medical Birth Register. *Eur J Epidemiol* 2018;33:27–36.
- 15 Schmidt M, Schmidt SA, Sandegaard JL, et al. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol* 2015;7:449–90.
- 16 Statistics Denmark DISCO-88. *The classification of occupations of Statistics Denmark*: Statistics Denmark, 1996.
- 17 Statistics Denmark DISCO-08. *The classification of occupations of Statistics Denmark*: Statistics Denmark, 2011.
- 18 Agenor A, Bhattacharya S. Infertility and miscarriage: common pathways in manifestation and management. *Womens Health* 2015;11:527–41.
- 19 Jørgensen T, Mortensen LH, Andersen AM. Social inequality in fetal and perinatal mortality in the Nordic countries. *Scand J Public Health* 2008;36:635–49.
- 20 Andersen PK, Geskus RB, de Witte T, et al. Competing risks in epidemiology: possibilities and pitfalls. *Int J Epidemiol* 2012;41:861–70.
- 21 Fine JP, Gray RJ. A Proportional hazards model for the redistribution of a competing risk. *J Am Stat Assoc* 1999;94:496–509.
- 22 Hansen ML, Thulstrup AM, Juhl M, et al. Occupational exposures and sick leave during pregnancy: results from a Danish cohort study. *Scand J Work Environ Health* 2015;41:397–406.
- 23 Ariansen AM. Age, occupational class and sickness absence during pregnancy: a retrospective analysis study of the Norwegian population registry. *BMJ Open* 2014;4:e004381.
- 24 Hjøllund NH, Bonde JP, Jensen TK, et al. Male-mediated spontaneous abortion among spouses of stainless steel welders. *Scand J Work Environ Health* 2000;26:187–92.
- 25 Lidsgaard O. Early pregnancy and miscarriage. Annual report from year 2013. [Tidlig graviditet og abort]: the Danish National Clinical Database on early pregnancies and miscarriages. 2014.
- 26 Mocevic E, Svendsen SW, Jørgensen KT, et al. Occupational lifting, fetal death and preterm birth: findings from the Danish National Birth Cohort using a job exposure matrix. *PLoS One* 2014;9:e90550.
- 27 Juhl M, Strandberg-Larsen K, Larsen PS, et al. Occupational lifting during pregnancy and risk of fetal death in a large national cohort study. *Scand J Work Environ Health* 2013;39:335–42.
- 28 Lawson CC, Johnson CY, Chavarro JE, et al. Work schedule and physically demanding work in relation to menstrual function: the Nurses' Health Study 3. *Scand J Work Environ Health* 2015;41:194–203.
- 29 Whelan EA, Lawson CC, Grajewski B, et al. Work schedule during pregnancy and spontaneous abortion. *Epidemiology* 2007;18:350–5.

- 30 Burgoyne PS, Holland K, Stephens R. Incidence of numerical chromosome anomalies in human pregnancy estimation from induced and spontaneous abortion data. *Hum Reprod* 1991;6:555–65.
- 31 Axelsson G, Ahlborg G, Bodin L. Shift work, nitrous oxide exposure, and spontaneous abortion among Swedish midwives. *Occup Environ Med* 1996;53:374–8.
- 32 Zhu JL, Hjøllund NH, Andersen AM, et al. Shift work, job stress, and late fetal loss: the National Birth Cohort in Denmark. *J Occup Environ Med* 2004;46:1144–9.
- 33 Swan SH, Beaumont JJ, Hammond SK, et al. Historical cohort study of spontaneous abortion among fabrication workers in the Semiconductor Health Study: agent-level analysis. *Am J Ind Med* 1995;28:751–69.
- 34 Hemminki K, Kyyrönen P, Lindbohm ML. Spontaneous abortions and malformations in the offspring of nurses exposed to anaesthetic gases, cytostatic drugs, and other potential hazards in hospitals, based on registered information of outcome. *J Epidemiol Community Health* 1985;39:141–7.
- 35 Infante-Rivard C, David M, Gauthier R, et al. Pregnancy loss and work schedule during pregnancy. *Epidemiology* 1993;4:73–5.
- 36 Axelsson G, Rylander R, Molin I. Outcome of pregnancy in relation to irregular and inconvenient work schedules. *Br J Ind Med* 1989;46:393–8.
- 37 Feodor Nilsson S, Andersen PK, Strandberg-Larsen K, et al. Risk factors for miscarriage from a prevention perspective: a nationwide follow-up study. *BJOG* 2014;121:1375–85.
- 38 Fenster L, Hubbard AE, Windham GC, et al. A prospective study of work-related physical exertion and spontaneous abortion. *Epidemiology* 1997;8:66–74.
- 39 Ekblad M, Gissler M, Korkeila J, et al. Trends and risk groups for smoking during pregnancy in Finland and other Nordic countries. *Eur J Public Health* 2014;24:544–51.
- 40 Christensen AI, Ekholm O, Davidsen E, et al. *Health and morbidity in Denmark 2010 and development since 1987. [Sundhed og sygelighed i Danmark 2010 og udviklingen siden 1987]*: National Institute of Public Health, 2012.