02E.4 EVALUATION OF EXPOSURE ASSESSMENT METHODS IN EPIDEMIOLOGICAL STUDIES: THE WELDING EXAMPLE

^{1,2}Susan Peters*, ³Jerome Lavoue, ⁴Marissa Baker, ¹Hans Kromhout. ¹*Institute for Risk Assessment Sciences, Utrecht University, Utrecht, Netherlands*; ²*University Medical Centre Utrecht, Utrecht, Netherlands*; ³*University of Montreal, Montreal, Canada*; ⁴*University of Washington, Seattle, USA*

10.1136/OEM-2019-EPI.55

Exposure assessment quality is a fundamental consideration in the design and evaluation of observational studies. High quality exposure assessment is particularly relevant for outcomes with long latency, such as cancer, where detailed information on past exposures are often missing and must therefore be estimated.

For the IARC Monograph on welding, the exposure group provided an overview of assessment methods used in the key epidemiological studies. Strengths and weaknesses of each study were assessed, along with their potential effects on interpretation of risk estimates.

For the association between lung cancer and welding fume exposure, 9 cohort and 10 case-control studies were reviewed. For ocular melanoma and ultraviolet radiation (UVR) from welding, 7 case-control studies were reviewed. Quality criteria were: full occupational histories, and standardized, blinded and quantitative exposure assessment. Additional criteria for lung cancer: specifically assessing welding fumes and using information on welding tasks. For ocular melanoma: assessing artificial and solar radiation separately, taking into account eye burns, eye protection and welding type.

Exposure assessment of welding fumes by applying a 'welding-exposure matrix' (n=2) or welding-specific questionnaires (n=3) were considered highest quality, followed by case-by-case expert assessment (n=5) or general job-exposure matrices (JEMs, n=4). Job title alone was considered less informative (n=5). For exposure to UVR, JEMs were most informative (n=2), followed by self-reported eye burns and self-reported exposure from specific welding types (n=2), although caution is advised regarding recall bias. Assessing welding fume exposure or ever exposure to welding arcs as proxy for UVR was considered less informative. For both exposures, ever versus never welder, or assessments based on data collected from proxies, were considered least informative.

The overall evaluation was that there is sufficient evidence in humans for the carcinogenicity of welding fumes and ultraviolet radiation from welding.

02E.5 CHALLENGES IN PESTICIDE EXPOSURE ASSESSMENT FOR EPIDEMIOLOGIC STUDIES

Laura Beane Freeman*. *Occupational and Environmental Epidemiology Branch, US National Cancer Institute, Rockville, USA*

10.1136/OEM-2019-EPI.56

Pesticides are used extensively worldwide to control weeds, insects and other pests, some of which have been linked to numerous adverse health outcomes, including cancer. Their ubiquity makes it imperative to conduct high quality epidemiologic studies of these chemicals for hazard identification

and risk assessment. There are hundreds of pesticides and pesticide active ingredients registered for use worldwide. However, only a handful have been formally evaluated for carcinogenicity, partially due to the paucity of high quality studies with information on specific active ingredients. Some of the challenges related to assessing exposure to these chemicals are not unique, but there are some particular circumstances that make evaluation of pesticides particularly challenging. Although agricultural practices are changing in many parts of the world, pesticides have traditionally been applied on family farms or small workplaces with few employees. Many studies have relied on the self-report of pesticides; however, depending on the specific work practices, the workers applying the pesticides may not know what chemicals are being applied and may not be able to report the specific active ingredients directly. Other studies have inferred exposure to individual chemicals using a variety of methods, including individual reports of general pesticide application, the crops being grown and combining these data with auxiliary information, such as sales or registration records, application diaries or the use of local experts to assign probability of exposure. In addition to the specific active ingredient, the intensity of exposure is related to other factors, including those associated with application methods and the use of personal protective equipment. Pesticides are often applied in combination with one another, or in mixtures and applicators may apply or be exposed to multiple chemicals through a growing season. Despite these, and other challenges, high quality exposure assessment is critical to studies of these important chemicals.

02E.6 DEFINING SHIFT WORK EXPOSURE IN EPIDEMIOLOGICAL STUDIES OF CANCER: WHERE HAVE WE BEEN, AND WHAT COMES NEXT?

Amy Hall*. *International Agency for Research on Cancer, Lyon, France*

10.1136/OEM-2019-EPI.57

In the epidemiological context, 'exposure' to shift work encompasses a variety of social, workplace, and individual characteristics. This complexity lends itself to coarse exposure assessment and assignment, which increases misclassification error and often reduces contrast between comparison groups. In early epidemiological studies of shift work, broad exposure assignment, such as 'day' versus 'night' worker, was commonly used; a technique that ignored various characteristics (e.g., shift intensity, timing, and work history) with potential impacts on long-term health risks.

In 2007, The International Agency for Research on Cancer classified 'shift work that involves circadian disruption' as a probable human carcinogen (Group 2A), on the basis of limited human evidence and sufficient evidence in experimental animals. In this evaluation, a major limitation of the epidemiological studies reviewed was the definition of shift work. A subsequent workshop convened by IARC considered how shift work should be assessed and quantified to improve the validity of future shift work and cancer studies. Since the publication of this working group's recommendations in 2011, the role of strong exposure assessment as a vital component of informative epidemiological studies of shift work has increasingly been recognized.

As part of a proposed symposium on the importance of exposure assessment in epidemiology for hazard identification and risk characterization, this presentation will summarize progress to date in the adoption of stronger exposure assessment methodologies in epidemiological studies of shift work, and offer ideas for next steps. A particular focus will be placed on the refinement of measurement techniques to assess long-term exposure to shift work (e.g., roster-based assessment and validity studies), and on quantitative considerations of important co-exposures (e.g., light-at-night).

Occupational Respiratory Disease

03A.1 ASSESSING THE UNDER-ESTIMATION OF OCCUPATIONAL RESPIRATORY DISEASES IN TAIWAN: ANALYSES OF DISEASE BURDENS AND HEALTHCARE COSTS

¹Chung-Yen Chen*, ²Yawen Cheng. ¹Taipei City Hospital, Renai Branch, Taipei City, Taiwan; ²Institute of Health Policy and Management, National Taiwan University, Taipei City, Taiwan

10.1136/OEM-2019-EPI.58

Background Respiratory diseases, including pneumoconiosis, asthma, chronic obstructive pulmonary disease (COPD) and respiratory malignancies, are featured by their multi-etiological nature and long latency periods, adding to the difficulties in recognizing their work-relatedness. Due to their heavy overall disease burdens and high healthcare spending, examining the fraction of respiratory diseases attributable to occupational factors is helpful in understanding the magnitude of under-estimation of occupational injuries and diseases.

Methods Prevalence rates of various occupational exposures were assessed retrospectively with self-reported surveys or job-exposure matrices. Relative risks of these exposures were drawn from international epidemiological literatures. The above two parameters were used to calculate population attributable risk percentages (PAR%), and with that the numbers of visiting and amounts of payment attributable to occupational factors were estimated based on the claim data of the National Health Insurance (NHI). The estimates were compared with the data of actual payment by the Workers' Compensation Insurance (WCI).

Results We estimated that around 15 000 patients visited medical facilities for occupational respiratory diseases in 2015, costing a total of 10 million USD. In contrast, less than 200 cases were approved by WCI in the same year. A 100-fold gap between the estimated and actual payment was also noted. Estimation analyses further indicated that 9.6% of cases and 48.1% of healthcare costs were asbestos-related.

Conclusion For occupational respiratory diseases with long latency periods and great causal complexity, the scope of under-estimation was substantial, and their medical expenses had been largely paid by NHI rather than WCI. To increase the visibility of occupational respiratory diseases, workplace exposure assessment and disease surveillance should be improved and public awareness of occupational diseases should be raised.

03A.3 DO AIRWAY INFLAMMATION AND AIRWAY RESPONSIVENESS MARKERS AT START OF APPRENTICESHIP PREDICT THEIR EVOLUTION DURING INITIAL TRAINING? A LONGITUDINAL STUDY

¹Valérie Demange*, ²Denis Zmirou-Navier, ³Abraham Bohadana, ¹Pascal Wild. ¹Institut National de Recherche et de Sécurité, Vandoeuvre-les-nancy, France; ²Ecole des Hautes Etudes de Santé Publique, Rennes, France; ³Pulmonary Institute, Jérusalem, Israël

10.1136/OEM-2019-EPI.59

The natural history of airway inflammation and symptoms in occupations at risk of asthma is still not fully understood. We aimed to study the evolution during apprenticeship of inflammation markers, bronchial hyperresponsiveness (BHR) and symptoms in at-risk subgroups as defined from measurements of markers made shortly after the start of training.

Respiratory symptoms, FEV₁ and airway resistance post-bronchial challenge (MBC) test results, fractional exhaled nitric oxide (FeNO) measurements, and eosinophils in nasal lavage fluid were longitudinally investigated in apprentice bakers, pastry-makers and hairdressers. Four baseline risk groups were defined, based on, (i) a high level of FeNO (NO), (ii) eosinophils > 1% (Eosino), (iii) a ≥ 15% decrease in FEV₁ during the MBC test (HR), and (iv) a ≥ 50% increase in the resistance (Resist). The statistical analysis relied on mixed models.

At baseline, the inflammation markers were related to the MBC markers. There was no evidence to suggest that the baseline risk groups predict a differential evolution of the airway inflammation and bronchial responsiveness markers, or the asthma-like symptoms considered. The baseline risk groups defined from MBC test predicted the levels of MBC markers. Similarly, the baseline risk groups based on eosinophilic inflammation predicted the levels of markers for eosinophilia. Twelve possible new asthma cases were identified, only the HR risk group predicted their occurrence.

Among this young population, at-risk groups based on initial high levels of inflammation markers did not experience any worsening during the follow-up. Initial BHR predicted consistently high levels of all markers considered and occurrence of possible asthma. These results have been published in BMC Pulmonary Medicine (Demange *et al.* *BMC Pulm Med.* 2018 July 11;18(1):113).

03A.4 ABSTRACT WITHDRAWN

03A.5 ENVIRONMENTAL DUST EXPOSURE FROM GOLD MINE WASTE DUMPS AND RESPIRATORY HEALTH EFFECTS IN JOHANNESBURG, SOUTH AFRICA

^{1,2}Samantha Iyaloo*, ^{1,2}Tahira Kootbodien, ^{1,2,3}Nisha Naicker, ^{2,3,4}Angela Mathee, ^{1,2}Spo Kgalamono, ^{1,2}Kerry Wilson, ^{1,2}David Rees. ¹National Institute for Occupational Health (NIOH), National Health Laboratory Services (NHLS), Johannesburg, South Africa; ²School of Public Health, University of the Witwatersrand, Johannesburg, South Africa; ³Environmental Health Department, Faculty of Health Sciences, University of Johannesburg, Johannesburg, South Africa; ⁴Environment and Health Research Unit, South African Medical Research Council, Johannesburg, South Africa

10.1136/OEM-2019-EPI.60

Background Thousands of people living in close proximity to gold mine waste dumps (GMWDs), are exposed to wind-