defined PTSD IRR 3.3 (95% CI 1.8–6.0), major depression 2.1 (1.3–3.5) and alcohol disorder 1.8 (1.0–3.3) at Wave 1. The adjusted IRR of incident CIDI-defined PTSD 3.7 (2.2–6.3) major depression 2.0 (1.2–3.2) and AUD 2.1 (1.1–3.8) at Wave 2 were increased for those with, compared to those without, MSI at Wave 1.

Conclusions Longitudinally, psychological disorders, in particular PTSD and depression, were found to be a risk factor for the development of MSI and the presence of MSI was a risk factor for the development of psychological disorders, in particular PTSD and depression. These findings have important implications for clinical and occupational health practice, service provision and longer term health in military and veteran populations.

05C.3 THE MENTAL HEALTH IMPACT OF DEPLOYMENT TO IRAQ AND AFGHANISTAN: WHAT DOES THE CURRENT DATA SHOW?

¹Sharon Stevelink*, ¹Margaret Jones, ¹Lisa Hull, ¹David Pernet, ¹Shirlee MacCrimmon, ²Laura Goodwin, ¹Deirdre MacManus, ^{1,3}Dominic Murphy, ⁴Norman Jones, ¹Neil Greenberg, ¹Roberto J Rona, ^{1,4}Nicola T Fear, ¹Sir Simon Wessely. ¹King's College London, King's Centre for Military Health Research, London, UK; ²University of Liverpool, Liverpool, UK; ³Combat Stress, London, UK; ⁴Academic Department for Military Mental Health, London, UK

10.1136/OEM-2019-EPI.124

The longer term mental health outcomes of UK serving and ex-serving personnel have been the subject of much speculation. The latest research findings from the third phase of a longitudinal study will be presented, which is the only military cohort study of its kind in the UK. Data was collected among 8093 personnel between 2014 and 2016. The study examined the prevalence of mental disorders and alcohol misuse, whether rates differed between serving and ex-serving regular personnel and the impact of various deployment exposures. The prevalence of probable posttraumatic stress disorder was 6.2% (95% confidence interval (CI) 5.5–6.9), 21.9% (95% CI 20.8–23.0) for common mental disorders and 10.0% (95% CI 9.2–10.9) for alcohol misuse. Deployment to Iraq or Afghanistan and self-reported role during deployment were associated with significantly worse mental health outcomes and alcohol misuse in ex-serving regular personnel but not in currently serving regular personnel. There was no association with number of deployments for any outcome. The findings highlight the importance of the continued monitoring of following personnel throughout their military career and beyond. The implications of the study outcomes will be discussed in the light of the mental health provision for serving and ex-serving personnel.

05C.4 THE ARMED SERVICES TRAUMA REHABILITATION OUTCOME STUDY (ADVANCE): A COHORT STUDY OF PHYSICAL BATTLEFIELD TRAUMA CASUALTIES

¹Nicola Fear*, ¹Daniel Dyball, ²Melanie Chesnokov, ³Christopher Boos, ²Anthony Bull, ²Paul Cullinan, ⁴Group Captain Alex Bennett. ¹King's College London, London, UK; ²Imperial College London, London, UK; ³Bournemouth University, Bournemouth, UK; ⁴Ministry of Defence, Loughborough, UK

10.1136/OEM-2019-EPI.125

The recent Iraq and Afghanistan conflicts represent a unique time in UK battlefield medicine; with more service personnel surviving complex traumas than any previous time. During these recent conflicts, approximately 830 UK service personnel were very seriously or seriously injured, and over 9000 were aero-medically evacuated. Little is known about the long-term health of these injured personnel. The Armed Services Trauma Rehabilitation Outcome Study (ADVANCE) is a longitudinal cohort study investigating the long-term medical and psychosocial outcomes of physical battlefield trauma casualties. 1200 male subjects who deployed to Afghanistan (600 injured and 600 non-injured individuals' frequency-matched by age, rank, service and role-in-theatre) will be followed up over a 20 year period. Primary objectives are to assess cardiovascular risk, cardiovascular events and osteoarthritis of the hip and knee. Secondary objectives are: musculoskeletal disease; pain; mental health; employment outcomes; functional status; quality of life; relationship status; sexual functioning; lung function; hearing loss and all-cause mortality. Data collection started in August 2015 and 705 participants have been recruited (376 non-injured and 329 injured). Most common injuries include single leg amputation, bilateral leg amputation (below knee), genital injuries and shrapnel injuries. Approximately 83% of participants are from the Army, 14% from Naval Services (including Royal Marines) and 3% from the Royal Air Force. The sample overall is comprised of approximately 14% Officers – with the other 86% holding lower ranks. Median age of the sample is 33 years at point of recruitment. Baseline data collection is due to be completed by the end of 2019, when follow data collection will commence. There are no other studies of this type currently being conducted and the results will have far reaching implications for the military and veteran community but also for members of the general population who have experienced a severe physical trauma.

05C.5 SARIN EXPOSURES IN A COHORT OF BRITISH MILITARY PARTICIPANTS IN HUMAN EXPERIMENTAL RESEARCH AT PORTON DOWN 1945–1987

¹Thomas J Keegan*, ²Lucy M Carpenter, ³Claire Brooks, ⁴Toby Langdon, ⁴Katherine M Venables. ¹Lancaster University, Lancaster, UK; ²Nuffield College, University of Oxford, Oxford, UK; ³Oxford Department of Oncology, Oxford, UK; ⁴Nuffield Department of Population Health, University of Oxford, Oxford, UK

10.1136/OEM-2019-EPI.126

Background The effects of exposure to chemical warfare agents in humans are topical. Porton Down, near Salisbury, is the UK's centre for research on chemical warfare. Since WWI, a programme of experiments involving approximately 30 000 participants drawn from the UK armed services has been undertaken there.

Objectives Our aim is to report on exposures to nerve agents, particularly sarin, using detailed exposure data in a cohort of servicemen who attended Porton Down.

Methods We have used existing data on exposures to UK servicemen who attended the human volunteer programme at Porton Down to examine exposures to nerve agents in general and to sarin in particular.

Results Six principal nerve agents were tested on humans, all between 1945 and 1987. Of the 4299 nerve agent tests recorded, 3511 (82%) were with sarin, most commonly in an exposure chamber, with inhalation being the commonest