incidence, trends and type of occupational disease (OD) and ii) incapacity for work due to ODs. It is hypothesized that workers in lower and higher SEP still differ in health disparities from an occupational perspective.

Methods From a Dutch cohort of occupational physicians (OPs), ODs assessed by OPs were retrieved over a seven year period (2010–2016) for lower and higher SEP groups. Incidence and type of OD and incapacity for work were determined for the total number of ODs and six frequently occurring ODs. Trends in incidence were estimated using a multilevel negative binominal regression model.

Result In total, 54 per 100,000 workers in elementary occupations, machine operating and assembly jobs, as well as managerial and intellectual jobs during 2016 had an OD diagnosed and reported by an OP, from which 98 per 100,000 were for lower SEP, and 36 per 100,000 for higher SEP. Among the lower SEP, musculoskeletal disorders (37%) and noise-induced hearing loss (NIHL) (32%) comprised two-thirds of the OD diagnoses. Among the higher SEP, distress/burnout comprised 60% of the OD diagnoses, with an increasing trend (6%; 95% CI: 3%–8%). Incapacity for work due to work-related low back disorders (69% vs 9%) and shoulder-, arm- and wrist disorders (89% vs 47%) differed significantly between workers in lower compared to higher SEP.

Discussion Occupational diseases occur at a 2.7 higher incidence rate for workers in lower SEP compared with higher SEP. Incapacity for work due to work-related musculoskeletal disorders are higher for workers in lower SEP compared with higher SEP, suggesting fewer opportunities to modify work tasks and working circumstances for lower SEP. Psychosocial risk factors constitute the greatest problem for workers in higher SEP, resulting in distress/burnout, accompanied by incapacity for work.

05E.4 BRIDGING THE GAP BETWEEN CLINICAL CASES AND EPIDEMIOLOGICAL EVIDENCE

¹Lode Godderis*, ¹Jelena Bakusic, ¹Sara Pauwels, ²Henk Van der Molen, ²Annet Lenderink. ¹KU Leuven, Leuven, Belgium; ²AMC/University of Amsterdam, Amsterdam, The Netherlands

10.1136/OEM-2019-EPI.135

One of the current gaps in the prevention of work-related diseases (WRDs) is the missing link between the assessment of occupational hazards introduced at the workplace, clinical alerts, epidemiological studies and policy actions. Alert systems aim to bridge this gap, by collecting information on diseases and exposures to raise alerts to different stakeholders and trigger timely prevention.

This project started with a review of international alert systems to identifying good practices and learning about prerequisites, drivers and obstacles to implement alert and sentinel systems. Next, the findings led to the implementation of an integrated approach consisting of alert systems on three levels in the Belgian context:

1. Signal detection and assessment: suspected clinical cases of new WRDs (new exposure–disease combinations) can be reported through an online platform called ‘SIGNAAL’. Each reported case is followed by an extensive assessment of

exposure– and work–relatedness performed by clinical experts. Since the start, 22 cases have been reported to the platform.

2. Signal strengthening through a network of occupational health physicians. This sentinel approach has been introduced through the PROBE system, in which 47 occupational physicians participated. During the periodic health examinations of workers, the physicians filled in a web survey regarding occupational exposure of a random sample of workers to 22 selected hazardous chemicals during the last working week. Results of the first study showed that 47% of workers were exposed to at least one chemical product from the list, with diesel exhaust being the most frequently reported substance (n=91; 14% of workers).
3. Alerts to public health authorities are mainly communicated through collaboration with Federal Public Services and Federal Agency for Occupational Risks.

These agencies provide support in the maintenance of systems and in turn, data from the systems is used as input for potential preventive strategies on company and societal level.

Occupational Health in Women

06A.1 AGGREGATION OF WORK-RELATED HEALTH PROBLEMS THROUGHOUT WORKING-LIFE IN A POPULATION-BASED SAMPLE OF WOMEN

¹Joana Amaro*, ^{1,2}Teresa Monjardino, ^{1,2}Raquel Lucas, ^{3,4,5}Mònica Ubalde-Lopez, ^{1,2}Henrique Barros. ¹EPIUnit – Instituto de Saúde Pública, Universidade do Porto, Rua das Taipas, nº 135, 4050–600 Porto, Porto, Portugal; ²Departamento de Ciências da Saúde Pública e Forenses e Educação Médica, Faculdade de Medicina, Universidade do Porto, Porto, Portugal; ³CISAL – Center for Research in Occupational Health, Pompeu Fabra University (UPF), Barcelona, Spain; ⁴CIBERESP, CIBER in Epidemiology and Public Health, Madrid, Spain; ⁵IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain

10.1136/OEM-2019-EPI.136

Introduction Multimorbidity potentiates a spectrum of adverse health outcomes that surpass those of individual diseases. However, little is known on disease aggregation related to occupational exposures.

Objective To assess the impact of life course occupational experience on health problem aggregation.

Methods We evaluated 4330 adult women at the 10-year-old follow-up wave of the population-based birth cohort Generation XXI, using a self-administered version of the Labour Force Survey item inquiring whether they had ever had a ‘physical or mental health problem that was caused or made worse by your current work or any previous work’ with 11 close-ended and one open-ended response options. Disease aggregation was assessed using principal components (PC) analysis and component scores were summarized by sociodemographic, anthropometric and work-related characteristics.

Results We identified five components to describe disease aggregation which accounted for 54.7% of observed variance. PC1 gathered all items on musculoskeletal disorders (back, upper and lower limb); participants with lower educational level, higher BMI, blue-collar jobs, working in the private sector, and with a history of occupational accidents scored higher in this component. PC2 gathered the item on mental disorders