worker's maximum BL, were found for brain cancer (malignant and benign combined), Hodgkins's lymphoma, lung cancer, and rectal cancer, while significant negative trends were found for colon cancer and melanoma. A borderline significant positive trend ($0.05 \leq p \leq 0.10$) was found for esophageal cancer. Significant interactions by country were found only for lung cancer, with Finland showing a strong positive trend and the UK showing only a modest trend. However, in general trends were marked in Finland and weak or inconsistent in the UK.

Conclusions We found strong positive incidence trends with increasing blood lead level, for several outcomes in internal analysis. Two of these, lung and brain cancer, were a priori suspected sites. Two of these outcomes are associated with smoking (lung and esophageal cancer), for which we had no data; however, we had no a priori reason to believe smoking differed between workers with different BL levels.

01C.5 ASSESSMENT AND ASSIGNMENT OF EXPOSURE TO ASBESTOS FOR AN INDUSTRIAL COHORT OF CHRYSOTILE MINERS AND PROCESSORS

¹Hans Kromhout*, ²Eleonora Feletto, ³Monika Moissonnier, ⁴Sara J Schonfeld, ³Ann Olsson, ^{5,6}Evgeny V Kovalevskiy, ^{5,6}Igor V Bukhtiyarov, ⁷Sergey V Kashanskiy, ³Joachim Schüz. ¹Utrecht University, Utrecht, Netherlands; ²Cancer Research Division, Cancer Council New South Wales, Woolloomooloo, Australia; ³Section of Environment and Radiation, International Agency for Research on Cancer, Lyon, France; ⁴Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, USA; ⁵Izmerov Research Institute of Occupational Health, Moscow, Russian Federation; ⁶SI.M. Sechenov First Moscow State Medical University (Sechenov University), Russian Federation; ⁷Yekaterinburg Medical Research Center for Prophylaxis and Health Protection in Industrial Workers, Yekaterinburg, Russian Federation

10.1136/OEM-2019-EPI.21

Introduction Historical dust concentrations are available for an occupational cohort study of workers active for 12 months or more between 1975 and 2010 in a chrysotile mine and processing factories in Asbest, Russian Federation. Their occupational histories were ascertained back to as early as the 1930s. A cohort specific job-exposure matrix (JEM) to estimate exposure to asbestos dust and fibre was elaborated.

Methods Almost 1 00 000 recorded dust concentrations were used to develop an asbestos dust JEM and previously derived conversion factors were applied to estimate an asbestos fibre JEM. Where dust concentrations were not available, linear mixed models were used to impute missing data. Both JEMs were applied to the occupational histories of over 30 000 individual workers (over 35% female workers) based on job title and year worked.

Results Assigned exposures varied over time with higher levels in the earlier years of activity. Approximately 97% of 2 00 000 person-years in the factories and 89% of 3 15 000 person-years in the mine had exposure assigned based on actual measurements. The median cumulative dust exposure for the exposed cohort was almost 50 mg/m³-years, with women slightly lower than men. The median cumulative fibre exposure for was 37 fibres/cm³-years for both men and women.

Discussion and conclusion A key strength of this study is the availability of high-quality measurement data covering workers' occupational histories. The dust and fibre JEMs enable estimation of annual profession-specific exposure levels that will form the basis of quantitative exposure estimates in the study and consequently quantitative exposure-response analyses.

01C.6 IS ADJUSTMENT FOR SMOKING NEEDED IN A COHORT STUDY OF CANCER MORTALITY AMONG CHRYSOTILE ASBESTOS FACTORY AND MINE WORKERS?

¹Ann Olsson*, ¹Madar Talibov, ²Evgeny V Kovalevskiy, ¹Graham Byrnes, ³Sara Schonfeld, ⁴Eleonora Feletto, ⁵Sergey V Kashanskiy, ¹Monika Moissonnier, ¹Evgenia Ostroumova, ¹Kurt Straif, ⁶Hans Kromhout, ²Igor V Bukhtiyarov, ¹Joachim Schüz. ¹International Agency For Research On Cancer, Lyon, France; ²Izmerov Research Institute of Occupational Health, Moscow, The Russian Federation; ³National Cancer Institute, Division of Cancer Epidemiology and Genetics, Radiation Epidemiology Branch, Bethesda, USA; ⁴Cancer Council NSW, Woolloomooloo, Australia; ⁵Yekaterinburg Medical Research Center for Prophylaxis and Health Protection in Industrial Workers, Yekaterinburg, The Russian Federation; ⁶Institute for Risk Assessment Sciences, Utrecht, The Netherlands

10.1136/OEM-2019-EPI.22

Introduction A retrospective cohort study of 35 840 employees is currently being conducted in a chrysotile mine and its processing facilities in Asbest, the Russian Federation. The primary aim is to quantify exposure-response relationships for cancers potentially associated with chrysotile exposure. Some of those cancers are also tobacco-related; however individual-level information on tobacco use is not available for the full cohort. To address this gap, a cross-sectional study of current and retired workers from JSC Uralasbest was conducted to assess the relationship between smoking status and workers' exposure to chrysotile.

Methods Self-administered questionnaires were completed by current workers during meetings organized by occupational safety specialists. Retired workers filled in questionnaires during Veterans' meetings or were interviewed via telephone or at home. Estimates of exposure to chrysotile were available for 999 current and 3795 retired workers who were linked to the cohort study.

Results Among the 7451 respondents (n=3698 men and n=3753 women), 66% of men and 9% of women were ever-smokers. Smoking prevalence was stable across birth decades in men, but increased from <10% in women born before 1960 to 30% in women born after 1980. Among ever-smokers, men smoked more cigarettes per day than women. The smoking prevalence was similar in exposed and non-exposed men, and did not increase with exposure category. The greatest difference in the proportion of smokers among women was observed between non-exposed (4%) and all exposed categories combined (7.5%), whereas there was little variation across categories of exposure (7%–8%). Overall, the self-reported smoking prevalence may have been underestimated, especially in the older age categories due to survivor bias.

Conclusion While no adjustments for smoking among men appear necessary in the future analyses, including smoking propensity by birth cohort for women may be useful.