Conclusion Health related absence contributes to significant loss in productive work-time. There is scope to reduce these losses by improvement in health situation. Our study provided key information and trends for planning evidence based health promotion programmes in the industry.

P.2.17 CASE REPORT: THREE OCCUPATIONAL DISEASES IN A NAIL TECHNICIAN

^{1,2}Shlomo Moshe*, ^{1,2}Ayala Krakov. ¹Maccabi Healthcare Services, Holon, Israel; ²Sackler Faculty of Medicine, School of Public Health, Tel-Aviv University, Tel Aviv, Israel

10.1136/OEM-2019-EPI.250

Introduction In cosmetics, acrylic glues are used to apply nail and eyelash extensions. Acrylates are highly irritant materials and may cause local irritation (Contact Dermatitis – CD), respiratory irritation (Asthma) and systemic irritation (Urticaria). This case report describes these three occupational diseases.

Clinical presentation A 48-year-old female, normally healthy, has been working in the nail beauty field for 11 years. Lately, following an increase of work, she started coughing every morning and feeling shortness of breath which was worsen at the start of work, and improved on vacations. In addition, the patient suffered from skin tenderness in her fingers. An examination of the hands showed a typical picture of CD in left hand fingers 4–5.

Methods and results Specific Patch Tests showed sensitivity to Hydroxyethyl Methacrylate and Hydroxypropyl Methacrylate. A methacholine challenge test showed a 22% decrease in FEV1 at 3.94µmol methacholine (i.e. positive test). The patient performed a Peak Expiratory Flow test (PEF). After a week in the nail salon her respiratory and dermatological status worsened greatly, including development of contact urticaria. She was instructed to stop immediately the exposure and start taking p/o steroids. Due to her 3 occupational diseases she was removed from work and latter was recognized as having 3 occupational disease by the Israeli National Insurance which supported her vocational rehabilitation.

Discussion Acrylic glues are hazardous agents which can cause several occupational diseases and require occupational environment monitoring yearly according to Israeli regulations. The TLV-TWA is 2ppm. Small private businesses like the patients' place don't perform environmental monitoring or use personal or environmental protective equipment which endanger its workers (usually low social class women) who are not aware of the dangers in this industry. More public attention should be given to the risk in this industry.

P.2.18 THE CURRICULUM IMPLEMENTATION OF OCCUPATIONAL MEDICINE IN ISRAEL

Shlomo Moshe*, ¹Maccabi Healthcare Services, Occupational Medicine Department, Holon, Israel; ²Sackler Faculty of Medicine, School of Public Health, Department of Environmental and Occupational Health, Tel Aviv University, Tel Aviv, Israel

10.1136/OEM-2019-EPI.251

Background At the early 1990s occupational medicine (OM) was recognized as a specialty in Israel. Fifteen years later after gathering problems and shortcomings of the first curriculum, the Israeli Association of occupational medicine (IAOM) defined the framework for a new curriculum in order to have a modern and better curriculum.

Methods In the early 2000 an expert committee appointed by the IAOM suggested a new syllabus, based on the scientific literature. The recommendations were sent to all experts in OM in Israel, and other professional members like industrial hygienist. We got several responses. Most people suggested additional standards.

Results The curriculum includes 54 months of training programme. The characteristics of the new curriculum are: interaction between theory and practice; fields of knowledge, learning process; competencies which are needed; and multi-disciplinarity. We added an obligatory walkthrough list of industries. Three of the walkthrough reports consists a part of the final oral examinations.

Conclusions The aim of the curriculum was to produce a comprehensive approach in OM competencies. The curriculum was adopted by the Israeli Medical Association and Ministry of Health since 2010.

P.2.19 EFFECT OF COLLABORATIVE SUPPORT OF OCCUPATIONAL AND CLINICAL PHYSICIANS - EVALUATION OF OUTCOME DUE TO TREATMENT DROP OUT IN NON-COMMUNICABLE DISEASES

^{1,2}Go Muto. ¹Harvard T.H.Chan School of Public Health, Boston, USA; ²Kitasato University, Dep of Hygiene, School of Medicine, Sagami-hara/Kanagawa, Japan

10.1136/OEM-2019-EPI.252

Objectives To develop a prediction model of outcome due to drop out in treatment for non-communicable diseases (NCDs) in Japan as well as to evaluate the effectiveness of collaboration between occupational and clinical physicians.

Methods Using both claim and health check-up data (JMDC) of beneficiaries aged 20 to 74 years who were covered by health insurance societies between 2008 and 2016, we identified high risk 5 33 955 examinees of health checkups who met the following criteria and had no claim data on the corresponding items 4 months prior to the health checkups. The criteria was either blood pressure, blood glucose level, or blood lipid level provided by Japanese government.

Results The cumulative non-visiting rates at 12 months after health check-ups were 84.4%. The rate of drop off during treatment was over 70%. As for HT, DM, and DL, the cumulative non-visiting rates at 12 months were 84.3, 67.9, and 86.1%, respectively. When focusing on the high risk group, the rate was 74.0%, since the rate for extremely high risk DM (HbA1c≥8.4%) was 51.9%.

Conclusion Over 80% high risk examinees of non-communicable diseases did not visit to medical institutions during 1 year period after health check-ups, although approximately half of the individuals with extremely high risk of DM visited to physicians at least one time. Finally, we developed a predictive scoring model for visiting to medical institutions as well as the risk of drop out from treatment, and it was well validated from both discrimination and calibration viewpoints. In addition, we evaluated the effectiveness of strategy for preventing treatment drop off through collaborative support from occupational and clinical physicians. This strategy enables public/occupational health staffs to individualize population for strong intervention and to support harmonization of treatment and work in NCDs.