

## LETTER

## How should data on airborne transmission of SARS-CoV-2 change occupational health guidelines?

Sir,

On the 6th of April, the WHO provided guidance about personal protective equipment (PPE) for healthcare workers (HCWs) in healthcare settings areas that have reported cases of COVID-19, stating that HCWs working in settings where aerosol-generating procedures were performed should wear a particulate respirator (N95 or FFP2), whereas in their previous recommendations, such masks were restricted to those who actually performed such procedures. This a step forward, although insufficient, towards the recognition of aerosol transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and its consequences in terms of PPE choice.

COVID-19 has been considered as a droplet transmitted disease by WHO, which means that it can be prevented using surgical masks and hand hygiene. However, there is now increasing evidence suggesting that SARS-CoV-2 may also be found in droplet nuclei, defined by WHO as respiratory droplets smaller than 5 µm, which can travel on long distances and remain in suspension in the air for a long time.<sup>1,2</sup>

SARS-CoV-2 RNA is easily detected in nasopharyngeal swabs during the presymptomatic stage of the disease and asymptomatic patients may shed droplet nuclei that contain the virus. Two studies performed in medical units where patients with COVID-19 were hospitalised have found SARS-CoV-2 on surfaces and in the air of patients' rooms, and even in the medical staff office, with peaks in the submicrometre size.<sup>2,3</sup> Although it is widely acknowledged that the identification of viral RNA does not necessarily imply the presence of an infectious virus, experimental data have demonstrated that SARS-CoV-2 can survive in an aerosol for up to 3 hours.<sup>4</sup>

Furthermore, investigations of COVID-19 clusters in various environments such as restaurants, ships or buses have concluded that direct contact transmission was insufficient to explain all cases and that airborne transmission was likely.

Indeed, the use of PPE and infection control training has been associated with decreased infection risk for HCWs.<sup>5</sup> However, droplet and contact precautions have not been sufficient in many cases. A recent study on anaesthesiologists performing spinal anaesthesia showed that the use of category 3 PPE (including self-contained breathing apparatus) as defined by the EU Regulation 2016/425 reduced the risk of contamination by 95% compared with category 1 PPE (including surgical mask), which are known to offer little protection against droplet nuclei.<sup>6</sup>

Therefore, it is more and more challenging to rule out the risk of airborne transmission of COVID-19. Given the severity of the disease that is concerning HCWs more than any other occupational group, it seems reasonable to adjust occupational health guidelines.

Systematic use of N95 or FFP2 respirators should be discussed when caring for a patient with COVID-19, as a piece of a broader strategy that must include education, fit-checking, frequent hand hygiene, respiratory etiquette, organisational factors and engineering controls (maximising ventilation and avoiding recirculation). As many clusters have started in professional settings such as factories and have spread up in the communities, protecting workers against COVID-19 is of utmost importance for the public health response to the pandemic.

Jean François Gehanno,<sup>1,2</sup>  
 Vincent Bonneterre,<sup>3,4</sup> Pascal Andujar,<sup>5,6,7</sup>  
 Jean-Claude Pairon,<sup>5,6,7</sup> Christophe Paris,<sup>8,9</sup>  
 Audrey Petit,<sup>9,10</sup> Catherine Verdun-Esquer,<sup>11</sup>  
 Alexis Descatha,<sup>9,12</sup>  
 Quentin V Durand-Moreau,<sup>13</sup>  
 Patrick Brochard<sup>14</sup>

<sup>1</sup>Department of occupational Medicine, Rouen University Hospital, Rouen, France

<sup>2</sup>Laboratoire d'informatique médicale et d'ingénierie des connaissances en e-santé, LIMICS, Sorbonne Université, Inserm, université Paris 13, Paris, France

<sup>3</sup>TIMC Research Laboratory, Grenoble Alpes University, La Tronche cedex, France

<sup>4</sup>Occupational & Environmental Diseases Centre, Grenoble cedex 09, France

<sup>5</sup>U955, Equipe 4, INSERM, Creteil, France

<sup>6</sup>Service de Pneumologie et de Pathologie Professionnelle, Centre Hospitalier Intercommunal de Créteil, Créteil, France

<sup>7</sup>Faculté de Santé, Université Paris-Est Créteil, Créteil, France

<sup>8</sup>Service de pathologie professionnelle et de l'environnement, CHU Rennes, Rennes, France

<sup>9</sup>UMR\_S1085 Ester team, UNIV Angers, CHU Angers, Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail), Angers, France

<sup>10</sup>Department of Occupational Health, University Hospital of Angers, Angers, France

<sup>11</sup>CHU de Bordeaux, Bordeaux, France

<sup>12</sup>CAPTIV CDC, CHU Angers Pôle A Vasculaire, Angers, France

<sup>13</sup>Division of Preventive Medicine, University of Alberta, Edmonton, Alberta, Canada

<sup>14</sup>Université Bordeaux, Bordeaux, France

**Correspondence to** Professor Alexis Descatha, U1085 (Irset) Ester, University of Angers, Angers 49000, France; Alexis.Descatha@inserm.fr

**Twitter** Quentin V Durand-Moreau @qdurandmoreau

**Contributors** All authors have participated in writing the letter and approved it.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; internally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.



**To cite** Gehanno JF, Bonneterre V, Andujar P, et al. *Occup Environ Med* 2020;**77**:736.

Received 15 May 2020

Revised 15 June 2020

Accepted 18 June 2020

Published Online First 30 June 2020

*Occup Environ Med* 2020;**77**:736.

doi:10.1136/oemed-2020-106707

### ORCID iDs

Vincent Bonneterre <http://orcid.org/0000-0003-2353-7102>

Alexis Descatha <http://orcid.org/0000-0001-6028-3186>

Quentin V Durand-Moreau <http://orcid.org/0000-0003-1168-4201>

### REFERENCES

- World Health Organization., Pandemic and Epidemic Diseases, World Health Organization. Infection prevention and control of epidemic- and pandemic-prone acute respiratory infections in health care: WHO guidelines, 2014. Available: [http://apps.who.int/iris/bitstream/10665/112656/1/9789241507134\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/112656/1/9789241507134_eng.pdf?ua=1) [Accessed 24 Apr 2020].
- Gehanno JF, Bonneterre V, Andujar P, et al. [Evidences for a possible airborne transmission of SARS-CoV-2 in the COVID-19 crisis]. *Arch des mal profs*. In Press 2020.
- Liu Y, Ning Z, Chen Y, et al. Aerodynamic analysis of SARS-CoV-2 in two Wuhan hospitals. *Nature* 2020;1–6.
- van Doremalen N, Bushmaker T, Munster VJ. Stability of middle East respiratory syndrome coronavirus (MERS-CoV) under different environmental conditions. *Euro Surveill* 2013;18. doi:10.2807/1560-7917.ES2013.18.38.20590. [Epub ahead of print: 19 Sep 2013].
- Lai X, Wang M, Qin C, et al. Coronavirus disease 2019 (COVID-2019) infection among health care workers and implications for prevention measures in a tertiary hospital in Wuhan, China. *JAMA Netw Open* 2020;3:e209666.
- Zhong Q, Liu YY, Luo Q, et al. Spinal anaesthesia for patients with coronavirus disease 2019 and possible transmission rates in anaesthetists: retrospective, single-centre, observational cohort study. *Br J Anaesth* 2020;124:670–5.